

Mary R. Suchyta
Callie J. Beck
Colin W. Key
Al Jephson
Ramona O. Hopkins

Substance dependence and psychiatric disorders are related to outcomes in a mixed ICU population

Published online: 16 December 2008
© Springer-Verlag 2008

The online version of the original article can be found under doi:[10.1007/s00134-008-1263-9](https://doi.org/10.1007/s00134-008-1263-9).

M. R. Suchyta · A. Jephson · R. O. Hopkins
Department of Medicine, Pulmonary
and Critical Care Division, LDS Hospital,
Salt Lake City, UT, USA
e-mail: mary.suchyta@mail.org

M. R. Suchyta · A. Jephson · R. O. Hopkins (✉)
Department of Pulmonary and Critical Care Medicine,
Intermountain Medical Center,
5121 South Cottonwood Street, Murray,
UT 84107, USA
e-mail: mona_hopkins@byu.edu
Tel.: +1-801-5074569
Fax: +1-801-4220602

C. J. Beck · C. W. Key · R. O. Hopkins
Psychology Department, 1001 SWKT,
Brigham Young University,
Provo, UT 84602, USA

R. O. Hopkins
Neuroscience Center, Brigham Young University,
Provo, UT, USA

Erratum to: Intensive Care Med (2008) 34:2264–2267 DOI 10.1007/s00134-008-1263-9

The legends to Figs. 1 and 2 were unfortunately transposed.

The figures are correct.

The corrected figures are given here.




Fig. 1 The prevalence of substance dependence and psychiatric disorders in critically ill patients (black bars) compared to Utah population data (gray bars). Significant differences are denoted with * for p values of <0.01 and ** for p values of <0.001 . Error bars denote the 95% confidence intervals for the percentages (proportions)




Fig. 2 The prevalence of psychiatric disorders by disorder in critically ill patients (black bars) compared to United States population data (gray bars). Significant differences are denoted with * for p values of <0.01 and ** for p values of <0.001 . Error bars denote the standard error of the percentages (proportions)