

Sean K. Gorman
Michelle Ho Chung
Richard S. Slavik
Peter J. Zed
Kerry Wilbur
Vinay K. Dhingra

Erratum to: A critical appraisal of the quality of critical care pharmacotherapy clinical practice guidelines and their strength of recommendations

Published online: 24 August 2010
© Copyright jointly held by Springer and ESICM 2010

The online version of the original article can be found under doi:[10.1007/s00134-010-1786-8](https://doi.org/10.1007/s00134-010-1786-8).

Electronic supplementary material The online version of this article (doi:[10.1007/s00134-010-1976-4](https://doi.org/10.1007/s00134-010-1976-4)) contains supplementary material, which is available to authorized users.

S. K. Gorman (✉) · P. J. Zed
Department of Pharmacy, Capital District Health Authority,
College of Pharmacy, Dalhousie University,
c/o Rm 2043 Victoria Building, 1276 South Park Street,
Halifax, NS B3H 2Y9, Canada
e-mail: sean.gorman@dal.ca
Tel.: +1-902-4736576
Fax: +1-902-4731606

M. H. Chung
Department of Pharmacy, St. Paul's Hospital,
1081 Burrard Street, Vancouver, BC V6Z 1Y6, Canada

R. S. Slavik
Pharmacy Department, Interior Health Authority,
Faculty of Pharmaceutical Sciences,
University of British Columbia, 200-1835 Gordon Drive,
Kelowna, BC V1Y 3H5, Canada

P. J. Zed
Department of Emergency Medicine, Dalhousie University,
Halifax Infirmary, Suite 355 (Room 345), 1796 Summer Street,
Halifax, NS B3H 3A7, Canada

K. Wilbur
College of Pharmacy, Qatar University,
P.O. Box 2713, Doha, Qatar

V. K. Dhingra
Intensive Care Unit, Vancouver General Hospital,
855 West 12th Avenue, Vancouver, BC V5Z 1M9, Canada

V. K. Dhingra
Faculty of Medicine, University of British Columbia,
Vancouver, BC V5Z 1M9, Canada

Erratum to: Intensive Care Med DOI:[10.1007/s00134-010-1786-8](https://doi.org/10.1007/s00134-010-1786-8)

Some data from Appendix 1 were lost during the editorial process. The corrected appendix is reproduced here. Additional electronic supplementary material has also been supplied.

Appendix 1 Clinical practice guidelines with AGREE recommendation, standardized domain scores, and breakdown of item scores

Guideline title	AGREE recommendation	Standardized domain scores ^a	Number of items scoring 1–2	Number of items scoring 3–4
Guidelines for the management of penetrating brain injury [9]	Not recommended	Domain 1: 67% Domain 2: 25% Domain 3: 61% Domain 4: 58% Domain 5: 3% Domain 6: 8%	12	11
Guidelines for the management of severe traumatic brain injury [10]	Strongly recommended	Domain 1: 72% Domain 2: 31% Domain 3: 93% Domain 4: 92% Domain 5: 17% Domain 6: 25%	7	16
Blunt cerebrovascular injury practice management guidelines [11]	Not recommended	Domain 1: 67% Domain 2: 17% Domain 3: 57% Domain 4: 60% Domain 5: 8% Domain 6: 0%	12	11
Guidelines for the management of aneurysmal subarachnoid hemorrhage [12]	Not recommended	Domain 1: 31% Domain 2: 6% Domain 3: 25% Domain 4: 48% Domain 5: 6% Domain 6: 0%	17	6
Guidelines for the management of spontaneous intracerebral hemorrhage [13]	Recommended with alterations	Domain 1: 33% Domain 2: 29% Domain 3: 58% Domain 4: 73% Domain 5: 3% Domain 6: 58%	11	12
Guideline on the management of status epilepticus [14]	Recommended with alterations	Domain 1: 44% Domain 2: 21% Domain 3: 63% Domain 4: 67% Domain 5: 0% Domain 6: 38%	11	12
Clinical practice guidelines for sustained use of sedatives and analgesics in the critically ill adult [15]	Recommended with alterations	Domain 1: 47% Domain 2: 38% Domain 3: 57% Domain 4: 88% Domain 5: 28% Domain 6: 46%	9	14
Clinical practice guidelines for sustained neuromuscular blockade in the adult critically ill patient [16]	Recommended with alterations	Domain 1: 50% Domain 2: 38% Domain 3: 61% Domain 4: 81% Domain 5: 25% Domain 6: 38%	8	15
Evidence-based clinical practice guideline for the prevention of ventilator-associated pneumonia [17]	Strongly recommended	Domain 1: 75% Domain 2: 65% Domain 3: 95% Domain 4: 79% Domain 5: 39% Domain 6: 96%	4	19
Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia [18]	Recommended with alterations	Domain 1: 64% Domain 2: 25% Domain 3: 39% Domain 4: 75% Domain 5: 22% Domain 6: 67%	12	11

Appendix 1 continued

Guideline title	AGREE recommendation	Standardized domain scores ^a	Number of items scoring 1–2	Number of items scoring 3–4
Consensus guidelines on the management of community-acquired pneumonia in adults [19]	Strongly recommended	Domain 1: 81% Domain 2: 44% Domain 3: 55% Domain 4: 90% Domain 5: 56% Domain 6: 71%	7	16
The American-European consensus conference on ARDS, part 2 [20]	Not recommended	Domain 1: 42% Domain 2: 19% Domain 3: 17% Domain 4: 35% Domain 5: 11% Domain 6: 0%	17	6
Hospital management of adults with severe acute respiratory syndrome (SARS) if SARS re-emerges [21]	Not recommended	Domain 1: 44% Domain 2: 17% Domain 3: 13% Domain 4: 40% Domain 5: 0% Domain 6: 0%	20	3
Inhaled nitric oxide therapy in adults: European expert recommendations [22]	Not recommended	Domain 1: 39% Domain 2: 19% Domain 3: 23% Domain 4: 40% Domain 5: 3% Domain 6: 83%	16	7
Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock [23]	Strongly recommended	Domain 1: 69% Domain 2: 42% Domain 3: 71% Domain 4: 92% Domain 5: 36% Domain 6: 100%	6	17
Practice parameters for hemodynamic support of sepsis in adult patients [24]	Not recommended	Domain 1: 61% Domain 2: 38% Domain 3: 35% Domain 4: 71% Domain 5: 6% Domain 6: 4%	14	9
Evidence-based colloid use in the critically ill [25]	Recommended with alterations	Domain 1: 53% Domain 2: 21% Domain 3: 51% Domain 4: 38% Domain 5: 8% Domain 6: 46%	13	10
Guidelines for cardiopulmonary resuscitation and emergency cardiovascular care [26]	Strongly recommended	Domain 1: 64% Domain 2: 38% Domain 3: 62% Domain 4: 77% Domain 5: 33% Domain 6: 67%	8	15
Guidelines for stress ulcer prophylaxis [27]	Strongly recommended	Domain 1: 75% Domain 2: 46% Domain 3: 73% Domain 4: 73% Domain 5: 39% Domain 6: 63%	6	17
Guidelines for the selection of anti-infective agents for complicated intra-abdominal infections [28]	Recommended with alterations	Domain 1: 89% Domain 2: 50% Domain 3: 50% Domain 4: 75% Domain 5: 42% Domain 6: 50%	8	15

Appendix 1 continued

Guideline title	AGREE recommendation	Standardized domain scores ^a	Number of items scoring 1–2	Number of items scoring 3–4
Guidelines on antimicrobial therapy for intra-abdominal infections [29]	Recommended with alterations	Domain 1: 89% Domain 2: 27% Domain 3: 79% Domain 4: 88% Domain 5: 0% Domain 6: 4%	9	14
Management of critically ill patients with severe acute pancreatitis [30]	Not recommended	Domain 1: 83% Domain 2: 29% Domain 3: 30% Domain 4: 67% Domain 5: 8% Domain 6: 8%	13	10
Guidelines for the prevention of intravascular catheter-related infections [31]	Recommended with alterations	Domain 1: 72% Domain 2: 50% Domain 3: 32% Domain 4: 71% Domain 5: 42% Domain 6: 54%	12	11
Guidelines for the management of intravascular catheter-related infections [32]	Not recommended	Domain 1: 67% Domain 2: 29% Domain 3: 29% Domain 4: 85% Domain 5: 25% Domain 6: 8%	12	11

^a AGREE Domain 1: Scope and purpose; AGREE Domain 2: Stakeholder involvement; AGREE Domain 3: Rigor of development; AGREE Domain 4: Clarity and presentation; AGREE Domain 5: Applicability; AGREE Domain 6: Editorial independence