

Jerry P. Nolan
Jasmeet Soar
Alain Cariou
Tobias Cronberg
Véronique R. M. Moulaert
Charles D. Deakin
Bernd W. Bottiger
Hans Friberg
Kjetil Sunde
Claudio Sandroni

# Erratum to: European Resuscitation Council and European Society of Intensive Care Medicine 2015 guidelines for post-resuscitation care

Published online: 9 December 2015

© Springer-Verlag Berlin Heidelberg and ESICM 2015

The online version of the original article can be found under doi:10.1007/s00134-015-4051-3.

### J. P. Nolan (💌)

Department of Anaesthesia and Intensive Care Medicine, Royal United Hospital, Bath, UK e-mail: jerry.nolan@nhs.net

### I P Nolan

School of Clinical Sciences, University of Bristol, Bristol, UK

### J. Soar

Anaesthesia and Intensive Care Medicine, Southmead Hospital, Bristol, UK

# A. Cariou

Cochin University Hospital (APHP) and Paris Descartes University, Paris, France

# T. Cronberg

Division of Neurology, Department of Clinical Sciences, Lund University, Lund, Sweden

# V. R. M. Moulaert

Adelante, Centre of Expertise in Rehabilitation and Audiology, Hoensbroek, The Netherlands

### C. D. Deakin

Cardiac Anaesthesia and Cardiac Intensive Care and NIHR Southampton Respiratory Biomedical Research Unit, University Hospital, Southampton, UK

### B. W. Bottiger

Department of Anaesthesiology and Intensive Care Medicine, University Hospital of Cologne, Cologne, Germany

### H. Friberg

Division of Anesthesia and Intensive Care Medicine, Department of Clinical Sciences, Lund University, Lund, Sweden

### K. Sunde

Division of Emergencies and Critical Care, Department of Anaesthesiology, Oslo University Hospital and Institute of Clinical Medicine, University of Oslo, Oslo, Norway

# C. Sandroni

Department of Anaesthesiology and Intensive Care, Catholic University School of Medicine, Rome, Italy

# Erratum to: Intensive Care Med (2015) 41:2039–2056 DOI 10.1007/s00134-015-4051-3

A mistake was made in Fig. 2: in the lower blue box on the right-hand side, poor outcome is reported as being "very likely", whereas in fact it is "likely". The corrected Fig. 2 is supplied here. The authors regret this error and apologize for any inconvenience.




Fig. 2 Prognostication strategy algorithm. *EEG* electroencephalography, *NSE* neuron specific enolase, *SSEP* somatosensory evoked potentials, *ROSC* return of spontaneous circulation, *FPR* false positive rate, *CI* confidence interval