RETRACTION NOTE

Retraction Note to: Volume therapy in the intensive care patient – we are still confused, but..

Editorial Office of Intensive Care Medicine*

© 2020 Springer-Verlag GmbH Germany, part of Springer Nature

**Retraction note to: Intensive Care Med (2000) 26:
1181–1192
<https://doi.org/10.1007/s001340000625>**

The Editor-in-Chief has retracted this article [1] because a number of studies included in this review [2, 3, 4] (originally cited as references 98, 128, 129) have since been retracted. Consequently, this has rendered the content of the review unreliable.

Author Joachim Boldt has not responded to any correspondence from the publisher about this retraction.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Published online: 13 May 2020

References

1. Boldt J (2000) Volume therapy in the intensive care patient – we are still confused, but. *Intensive Care Med* 26:1181–1192. <https://doi.org/10.1007/s001340000625>
2. Boldt J, Müller M, Mentges D et al (1998) Volume therapy in the critically ill: is there a difference? *Intensive Care Med* 24:28–36. <https://doi.org/10.1007/s001340050511>
3. Boldt J, Heesen M, Muller M et al (1996) The effects of albumin versus hydroxyethyl starch solution on cardiorespiratory and circulatory variables in critically ill patients. *Anesth Analg* 83(2):254–261. <https://doi.org/10.1097/0000539-199608000-00010> 528.835 pt528.835 pt
4. Boldt J, Mueller M, Menges T et al (1996) Influence of different volume therapy regimens on regulators of the circulation in the critically ill. *Br J Anesth* 77(4):480–487. <https://doi.org/10.1093/bja/77.4.480>

*Correspondence:

<https://www.springer.com/journal/134/>

The original article can be found online at <https://doi.org/10.1007/s001340000625>.