


NARRATIVE REVIEW



Management of hematological patients requiring emergency chemotherapy in the intensive care unit

Antoine Lafarge^{1*} , Dara Chean¹, Livia Whiting¹ and Raphaël Clere-Jehl^{2,3} on behalf of Groupe de Recherche en Réanimation Respiratoire du patient d'Onco-Hématologie (GRRR-OH) and Clinical Research in Intensive Care and Sepsis - TRIal Group for Global Evaluation and Research in SEpsis (CRICS-TRIGGERSEP)

© 2024 The Author(s)

Abstract

Hematological malignancies may require rapid-onset treatment because of their short doubling time, notably observed in acute leukemias and specific high-grade lymphomas. Furthermore, in targeted onco-hematological scenarios, chemotherapy is deemed necessary as an emergency measure when facing short-term, life-threatening complications associated with highly chemosensitive hematological malignancies. The risks inherent in the disease itself, or in the initiation of treatment, may then require admission to the intensive care unit (ICU) to optimize monitoring and initial management protocols. Hyperleukocytosis and leukostasis in acute leukemias, tumor lysis syndrome, and disseminated intravascular coagulation are the most frequent onco-hematological complications requiring the implementation of emergency chemotherapy in the ICU. Chemotherapy must also be started urgently in secondary hemophagocytic lymphohistiocytosis. Tumor-induced microangiopathic hemolytic anemia and plasma hyperviscosity due to malignant monoclonal gammopathy represent infrequent yet substantial indications for emergency chemotherapy. In all cases, the administration of emergency chemotherapy in the ICU requires close collaboration between intensivists and hematology specialists. In this review, we provide valuable insights that aid in the identification and treatment of patients requiring emergency chemotherapy in the ICU, offering diagnostic tools and guidance for their overall initial management.

Keywords: Chemotherapy, Emergency, Acute leukemia, Lymphoma, ICU

Introduction

Some hematological malignancies have short tumor doubling times (TDT) and specific related complications incompatible with scheduled initiation of treatment, necessitating rapid-onset chemotherapy. This emergency is particularly evident in highly chemosensitive diseases like acute leukemias and high-grade lymphomas, characterized by short TDT, initial chemosensitivity, and

frequent life-threatening complications without prompt treatment [1–3].

Chemotherapy administration in the intensive care unit (ICU) for malignancy-related complications, once considered a double-edged sword, has emerged as a potentially lifesaving intervention, even in the presence of concurrent infections or organ failures [2, 4, 5]. While studies focusing on ICU chemotherapy in hematological patients report mortality rates ranging from 25% [6] to 32% [2], available data suggest encouraging long-term outcomes with a reported 12-month survival rate of 30% among cancer patients receiving ICU chemotherapy, with 70% achieving complete remission [1].

Beyond drug-related toxicities, chemotherapy can exacerbate tumor-associated complications, potentially

*Correspondence: antoine.lafarge@aphp.fr

¹ Médecine Intensive et Réanimation, APHP, Saint-Louis Hospital and Paris University, Paris, France

Full author information is available at the end of the article

leading to life-threatening conditions. Hence, the initiation of chemotherapy represents a precarious phase that may necessitate preemptive ICU admission for optimized monitoring and management [7].

The literature on this topic is limited, and the beneficial effects of chemotherapy administered during ICU stays remain a subject of debate [8]. However, anticancer therapy in the ICU is feasible and might serve as a bridge to cure for selected, well-informed hematological patients with reasonable prognostic expectations [6, 9]. Intensivists play a crucial role in identifying patients who could benefit from urgent chemotherapy, collaborating closely with hematologists, as timely diagnostic and therapeutic interventions may significantly impact prognosis [4, 10].

In this review, we will first discuss common scenarios requiring or complicating urgent chemotherapy, most of the time in an ICU setting: hyperleukocytosis and leukostasis in acute leukemia, tumor lysis syndrome (TLS), and disseminated intravascular coagulation (DIC). Chemotherapy must also be started urgently in secondary hemophagocytic lymphohistiocytosis (HLH) [11]. Although less frequent, tumor-induced microangiopathic hemolytic anemia (MAHA) and plasma hyperviscosity due to malignant monoclonal gammopathy also necessitate prompt initiation of chemotherapy.

The objective of this review is to offer clinicians guidance and support in identifying and managing patients requiring urgent chemotherapy.

Hematological complications requiring emergency chemotherapy in the ICU

Hyperleukocytosis and leukostasis in acute leukemia

Hyperleukocytosis is often an unexpected finding in outpatients or in the emergency ward, frequently accompanied by non-specific symptoms like fever, dyspnea, or mucocutaneous bleeding. Hyperleukocytosis is defined as a circulating white blood cell count greater than $100 \times 10^9/L$ [12], although the associated complications can occur as early as $50 \times 10^9/L$ and may require ICU management. Hyperleukocytosis is present at diagnosis in up to 20% and 30% of patients with acute myeloid leukemia (AML) and with acute lymphoblastic leukemia (ALL) [13], respectively. Hyperleukocytosis can also be found in chronic leukemias, but is less commonly associated with complications, except in the acceleration phase of chronic myeloid or myelomonocytic leukemia, which highlights the role of circulating myeloid blasts in the genesis of leukostasis [14].

Leukostasis is a hyperviscosity syndrome involving cellular components encountered in up to 30% of hyperleukocytic AML [15], especially with monocytic characteristics and FLT3-ITD mutation [16]. Leukostasis is

Take-home messages

In targeted onco-hematological scenarios, chemotherapy is deemed necessary as an emergency measure when facing short-term, life-threatening complications associated with highly chemosensitive hematological malignancies.

The administration of emergency chemotherapy in the intensive care unit requires close collaboration between intensivists and hematology specialists, and may contribute to improving the prognosis of hematological malignancies.

more rarely described in ALL, mostly in pediatric populations, and is associated with particularly high leukocytosis, greater than $400 \times 10^9/L$ [14]. In this context, leukostasis often manifests alongside multiorgan failure, tumor lysis syndrome, and severe coagulopathy.

The pathophysiology of leukostasis involves endothelial dysfunction, due to the adhesion of blast cells to the endothelium, followed by their massive migration across the endothelial barrier [17]. This phenomenon is initiated by the exaggerated expression of integrins and selectins by leukemia cells, which amplify endothelial dysfunction through the production of proinflammatory cytokines [18]. The endothelial dysfunction leads to a rupture of the vascular barrier, a perivascular tissue infiltration by the blasts, and hemorrhagic phenomena. The role of hyperleukocytosis-induced blood hyperviscosity itself has long been suggested [19], and although it is only part of the mechanisms involved in leukostasis, this hyperviscosity underscores the importance of minimizing red blood cell transfusions during the hyperleukocytic phase of acute leukemia [14].

Pulmonary involvement is the most common manifestation of leukostasis, occurring in up to 40% of cases [20]. Clinical signs are non-specific and may include cough, dyspnea, and hypoxemia. Chest radiographs show non-specific patterns, with frequent bilateral diffuse alveolar infiltrates [21]. Pulmonary leukostasis often requires the initiation of mechanical ventilation as it can evolve into acute respiratory distress syndrome [22].

Brain involvement occurs in up to 30% of patients with leukostasis and manifests as delirium, focal neurologic deficits, or seizures. Brain imaging shows ischemic or hemorrhagic lesions that may be localized or diffuse [23], and in which blood hyperviscosity appears to play an important role (Fig. 1).

Hyperleukocytosis, whether accompanied by leukostasis or not, is linked to a high mortality rate ranging from 25% to 50% in the initial weeks [17], which justifies ICU admission for close monitoring. Patients with hyperleukocytosis are at high-risk of respiratory deterioration after the initiation of induction chemotherapy, commonly termed acute lysis pneumopathy [22]. Effective




Fig. 1 Inaugural complications of acute myeloid leukemia requiring emergency chemotherapy. **A** Pulmonary leukostasis, with bilateral alveolar and interstitial opacities, related to hyperleukocytic ($230 \times 10^9/L$) acute monocytic leukemia (ex FAB-M5) at the time of diagnosis. **B** Pulmonary leukostasis, with ground-glass opacity in the right upper lobe, secondary to hyperleukocytic ($200 \times 10^9/L$) acute myelomonocytic leukemia (ex FAB-M4). **C** Fundus image with fluorescein angiography revealing hyperviscosity-related retinopathy with disseminated retinal hemorrhages and vascular dilation. **D, E** T2*-weighted and T2-flair MRI sequences revealing inaugural left fronto-temporal hemorrhage and cerebral edema linked to hyperleukocytic ($330 \times 10^9/L$) acute monocytic leukemia (ex FAB-M5). **F** Acute monocytic leukemia (ex FAB-M5): peripheral blood smear with numerous promonocytes that have folded and convoluted nuclei and a finely granulated cytoplasm and vacuoles. *DIC* disseminated intravascular coagulation, *FAB* French–American–British classification, *MRI* magnetic resonance imaging

management encompasses addressing organ failures and implementing treatments such as hyperhydration and urgent cytoreduction before introducing anthracycline and cytarabine-based induction chemotherapy (Table 1) [12].

Hydroxyurea has long been described as a reliable method to reduce cell counts, taper clinical leukostasis manifestations, and lower the risk of developing complications from the cytotoxic effects associated with definitive induction chemotherapy [24, 25]. The gradual reduction of cell counts with hydroxyurea has been

Table 1 Main hematological complications requiring emergency chemotherapy

Indication	Clinical and biological features	Etiologies	Initial management
Hyperleukocytosis in acute leukemia	Leukostasis (cerebral and/or pulmonary) Tumor lysis syndrome ^b DIC ^c	AML with monocytic features (ex FAB-M4, M5, or M3v) ^a ALL ⁺⁺ , AML ⁺ Acute promyelocytic leukemia ⁺⁺ (APL, ex FAB-M3)	Hyperhydration (3L/m ²) Visual LVEF assessment advised Avoid RBC transfusion initially Early cytereuction AML: hydroxyurea ALL: dexamethasone Induction chemotherapy
Tumor lysis syndrome ^b	Biological TLS ↑LDH ↑K ⁺ , ↑P, ↓Ca ²⁺ Hyperuricemia Clinical TLS Acute kidney injury Seizures Arrhythmia	Acute leukemia, according to hyperleukocytosis (lymphoblastic ⁺⁺) High grade non-Hodgkin lymphoma (Burkitt ⁺⁺ , diffuse large B-cell lymphoma)	Hyperhydration (3 L/m ²) Hydro-electrolyte rebalancing (hypocalcemia should not be corrected unless seizures occur) Rasburicase (↓ uric acid) Early renal replacement therapy Reducing tumor mass: Acute leukemia: cf. supra Lymphoma: corticosteroids Less aggressive first chemotherapy
Disseminated intravascular coagulation ^c	Blood tests Thrombocytopenia ↑INR ↑FDP or D-dimers ↓fibrinogen Clinical forms Hemorrhagic (cerebral ⁺⁺) Thrombotic	Hemorrhagic (hyperfibrinolytic) APL (ex FAB-M3) ⁺⁺⁺ AML (monocytic features ^a) Thrombotic (hypofibrinolytic) ALL (L-asparaginase ⁺⁺) Metastatic adenocarcinoma ^d	Platelet transfusion (> 20 G/L, 50 G/L if active bleeding) Etiological treatment: APL: ATRA ⁺⁺⁺ Other AML: cf. supra Lower platelet transfusion threshold Anticoagulation according to platelet count (LMWH or UFH) Dexamethasone (ALL induction phase)
Tumor-induced microangiopathic hemolytic anemia	Blood tests Thrombocytopenia Hemolysis, schistocytes Negative DAT Clinical forms Atypical forms (not fitting with HUS or TTP)	Metastatic adenocarcinoma ^d Non-Hodgkin lymphomas Differential diagnoses •TTP •Infectious causes •Bone marrow invasion •Drug-induced MAHA ^e	Exclusion of alternative diagnoses ADAMTS13 > 10% (rules out TTP) Search for sepsis and HIV Consider bone marrow biopsy Check medication history ^e Lymphoma: corticosteroids Rapid-onset chemotherapy
Malignancy-associated hemophagocytic lymphohistiocytosis	Biological ↑ferritin, ↑TG Cytopenias, DIC Hemophagocytosis Hepatic cytolysis Clinical Fever, lymphadenopathy Hepatosplenomegaly	Non-Hodgkin B- or T-cell lymphomas Differential diagnoses •Infection (EBV ⁺⁺ , sepsis) •Autoimmune (Lupus, Still)	High-dose dexamethasone Etoposide ⁺⁺ (lymphoma, EBV) Rituximab (B-cell lymphoma, EBV) Emergency chemotherapy (DEP or CHOEP regimen) ^f Ruxolitinib (hematological malignancies, HSCT)
Plasma hyperviscosity syndrome	Biological ↑protidemia ↑ monoclonal Ig (IgM ⁺⁺) ± cryoglobulinemia Clinical CNS symptoms Visual disturbances Mucosal bleeding Fundoscopic exam Dilated veins Hemorrhage	Myeloma (IgG, IgA) Waldenström macroglobulinemia ⁺⁺⁺ Others: rituximab IgM flare, Types I and II cryoglobulinemia, Sjögren syndrome	Hyperhydration Plasmapheresis (with albumin) Avoid RBC transfusion Treatment of underlying disease Dexamethasone Chemotherapy Waldenström: rituximab
Multiple myeloma-associated complications	Hypercalcemia Ion. Ca ²⁺ ≥ 1.33 mmol/L Acute kidney injury ↑ urine free light chain ↓ albumin excretion	Newly diagnosed or refractory multiple myeloma	Hyperhydration Consider early RRT (AKI) BDdoublet ^g Hypercalcemia: bisphosphonates (pamidronate ⁺⁺)

ADAMTS13 a disintegrin and metalloprotease with thrombospondin type I repeats-13, ALL acute lymphoblastic leukemia, AML acute myeloid leukemia, APL acute promyelocytic leukemia, ATRA all-trans retinoic acid, ion. Ca²⁺ ionized calcium, CNS central nervous system, DAT direct antiglobulin test, DIC disseminated intravascular coagulation, EBV Epstein-Barr virus, FAB French-American-British classification, FDP fibrin degradation products, HIV human immunodeficiency virus, HSCT hematopoietic stem cell transplantation, HUS hemolytic uremic syndrome, Ig immunoglobulin, INR international normalized ratio, K⁺ potassium, LDH lactate

Table 1 (continued)

dehydrogenase, *LMWH* low-molecular-weight heparin, *LVEF* left ventricular ejection fraction, *P* phosphorus, *RBC* red blood cell, *RRT* renal replacement therapy, *TG* triglycerides, *tiMAHA* tumor-induced microangiopathic hemolytic anemia, *TTP* thrombotic thrombocytopenic purpura, *TLS* tumor lysis syndrome, *UFH* unfractionated heparin

^a The French–American–British classification is no longer standard for the diagnosis, but distinguished acute myeloid leukemias based on cell morphology, with AMLs with monocytic features being represented by AML4 (acute myelomonocytic leukemia), AML5 (acute monocytic leukemia), and AML3v (microgranular variant of acute promyelocytic leukemia)

^b Tumor lysis syndrome is defined by the diagnostic criteria of Cairo and Bishop (see also Supplementary Table 1) [35]

^c Disseminated intravascular coagulation is diagnosed according to the *International Society on Thrombosis and Hemostasis (ISTH)* criteria (see also Table 2) [46]

^d The metastatic adenocarcinomas involved in paraneoplastic DIC and tiMAHA are mostly of digestive, pancreatic, mammary gland, prostatic, or pulmonary origin

^e Several anti-tumor treatments may induce MAHA (e.g., gemcitabine, proteasome inhibitors, and oxaliplatin therapy)

^f For the treatment of lymphoma-associated HLH, the main first-line chemotherapies include the DEP (dexamethasone, etoposide, and cisplatin) regimen or the CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, and prednisone) regimen

^g The BD doublet includes bortezomib (proteasome inhibitor) and dexamethasone

shown to decrease hospital mortality in patients with hyperleukocytic AML when compared to a matched population that received induction chemotherapy without prior cytoreduction [24]. However, response to hydroxyurea has not been shown to be predictive of favorable short- and long-term outcomes [25]. The use of hydroxyurea is recommended by international guidelines to reduce white blood cells (WBC) count below $25 \times 10^9/L$ although high-quality evidence to support this treatment is currently lacking [26].

Leukapheresis remains the frontline treatment for hyperleukocytosis and leukostasis in a large number of specialized centers worldwide [27]. While this procedure rapidly removes circulating blasts, it cannot eliminate cellular plugs already formed within vessels or inhibit cellular proliferation in the bone marrow. Although supported by several single-center retrospective studies [28], the largest study conducted to date found no discernible impact of leukapheresis on overall survival [15]. Additionally, the complications associated with central catheter insertion in patients with thrombocytopenia or DIC and the risk of catheter-related infection must be carefully considered before initiating leukapheresis. Various cytoreduction regimens have been described with no prospective data supporting the use of one strategy over another [29].

Alongside cytoreduction, the use of dexamethasone during induction chemotherapy has been suggested to reduce short-term mortality when compared with either a historical cohort or a propensity score-matched cohort that did not receive dexamethasone [30, 31]. No impact on infection occurrence has been demonstrated; however, there is a lack of prospective data supporting this claim. Dexamethasone has been shown to downregulate the expression of adhesion molecules involved in leukocytes adhesion, potentially alleviating the symptoms of leukostasis [32].

Tumor lysis syndrome

TLS is a life-threatening condition secondary to the release of tumor cells content into the bloodstream [33], including deoxyribonucleic acid (DNA) and its metabolites, intracellular ions (potassium, phosphorus), as well as evocative proteins like lactate dehydrogenase (LDH) and certain cytokines. The degradation of the purine bases of DNA (adenine, guanine) leads to the increase of plasmatic uric acid and the formation of uric acid crystals. Released phosphorus precipitates with calcium, which induces both hypocalcemia and the formation of calcium phosphate crystals [33, 34]. Thus, TLS-associated laboratory abnormalities include hyperuricemia, hyperphosphatemia, hypocalcemia, and hyperkalemia, which collectively define the biological TLS according to Cairo and Bishop (see supplementary Table 1). The combination of biological TLS with acute kidney injury (AKI), seizures (related to hypocalcemia), ventricular arrhythmia, or sudden cardiac death (related to hyperkalemia) defines the clinical TLS [35].

TLS arises in patients with chemosensitive malignancy and high tumor burden. It is therefore common in hyperleukocytic leukemia [17] or high-grade non-Hodgkin lymphomas such as Burkitt's lymphoma [36]. Although TLS mainly affects patients undergoing chemotherapy, spontaneous TLS might also occur. It is estimated that up to 40% of TLS happen spontaneously in high-risk patients [37]. Paradoxically, as a marker of high tumor burden, spontaneous TLS is an indication for urgent chemotherapy and close monitoring.

AKI severely affects the prognosis of TLS, and is associated with a 66% mortality rate at 6 months, compared to 21% in the absence of renal failure [37]. Thus, hyperhydration ($3L/m^2/day$ or $70 mL/kg$) is the treatment's cornerstone [33]. Alkalinization is no longer recommended because it is associated with the risk of precipitation of calcium phosphate crystals [38]. After stopping any phosphorus and potassium supplementation, the hydro-electrolyte rebalancing is aimed at

correcting hyperkalemia by using polystyrene sulfonate alone in the absence of electrocardiographic (ECG) signs, and by combining insulin-glucose in hyperkalemia with ECG signs to avoid alkalization. The efficacy of phosphate binders on hyperphosphatemia is disputed in this context, and hypocalcemia should not be supplemented (except in cases of convulsions, or hyperkalemia with ventricular arrhythmia) due to the precipitation of calcium with phosphorus [33]. Electrolyte disturbances (hyperkalemia, hyperphosphatemia) require initial monitoring in the ICU, including ECG monitoring.

Due to the risk of uratic nephropathy, the treatment includes a hypouricemic therapy, favoring rasburicase over allopurinol starting from the biological TLS stage [39]. However, ruling out glucose-6-phosphate dehydrogenase (G6PD) deficiency is essential beforehand to prevent rasburicase-induced methemoglobinemia.

Renal replacement therapy (RRT) should be discussed early [40], in addition to the usual criteria, in the event of persistent hyperphosphatemia, although no threshold could be determined by consensus [34]. In this context, the early consideration of hyperphosphatemia is part of an approach to prevent the onset or worsening of AKI. Clinical, ECG, and biological monitoring is essential before and during the initiation of anti-tumor treatment (cytoreduction or chemotherapy). The main biological abnormalities (serum potassium, phosphoremia, calcemia, uremia, creatinine, uric acid level, and eventually LDH) must be checked every 4–6 h during the acute phase [33].

Disseminated intravascular coagulation

DIC of tumor origin is encountered in different clinical forms depending on the causative malignancy, and is categorized into two groups: hyperfibrinolytic, associated with a hemorrhagic presentation, and antifibrinolytic, characterized by thrombotic manifestations [41]. The hemorrhagic phenotype is prominent in DIC induced by hematological malignancies and mainly concerns AML, while myeloproliferative disorders and asparaginase-treated ALL are preferentially associated with thrombotic phenotypes [41, 42].

Incidence of DIC among acute leukemias is estimated at 15–25% at the time of diagnosis [42]. The incidence is higher in the presence of hyperleukocytosis with leukostasis, especially in cases of AML that display a normal karyotype with FLT3-ITD mutation [43]. The pathophysiology of leukemia-induced DIC involves the production of tissue factor by leukemic cells, which triggers the coagulation cascade through the extrinsic pathway, and initiates multifactorial endothelial activation and dysfunction [44]. The hyperfibrinolytic and hemorrhagic presentation

is particularly common in acute promyelocytic leukemia (APL), in which leukemia cells activate plasminogen through the expression of tissue plasminogen activator (tPA) thereby exacerbating fibrinolysis [45]. DIC is initially present in up to 80% of APL cases [42] according to the International Society on Thrombosis and Hemostasis (ISTH) definition [46]. ISTH criteria include thrombocytopenia, prolonged prothrombin time (PT), decreased fibrinogen, and increased fibrin degradation products (FDPs), or D-dimers. Thrombocytopenia is a questionable criterion in this context, as it may result from bone marrow invasion. Hence, the Japanese Ministry of Health and Welfare (JMHW) has proposed a platelet-free score for the diagnosis of DIC in patients with hematological malignancies (Table 2) [47].

Severe clinical manifestations of DIC are represented by cerebral hemorrhages, followed by intra-alveolar hemorrhages [42]. In patients with active bleeding, the platelet transfusion threshold should attain $50 \times 10^9/L$ and fibrinogen replacement therapy is indicated to reach a level $>1.5 \text{ g/L}$ [42, 48]. Acute leukemia-associated DIC is an indication for emergency cytoreduction and chemotherapy (Table 1). The treatment of DIC specifically induced by APL includes trans-retinoic acid (ATRA), which should be administered as soon as the diagnosis is established. However, ATRA can cause a differentiation syndrome in which the massive transformation of blasts into mature cells induces acute febrile

Table 2 Definition of disseminated intravascular coagulation according to the criteria of the International Society on Thrombosis and Haemostasis [46] and the modified Japanese Ministry of Health and Welfare criteria [47]

Criteria	ISTH	JMHW ^a	Score
Clinical symptoms	NA	Organ failure	1
Platelet count (G/L)	> 100	NA	0
	50–100		1
	< 50		2
D-dimers or FDP	No increase	FDPs ≥ 20	0
	Moderate increase		1
	Severe increase		2
PT increase (s) ^b or PT ratio	< 3 s	1.25–1.67 (PT ratio)	0
	3–6 s	> 1.67 (PT ratio)	1
	> 6 s		2
Fibrinogen (g/L)	≥ 1	NA	0
	< 1		1
Diagnosis of DIC	≥ 5 points	≥ 4 points	

DIC disseminated intravascular coagulation, FDP fibrin degradation products, ISTH International Society on Thrombosis and Haemostasis, JMHW Japanese Ministry of Health and Welfare, NA not applicable, PT prothrombin time, s second

^a JMHW criteria are modified for patients with thrombocytopenia of central origin, including bone marrow invasion by hematological malignancies [47]. These modified criteria do not include points for platelet count or bleeding

^b The value of prothrombin time increase is given in second(s) above the normal upper limit

multiorgan failure and whose treatment is based on dexamethasone and cytoreduction by hydroxyurea [49]. Importantly, with the advent of targeted therapies for AML, such as inhibitors targeting isocitrate dehydrogenase (IDH-1/2) and FMS-like tyrosine kinase 3 (FLT3), differentiation syndrome is now increasingly reported in non-APL AML [50]. Thrombotic manifestations are more often encountered in ALL, mostly secondary to the prothrombotic effect of L-asparaginase [51]. Through reduced concentrations of asparagine, hepatic synthesis of antithrombin and fibrinogen is lowered, resulting in both a prothrombotic state and hypofibrinogenemia in asparaginase-treated ALL patients [42]. Thus, antithrombin concentrates are recommended by the ISTH to achieve a target level of 80% to 120% and low-molecular-weight heparin thromboprophylaxis is strongly suggested in patients with a platelet count greater than $30 \times 10^9/L$ without bleeding [51]. A lower threshold for platelet transfusion should be considered in asparaginase-treated ALL patients, but no precise guidelines have been published yet. Fibrinogen substitution is implemented in most studies to achieve a target level >0.5 g/L [51]. Unfractionated heparin or low-molecular-weight heparin is used preferentially to oral anticoagulants to treat thromboembolic events [52].

Tumor-induced microangiopathic hemolytic anemia

Tumor-induced MAHA is a differential diagnosis of paraneoplastic DIC in the presence of peripheral thrombocytopenia [53]. In approximately 10–15% of cases, MAHA and DIC can co-exist [53, 54], and they appear to constitute the two facets of tumor-induced endothelial activation and dysfunction [41]. While DIC firstly activates secondary hemostasis through the production of tissue factor by endothelial cells, the pathophysiology of MAHA relies primarily on excessive platelet adhesion and activation, in correlation with an enhanced von Willebrand factor (VWF) activity of endothelial origin. Overexpression of VWF often accompanies a partial decrease in a disintegrin and metalloprotease with thrombospondin type I repeats-13 (ADAMTS13) protease activity [55], the physiological role of which is to cleave and inhibit VWF. This process leads to the formation of microthrombi, resulting in thrombocytopenia by consumption, and secondarily to intravascular hemolysis with the presence of schistocytes, by fragmentation of red blood cells in contact with microthrombi. Erythrocyte fragmentation may also be related to disseminated microvascular metastases or widespread bone marrow involvement [54, 56]. Therefore, the diagnosis of MAHA is based on a triad of thrombocytopenia, LDH elevation and the presence of schistocytes, along with a negative

direct antiglobulin test. The search for schistocytes is a major diagnostic element and should be repeated in case of initial negativity [57].

Most cases of paraneoplastic MAHA occur in patients with solid tumors, followed in frequency by non-Hodgkin lymphomas [56, 58]. A notable characteristic of paraneoplastic MAHA is the prevalence of unclassified forms because they do not fit with hemolytic uremic syndrome (HUS) or thrombotic thrombocytopenic purpura (TTP) [58]. When available, ADAMTS13 activity is often decreased in tumor-induced MAHA, but does not reach the diagnostic threshold characterizing TTP (less than 10%), whether in solid tumors [59] or hematological malignancies [60]. This reinforces the idea that treatment strategies cannot be modeled on those used for TTP [61]. Corticosteroids, rituximab and caplacizumab cannot be universally recommended and should be considered on a case-by-case basis. In scenarios of suspected TTP without determined etiology, failure of plasmapheresis should raise the diagnosis of paraneoplastic MAHA, and in these circumstances bone marrow biopsy is recommended by some authors [56]. In the absence of recent clinical trials focused on paraneoplastic MAHA, only emergency chemotherapy has been shown to be associated with survival thus far [58]. Infectious or treatment-related etiologies must be ruled out, including gemcitabine, proteasome inhibitors (especially carfilzomib) and oxaliplatin therapy [54, 62]. Regarding immunomodulatory drugs, lenalidomide and checkpoint inhibitors may also trigger authentic immune-mediated TTP through the expression of an anti-ADAMTS 13 autoantibody [54]. Finally, hematopoietic stem cell transplantation (HSCT)-induced MAHA is a well-documented complication [54, 62]

Secondary hemophagocytic lymphohistiocytosis

Malignancy-associated secondary HLH is a hyperinflammatory state occurring in cancer patients. It results from an excessive stimulation of macrophages under the effect of an inappropriate cytokine production, especially interferon- γ by CD8⁺ T cells, accompanied by a loss of regulation by natural killer (NK) cells. Activated macrophages in turn produce cytokines in inappropriate quantities, including interleukin (IL) -6 and tumor necrosis factor (TNF), which is enhanced during cell lysis and boosted by tumor-infiltrating lymphocytes [11, 63]. Clinical signs are non-specific and include high fever, lymphadenopathy, and hepatosplenomegaly [11]. The hallmark biological signs combine cytopenia, hepatic cytolysis, hyperferritinemia, hypertriglyceridemia, and coagulation disorders, with hypofibrinogenemia being associated with mortality [63]. Hemophagocytosis, although frequently found in bone marrow aspiration smears, is not imperative for diagnosis. The diagnosis is

based on the HLH-2004 diagnostic criteria (supplementary Table 2) [64]. The HScore (HLH-probability calculator) is an alternative diagnostic tool available online (<http://saintantoine.aphp.fr/score/>) [65]. Therefore, HLH is highly suspected if 5 of 8 criteria are fulfilled, and the diagnosis of HLH should be consistent with the overall clinical assessment and patient's history [66]. Onco-hematological diseases account for 97% of malignancy-induced HLH, with non-Hodgkin lymphoma ranking first (roughly 74% of cases) followed by leukemia [11]. Secondary HLH is independently associated with mortality in lymphoma patients.

As untreated secondary HLH is systematically lethal [63], the administration of high-dose dexamethasone and emergency etoposide-based chemotherapy to eliminate activated T cells is a therapeutic priority in onco-hematological patients [66, 67]. Completion of the entire first-line chemotherapy regimen appropriate to the lymphoma subtype significantly improves patient prognosis. This includes combinations of rituximab, cyclophosphamide, doxorubicin, and vincristine in case of diffuse large B cell lymphoma (DLBCL) [68], resulting in an adapted regimen such as CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, and prednisone) regimen [66].

Etoposide must also be administered urgently in certain non-malignant underlying indications, especially in the case of Epstein-Barr virus (EBV)-induced HLH, which requires the rapid addition of rituximab to deplete EBV-infected B-cells [11]. However, first-line administration of emergency chemotherapy does not apply to all cases of secondary HLH. For this reason, the etiological assessment of HLH must be meticulous, with the main treatment remaining that of the underlying cause. Ruxolitinib, a selective Janus kinase (JAK) inhibitor, used as a cytokine-reducing therapy with minimal myelosuppressive effects, is currently being incorporated into different treatment regimens for HLH secondary to hematological malignancies and HSCT. Emapalumab, a monoclonal antibody targeting interferon-gamma, is increasingly used in various centers for the treatment of secondary HLH. However, its application derived from a narrow subset of pediatric patients with primary HLH, warrants further assessment [69]. These new etoposide-reducing protocols help limit treatment toxicities before the initiation of disease-specific chemotherapy [70].

Plasma hyperviscosity

In contrast to "cellular" hyperviscosity, plasma hyperviscosity syndrome (HVS) results from an abnormal protein increase, often arising from plasma cell disorders with immunoglobulin (Ig). Due to their pentameric structure and their high molecular weight, IgMs are more likely to

induce HVS, which explains the importance of Waldenström macroglobulinemia (WM) among etiologies [71, 72]. At diagnosis, 40% of WM patients present with HVS, compared to 3.4% of multiple myeloma (MM) patients [72, 73]. Notably, the presence of a cryoglobulin can trigger or worsen plasma hyperviscosity, due to the cold-induced polymerization of the immunoglobulin involved.

Clinical signs of plasma hyperviscosity are dominated by a triad [74], associating neurological abnormalities (headache, delirium, tinnitus, ataxia), visual disturbances (blurring, phosphenes, myodesopsia), and mucosal bleeding (epistaxis, gum). In a context of HVS suspicion or for asymptomatic patients with a high protein concentration, the diagnosis can be rapidly confirmed by ophthalmologic examination revealing delayed venule clearance and/or retinal hemorrhage [72, 73, 75]. Beyond major hyperproteinemia, serum protein electrophoresis reveals the monoclonal hypergammaglobulinemia, characterized by immunofixation, with levels of IgM, IgG, and IgA generally greater than 30, 40, and 60 g/L, respectively [71]. Cryoglobulinemia should be specifically screened, especially in the presence of IgM [76].

Main vital complications include thrombosis, digestive and cerebral bleedings. Thus, HVS represents a therapeutic emergency that should not be delayed for ophthalmic examination in symptomatic patients. Early management combines ICU admission, hyperhydration, and the elimination of all factors that may increase hyperviscosity, such as diuretics or red blood cell transfusions, except in case of active uncontrolled bleeding. Plasmapheresis, performed urgently at 40 mL/kg with albumin as the primary replacement fluid, except in cases of active hemorrhage, is essential [77]. One to three therapeutic plasma exchange(s) may be sufficient to resolve symptoms of HVS. In otherwise asymptomatic patients, retinal changes respond dramatically to a single TPE with marked or complete reversal of the fundoscopic findings [75].

As serum Ig levels will return to baseline in 4 weeks, etiological treatment should be initiated soon after plasma therapy, to avoid treatment clearance. For newly diagnosed WM with bulky disease, a combination of bendamustine and rituximab is recommended as first line therapy. Therapeutic alternatives include the dexamethasone-rituximab-cyclophosphamide combination in cases of low disease burden, as well as novel agents such as proteasome (bortezomib and carfilzomib) and Bruton's tyrosine kinase (BTK) inhibitors (ibrutinib) [72, 78]. The specific treatment of MM is detailed in the following paragraph.

Myeloma-associated complications

Beyond HVS, newly diagnosed or refractory MM typically presents with life-threatening complications such

as hypercalcemia and AKI that may require ICU admission [79]. Light-chain cast nephropathy (LCCN) is the main cause of reversible AKI in MM patients and a frequent mode of diagnosis. LCCN must be evoked by the combination of low urinary albumin excretion and high free light chains (FLCs) level (>500 mg/l) [79]. Renal recovery, which is a major predictive factor for survival, mostly depends on early reduction of serum FLCs, particularly in patients requiring RRT [80]. Therefore, specific treatment initiation is an emergency that should not be delayed by RRT and/or ICU admission [79].

Early management of MM-associated AKI is based on the combination of symptomatic measures and emergency chemotherapy with high-dose corticosteroids. By reducing the urinary concentration of FLCs and enhancing renal tubular flow, hyperhydration using saline fluids reduces the risk of intratubular precipitation [79]. The effectiveness of urine alkalinization remains controversial and should be avoided in hypercalcemic patients because of the risk of calcium phosphate precipitation [81].

Treatment of hypercalcemia relies on rehydration and intravenous bisphosphonates, adapted to estimated glomerular filtration rate (eGFR) value. While zoledronic acid is not contraindicated, pamidronate is preferred due to its lower risk of renal toxicity [82]. Despite controversial clinical benefit, calcitonin might be initiated concurrently, as bisphosphonates may require several days to exert their optimal therapeutic effects. Nephrotoxic treatments should be discontinued.

Due to its anti-inflammatory, cytotoxic, and catabolic properties, high-dose dexamethasone (40 mg/day) improves renal recovery [83] and should be administered immediately after the diagnosis [84]. Prudent hydration and reduced-dose (20 mg/day) dexamethasone might be considered in most frail patients (cardiovascular comorbidities, age > 80 years).

Initiating early anti-plasma cell chemotherapy is essential to reduce the tumor burden and monoclonal FLCs secretion. Currently, the standard of care in LCCN patients combines high-dose dexamethasone with the proteasome inhibitor bortezomib, whose efficacy and tolerance are established without dose adaptation, even in patients requiring RRT [83]. However, the broader use of lenalidomide, carfilzomib, or cyclophosphamide is constrained due to unfavorable toxicity profiles and limited renal metabolism within the LCCN setting [83, 85]. It is anticipated that in the coming years, the combination of bortezomib with the monoclonal anti-CD38 antibody daratumumab is likely to further improve both hematologic response and renal recovery [86].

In addition, extracorporeal removal of circulating FLCs through plasmapheresis or intensive hemodialysis using new-generation “high-cutoff” (HCO) protein-leaking dialyzers might be considered. This approach aims to prevent the persistent accumulation of FLCs in both vascular and extravascular compartments [80].

Conclusion

Indications for emergency chemotherapy are relatively rare, yet demand early recognition due to the potentially rapid and lethal nature presented of some malignant complications. Admission to the ICU must be considered for the implementation of emergency chemotherapy and overall initial management particularly when an initial worsening is expected, due to the disease itself and/or its treatment. Depending on the etiology, the rapid onset of the antineoplastic treatment may lead to complete recovery, justifying full-code management, especially in the case of initial treatment. The broadening of indications for ICU admission, requiring close interprofessional collaboration, may contribute to the major prognostic improvement reported in onco-hematological patients. Dedicated multicenter studies are warranted to assess both the efficacy and safety of emergency chemotherapies in the ICU.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1007/s00134-024-07454-z>.

Author details

¹ Médecine Intensive et Réanimation, APHP, Saint-Louis Hospital and Paris University, Paris, France. ² Médecine Intensive et Réanimation, Hôpital de Haute-pierre, University Hospital of Strasbourg, Strasbourg, France. ³ Laboratoire d'ImmunoRhumatologie Moléculaire, INSERM (French National Institute of Health and Medical Research), UMR_S1109, Centre de Recherche d'Immunologie et d'Hématologie, University of Strasbourg, Strasbourg, France.

Acknowledgements

This work received non-financial support from the Grrr-OH Network (Groupe de Recherche en Réanimation Respiratoire du patient d'Onco-Hématologie) and CRICS group (Clinical Research in Intensive Care and Sepsis). We warmly thank Elie Azoulay (Saint Louis Hospital, Paris, France), who had the initial idea for writing this review, for his helpful contribution. We are also grateful to Guillaume Dumas (Grenoble, France) and Sofiane Fodil (Saint Louis Hospital, Paris, France) for authorization to use fundus and peripheral blood smear images.

Author contributions

All the authors participated in the writing and editing of the manuscripts. AL and RCJ reviewed the clinical trials.

Declarations

Conflicts of interest

The authors declare no competing financial interests.

Open Access

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing,

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 27 February 2024 Accepted: 18 April 2024

Published online: 15 May 2024

References

1. Wohlfarth P, Staudinger T, Sperr WR, Bojic A, Robak O, Hermann A, Laczika K, Carlstrom A, Riss K, Rabitsch W, Bojic M, Knoebl P, Locker GJ, Obiditsch M, Fuhrmann V, Schellongowski P, Arbeitsgruppe fur hamato-onkologische Intensivmedizin der Osterreichischen Gesellschaft fur Internistische und Allgemeine Intensivmedizin und N (2014) Prognostic factors, long-term survival, and outcome of cancer patients receiving chemotherapy in the intensive care unit. *Ann Hematol* 93:1629–1636
2. Benoit DD, Depuydt PO, Vandewoude KH, Offner FC, Boterberg T, De Cock CA, Noens LA, Janssens AM, Decruyenaere JM (2006) Outcome in severely ill patients with hematological malignancies who received intravenous chemotherapy in the intensive care unit. *Intensive Care Med* 32:93–99
3. Song JU, Suh GY, Chung MP, Kim H, Kwon OJ, Jung CW, Kang WK, Park K, Jeon K (2011) Risk factors to predict outcome in critically ill cancer patients receiving chemotherapy in the intensive care unit. *Support Care Cancer* 19:491–495
4. Moors I, Pene F, Lengline E, Benoit D (2015) Urgent chemotherapy in hematological patients in the ICU. *Curr Opin Crit Care* 21:559–568
5. Vandijck DM, Benoit DD, Depuydt PO, Offner FC, Blot SI, Van Tilborgh AK, Nolle J, Steel E, Noens LA, Decruyenaere JM (2008) Impact of recent intravenous chemotherapy on outcome in severe sepsis and septic shock patients with hematological malignancies. *Intensive Care Med* 34:847–855
6. Pastores SM, Goldman DA, Shaz DJ, Kostecky N, Daley RJ, Peterson TJ, Tan KS, Halpern NA (2018) Characteristics and outcomes of patients with hematologic malignancies receiving chemotherapy in the intensive care unit. *Cancer* 124:3025–3036
7. Lengline E, Raffoux E, Lemiale V, Darmon M, Canet E, Boissel N, Schlemmer B, Dombret H, Azoulay E (2012) Intensive care unit management of patients with newly diagnosed acute myeloid leukemia with no organ failure. *Leuk Lymphoma* 53:1352–1359
8. de Oliveira MCF, Ferreira JC, Nassar Junior AP, Dettino ALA, Caruso P (2020) Impact of urgent chemotherapy in critically ill patients. *J Intensive Care Med* 35:347–353
9. Zerbib Y, Rabbat A, Fartoukh M, Bige N, Andrejak C, Mayaux J, De Prost N, Misset B, Lemiale V, Bruneel F, Maizel J, Ricome S, Jacobs F, Bornstain C, Dupont H, Baudin F, Azoulay E, Pene F, Groupe de Recherche sur la Reanimation Respiratoire en O-H (2017) Urgent chemotherapy for life-threatening complications related to solid neoplasms. *Crit Care Med* 45:e640–e648
10. Sekeres MA, Elson P, Kalaycio ME, Advani AS, Copelan EA, Faderl S, Kantarjian HM, Estey E (2009) Time from diagnosis to treatment initiation predicts survival in younger, but not older, acute myeloid leukemia patients. *Blood* 113:28–36
11. Vick EJ, Patel K, Prouet P, Martin MG (2017) Proliferation through activation: hemophagocytic lymphohistiocytosis in hematologic malignancy. *Blood Adv* 1:779–791
12. Dohner H, Weisdorf DJ, Bloomfield CD (2015) Acute myeloid leukemia. *N Engl J Med* 373:1136–1152
13. Desprez C, Kouatchet A, Marchand T, Mear JB, Tadie JM, Peterlin P, Chevalier P, Canet E, Couturier MA, Guillermin G, Bodenès L, Gyan E, Villate A, Ehrmann S, Lebreton A, Lester MA, Fronteau C, Larhantec G, Andre V, Riou J, Hunault-Berger M, Schmidt-Tanguy A, Orvain C (2023) Outcome of patients with newly diagnosed AML admitted to the ICU, including preemptive admission—a multi-center study. *Ann Hematol* 102:1383–1393
14. Macaron W, Sargsyan Z, Short NJ (2022) Hyperleukocytosis and leukostasis in acute and chronic leukemias. *Leuk Lymphoma* 63:1780–1791
15. Stahl M, Shallis RM, Wei W, Montesinos P, Lengline E, Neukirchen J, Bhatt VR, Sekeres MA, Fathi AT, König H, Luger S, Khan I, Roboz GJ, Cluzeau T, Martinez-Cuadron D, Raffoux E, Germing U, Umakanthan JM, Mukherjee S, Brunner AM, Miller A, McMahon CM, Ritchie EK, Rodriguez-veiga R, Itzykson R, Boluda B, Rabian F, Tormo M, Acuna-Cruz E, Rabinovich E, Yoo B, Cano I, Podoltsev NA, Bewersdorf JP, Gore S, Zeidan AM (2020) Management of hyperleukocytosis and impact of leukapheresis among patients with acute myeloid leukemia (AML) on short- and long-term clinical outcomes: a large, retrospective, multicenter, international study. *Leukemia* 34:3149–3160
16. Thiede C, Steudel C, Mohr B, Schaich M, Schakel U, Platzbecker U, Wermke M, Bornhauser M, Ritter M, Neubauer A, Ehninger G, Illmer T (2002) Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. *Blood* 99:4326–4335
17. Bewersdorf JP, Zeidan AM (2020) Hyperleukocytosis and leukostasis in acute myeloid leukemia: can a better understanding of the underlying molecular pathophysiology lead to novel treatments? *Cells* 9:2310
18. Stucki A, Rivier AS, Gikic M, Monai N, Schapira M, Spertini O (2001) Endothelial cell activation by myeloblasts: molecular mechanisms of leukostasis and leukemic cell dissemination. *Blood* 97:2121–2129
19. Steinberg MH, Charm SE (1971) Effect of high concentrations of leukocytes on whole blood viscosity. *Blood* 38:299–301
20. Porcu P, Danielson CF, Orazi A, Heerema NA, Gabig TG, McCarthy LJ (1997) Therapeutic leukapheresis in hyperleukocytic leukaemias: lack of correlation between degree of cyto-reduction and early mortality rate. *Br J Haematol* 98:433–436
21. Stefanski M, Jamis-Dow C, Bayerl M, Desai RJ, Claxton DF, Van de Louw A (2016) Chest radiographic and CT findings in hyperleukocytic acute myeloid leukemia: a retrospective cohort study of 73 patients. *Medicine (Baltimore)* 95:e5285
22. Moreau AS, Lengline E, Seguin A, Lemiale V, Canet E, Raffoux E, Schlemmer B, Azoulay E (2014) Respiratory events at the earliest phase of acute myeloid leukemia. *Leuk Lymphoma* 55:2556–2563
23. Algharras AA, Mamourian A, Coyne T, Mohan S (2013) Leukostasis in an adult with AML presenting as multiple high attenuation brain masses on CT. *J Clin Diagn Res* 7:3020–3022
24. Mamez AC, Raffoux E, Chevret S, Lemiale V, Boissel N, Canet E, Schlemmer B, Dombret H, Azoulay E, Lengline E (2016) Pre-treatment with oral hydroxyurea prior to intensive chemotherapy improves early survival of patients with high hyperleukocytosis in acute myeloid leukemia. *Leuk Lymphoma* 57:2281–2288
25. Cerrano M, Seegers V, Raffoux E, Rabian F, Sebert M, Itzykson R, Lemiale V, Ades L, Boissel N, Dombret H, Azoulay E, Lengline E (2020) Predictors and outcomes associated with hydroxyurea sensitivity in acute myeloid leukemia patients with high hyperleukocytosis. *Leuk Lymphoma* 61:737–740
26. Dohner H, Wei AH, Appelbaum FR, Craddock C, DiNardo CD, Dombret H, Ebert BL, Fenau P, Godley LA, Hasserjian RP, Larson RA, Levine RL, Miyazaki Y, Niederwieser D, Ossenkoppele G, Rolig C, Sierra J, Stein EM, Tallman MS, Tien HF, Wang J, Wierzbowska A, Lowenberg B (2022) Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood* 140:1345–1377
27. Stahl M, Pine A, Hendrickson JE, Litzow MR, Luger SM, Stone RM, Erba HP, Kim TK, Sekeres MA, Steensma DP, Komrokji RS, Gore SD, Zeidan AM (2018) Beliefs and practice patterns in hyperleukocytosis management in acute myeloid leukemia: a large U.S. web-based survey(). *Leuk Lymphoma* 59:2723–2726
28. Ventura GJ, Hester JP, Smith TL, Keating MJ (1988) Acute myeloblastic leukemia with hyperleukocytosis: risk factors for early mortality in induction. *Am J Hematol* 27:34–37

29. Zhao J, Bewersdorf JP, Jaszczur S, Kowalski A, Perreault S, Schiffer M, Gore S, Podoltsev N, Prebet T, Shallis R, Zeidan AM (2021) High dose cyclophosphamide for cytoreduction in patients with acute myeloid leukemia with hyperleukocytosis or leukostasis. *Leuk Lymphoma* 62:1195–1202
30. Azoulay E, Canet E, Raffoux E, Lengline E, Lemiale V, Vincent F, de Labarthe A, Seguin A, Boissel N, Dombret H, Schlemmer B (2012) Dexamethasone in patients with acute lung injury from acute monocytic leukaemia. *Eur Respir J* 39:648–653
31. Cerrano M, Chevret S, Raffoux E, Rabian F, Seberty M, Valade S, Itzykson R, Lemiale V, Ades L, Boissel N, Dombret H, Azoulay E, Lengline E (2023) Benefits of dexamethasone on early outcomes in patients with acute myeloid leukemia with hyperleukocytosis: a propensity score matched analysis. *Ann Hematol* 102:761–768
32. Nakagawa M, Bondy GP, Waisman D, Minshall D, Hogg JC, van Eeden SF (1999) The effect of glucocorticoids on the expression of L-selectin on polymorphonuclear leukocyte. *Blood* 93:2730–2737
33. Howard SC, Jones DP, Pui CH (2011) The tumor lysis syndrome. *N Engl J Med* 364:1844–1854
34. Zafrani L, Canet E, Darmon M (2019) Understanding tumor lysis syndrome. *Intensive Care Med* 45:1608–1611
35. Cairo MS, Bishop M (2004) Tumour lysis syndrome: new therapeutic strategies and classification. *Br J Haematol* 127:3–11
36. Darmon M, Vincent F, Camous L, Canet E, Bonmati C, Braun T, Caillot D, Cormillon J, Dimicoli S, Etienne A, Galicier L, Garnier A, Girault S, Hunault-Berger M, Marolleau JP, Moreau P, Raffoux E, Recher C, Thiebaud A, Thieblemont C, Azoulay E, Groupe de Recherche en Reanimation Respiratoire et O-H (2013) Tumour lysis syndrome and acute kidney injury in high-risk haematology patients in the rasburicase era. A prospective multicentre study from the Groupe de Recherche en Reanimation Respiratoire et Onco-Hematologique. *Br J Haematol* 162:489–497
37. Darmon M, Guichard I, Vincent F, Schlemmer B, Azoulay E (2010) Prognostic significance of acute renal injury in acute tumor lysis syndrome. *Leuk Lymphoma* 51:221–227
38. Jones GL, Will A, Jackson GH, Webb NJ, Rule S, Committee B, for Standards in H, (2015) Guidelines for the management of tumour lysis syndrome in adults and children with haematological malignancies on behalf of the British Committee for Standards in Haematology. *Br J Haematol* 169:661–671
39. Alakel N, Middeke JM, Schetelig J, Bornhauser M (2017) Prevention and treatment of tumor lysis syndrome, and the efficacy and role of rasburicase. *Onco Targets Ther* 10:597–605
40. Matuszkiewicz-Rowinska J, Malyszko J (2020) Prevention and treatment of tumor lysis syndrome in the era of onco-nephrology progress. *Kidney Blood Press Res* 45:645–660
41. Helms J, Iba T, Connors JM, Gando S, Levi M, Meziani F, Levy JH (2023) How to manage coagulopathies in critically ill patients. *Intensive Care Med* 49:273–290
42. Wang TF, Makar RS, Antic D, Levy JH, Douketis JD, Connors JM, Carrier M, Zwicker JI (2020) Management of hemostatic complications in acute leukemia: guidance from the SSC of the ISTH. *J Thromb Haemost* 18:3174–3183
43. Guo Z, Chen X, Tan Y, Xu Z, Xu L (2020) Coagulopathy in cytogenetically and molecularly distinct acute leukemias at diagnosis: comprehensive study. *Blood Cells Mol Dis* 81:102393
44. Ten Cate H, Leader A (2021) Management of disseminated intravascular coagulation in acute leukemias. *Hamostaseologie* 41:120–126
45. Kwaan HC, Weiss I, Tallman MS (2019) The role of abnormal hemostasis and fibrinolysis in morbidity and mortality of acute promyelocytic leukemia. *Semin Thromb Hemost* 45:612–621
46. Taylor FB Jr, Toh CH, Hoots WK, Wada H, Levi M, Scientific Subcommittee on Disseminated Intravascular Coagulation of the International Society on Thromb Haemostasis (2001) Towards definition, clinical and laboratory criteria, and a scoring system for disseminated intravascular coagulation. *Thromb Haemost* 86:1327–1330
47. Yanada M, Matsushita T, Suzuki M, Kiyoi H, Yamamoto K, Kinoshita T, Kojima T, Saito H, Naoe T (2006) Disseminated intravascular coagulation in acute leukemia: clinical and laboratory features at presentation. *Eur J Haematol* 77:282–287
48. Squizzato A, Hunt BJ, Kinasevitz GT, Wada H, Ten Cate H, Thachil J, Levi M, Vicente V, D'Angelo A, Di Nisio M (2016) Supportive management strategies for disseminated intravascular coagulation. An international consensus. *Thromb Haemost* 115:896–904
49. Sanz MA, Fenaux P, Tallman MS, Estey EH, Lowenberg B, Naoe T, Lengfelder E, Dohner H, Burnett AK, Chen SJ, Mathews V, Iland H, Rego E, Kantarjian H, Ades L, Avvisati G, Montesinos P, Platzbecker U, Ravandi F, Russell NH, Lo-Coco F (2019) Management of acute promyelocytic leukemia: updated recommendations from an expert panel of the European LeukemiaNet. *Blood* 133:1630–1643
50. Norsworthy KJ, Mulkey F, Scott EC, Ward AF, Przepiorka D, Charlab R, Dorff SE, Deisseroth A, Kazandjian D, Sridhara R, Beaver JA, Farrell AT, de Claro RA, Pazdur R (2020) Differentiation syndrome with ivosidenib and enasidenib treatment in patients with relapsed or refractory IDH-mutated AML: a U.S. Food and Drug Administration systematic analysis. *Clin Cancer Res* 26:4280–4288
51. Orvain C, Balsat M, Tavernier E, Marolleau JP, Pabst T, Chevallier P, de Gunzburg N, Cacheux V, Huguet F, Chantepie S, Caillot D, Chalandon Y, Frayre J, Bonmati C, Lheritier V, Ifrah N, Dombret H, Boissel N, Hunault-Berger M (2020) Thromboembolism prophylaxis in adult patients with acute lymphoblastic leukemia treated in the GRAALL-2005 study. *Blood* 136:328–338
52. Rank CU, Toft N, Tuckuviene R, Grell K, Nielsen OJ, Frandsen TL, Marquart HVH, Albertsen BK, Tedgard U, Hallbook H, Ruud E, Jarvis KB, Quist-Paulsen P, Huttunen P, Wartiovaara-Kautto U, Jonsson OG, Trakymiene SS, Griskevicius L, Saks K, Punab M, Schmiegelow K (2018) Thromboembolism in acute lymphoblastic leukemia: results of NOPHO ALL2008 protocol treatment in patients aged 1 to 45 years. *Blood* 131:2475–2484
53. Wada H, Matsumoto T, Suzuki K, Imai H, Katayama N, Iba T, Matsumoto M (2018) Differences and similarities between disseminated intravascular coagulation and thrombotic microangiopathy. *Thromb J* 16:14
54. Thomas MR, Scully M (2021) How I treat microangiopathic hemolytic anemia in patients with cancer. *Blood* 137:1310–1317
55. Colonne CK, Favaloro EJ, Pasalic L (2022) The intriguing connections between von Willebrand Factor, ADAMTS13 and cancer. *Healthcare (Basel)* 10(3):557
56. Decaestecker A, Hamroun A, Provot F, Rondeau E, Faguer S, Sallee M, Titeca-Beauport D, Rebibou JM, Forestier A, Azar R, Deltombe C, Wyncel A, Grange S, Fremaux Bacchi V, Cartery C (2023) Retrospective study of 59 cases of cancer-associated thrombotic microangiopathy: presentation and treatment characteristics. *Nephrol Dial Transplant* 38:913–921
57. Schmidt J, Zafrani L, Lemiale V, Stepanian A, Joly B, Azoulay E, Mariotte E (2021) The clinical picture of thrombotic microangiopathy in patients older than 60 years of age. *Br J Haematol* 192:e25–e28
58. Lechner K, Obermeier HL (2012) Cancer-related microangiopathic hemolytic anemia: clinical and laboratory features in 168 reported cases. *Medicine (Baltimore)* 91:195–205
59. Fontana S, Gerritsen HE, Kremer Hovinga J, Furlan M, Lammle B (2001) Microangiopathic haemolytic anaemia in metastasizing malignant tumours is not associated with a severe deficiency of the von Willebrand factor-cleaving protease. *Br J Haematol* 113:100–102
60. Yui JC, Garceau D, Jhaveri KD, Wanchoo R, Bijol V, Glezerman I, Hassoun H, Dispenzieri A, Russell SJ, Leung N (2019) Monoclonal gammopathy-associated thrombotic microangiopathy. *Am J Hematol* 94:E250–E253
61. Azoulay E, Bauer PR, Mariotte E, Russell L, Knoebel P, Martin-Loeches I, Pene F, Puxty K, Povoja P, Barratt-Due A, Garnacho-Montero J, Wendon J, Munshi L, Benoit D, von Bergwelt-Baildon M, Maggiorini M, Coppo P, Cataland S, Veyradier A, Van de Louw A, Nine-i I (2019) Expert statement on the ICU management of patients with thrombotic thrombocytopenic purpura. *Intensive Care Med* 45:1518–1539
62. Valerio P, Barreto JP, Ferreira H, Chuva T, Paiva A, Costa JM (2021) Thrombotic microangiopathy in oncology—a review. *Transl Oncol* 14:101081
63. Valade S, Mariotte E, Azoulay E (2020) Coagulation disorders in hemophagocytic lymphohistiocytosis/macrophage activation syndrome. *Crit Care Clin* 36:415–426
64. Henter JI, Horne A, Arico M, Egeler RM, Filipovich AH, Imashuku S, Ladisch S, McClain K, Webb D, Winiarski J, Janka G (2007) HLH-2004: diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer* 48:124–131
65. Fardet L, Galicier L, Lambotte O, Marzac C, Aumont C, Chahwan D, Coppo P, Hejblum G (2014) Development and validation of the HScore, a score for the diagnosis of reactive hemophagocytic syndrome. *Arthritis Rheumatol* 66:2613–2620

-
66. La Rosee P, Horne A, Hines M, von Bahr GT, Machowicz R, Berliner N, Birndt S, Gil-Herrera J, Girschikofsky M, Jordan MB, Kumar A, van Laar JAM, Lachmann G, Nichols KE, Ramanan AV, Wang Y, Wang Z, Janka G, Henter JI (2019) Recommendations for the management of hemophagocytic lymphohistiocytosis in adults. *Blood* 133:2465–2477
 67. Arca M, Fardet L, Galicier L, Riviere S, Marzac C, Aumont C, Lambotte O, Coppo P (2015) Prognostic factors of early death in a cohort of 162 adult haemophagocytic syndrome: impact of triggering disease and early treatment with etoposide. *Br J Haematol* 168:63–68
 68. Li B, Guo J, Li T, Gu J, Zeng C, Xiao M, Zhang W, Li Q, Zhou J, Zhou X (2021) Clinical characteristics of hemophagocytic lymphohistiocytosis associated with non-Hodgkin B-cell lymphoma: a multicenter retrospective study. *Clin Lymphoma Myeloma Leuk* 21:e198–e205
 69. Garonzi C, Chinello M, Cesaro S (2021) Emapalumab for adult and pediatric patients with hemophagocytic lymphohistiocytosis. *Expert Rev Clin Pharmacol* 14:527–534
 70. Stalder G, Suffiotti M, Segot A, Noto A, Pantaleo G, Spertini O, Obeid M (2023) Response-adjusted regimen combining ruxolitinib, etoposide and dexamethasone (adRED) in adult patients with acute myeloid leukemia-associated hemophagocytic lymphohistiocytosis: a single-center pilot trial. *Haematologica* 108:234–239
 71. Gustine JN, Meid K, Dubeau T, Hunter ZR, Xu L, Yang G, Ghobrial IM, Treon SP, Castillo JJ (2017) Serum IgM level as predictor of symptomatic hyperviscosity in patients with Waldenstrom macroglobulinaemia. *Br J Haematol* 177:717–725
 72. Weaver A, Rubinstein S, Cornell RF (2020) Hyperviscosity syndrome in paraprotein secreting conditions including Waldenstrom macroglobulinemia. *Front Oncol* 10:815
 73. Debureau PE, Harel S, Parquet N, Lemiale V, Siguret V, Goubeau L, Morin F, Royer B, Cuccuini W, Elessa D, Theves F, Brignier AC, Azoulay E, Arnulf B, Talbot A (2022) Prognosis of hyperviscosity syndrome in newly diagnosed multiple myeloma in modern-era therapy: a real-life study. *Front Immunol* 13:1069360
 74. Fahey JL, Barth WF, Solomon A (1965) Serum hyperviscosity syndrome. *JAMA* 192:464–467
 75. Dumas G, Gabarre P, Bigne N, Maury E (2018) Hyperviscosity syndrome. *Intensive Care Med* 44:1151–1152
 76. Stone MJ (2009) Waldenstrom's macroglobulinemia: hyperviscosity syndrome and cryoglobulinemia. *Clin Lymphoma Myeloma* 9:97–99
 77. Bauer PR, Ostermann M, Russell L, Robba C, David S, Ferreyro BL, Cid J, Castro P, Juffermans NP, Montini L, Pirani T, Van De Louw A, Nielsen N, Wendon J, Brignier AC, Schetz M, Kielstein JT, Winters JL, Azoulay E, Nine I (2022) Plasma exchange in the intensive care unit: a narrative review. *Intensive Care Med* 48:1382–1396
 78. Gertz MA (2017) Waldenstrom macroglobulinemia: 2017 update on diagnosis, risk stratification, and management. *Am J Hematol* 92:209–217
 79. Bridoux F, Leung N, Belmouaz M, Royal V, Ronco P, Nasr SH, Ferman J, International K, Monoclonal Gammopathy Research G (2021) Management of acute kidney injury in symptomatic multiple myeloma. *Kidney Int* 99:570–580
 80. Hutchison CA, Cockwell P, Stringer S, Bradwell A, Cook M, Gertz MA, Dispenzieri A, Winters JL, Kumar S, Rajkumar SV, Kyle RA, Leung N (2011) Early reduction of serum-free light chains associates with renal recovery in myeloma kidney. *J Am Soc Nephrol* 22:1129–1136
 81. Sathick IJ, Drosou ME, Leung N (2019) Myeloma light chain cast nephropathy, a review. *J Nephrol* 32:189–198
 82. Perazella MA, Markowitz GS (2008) Bisphosphonate nephrotoxicity. *Kidney Int* 74:1385–1393
 83. Bridoux F, Arnulf B, Karlin L, Blin N, Rabot N, Macro M, Audard V, Belhadj K, Pegourie B, Gobert P, Cornec Le Gall E, Joly B, Karras A, Jaccard A, Augeul-Meunier K, Manier S, Royer B, Caillot D, Tiab M, Delbes S, Suarez F, Vigneau C, Caillard S, Arakelyan-Laboure N, Roos-Weil D, Chevret S, Ferman JP (2020) Randomized trial comparing double versus triple bortezomib-based regimen in patients with multiple myeloma and acute kidney injury due to cast nephropathy. *J Clin Oncol* 38:2647–2657
 84. Bayraktar UD, Warsch S, Pereira D (2011) High-dose glucocorticoids improve renal failure reversibility in patients with newly diagnosed multiple myeloma. *Am J Hematol* 86:224–227
 85. Dimopoulos MA, Roussou M, Gavriatopoulou M, Psimenou E, Ziogas D, Eleutherakis-Papaikovou E, Fotiou D, Migkou M, Kanellias N, Panagiotidis I, Ntalianis A, Papadopoulou E, Stamatelopoulos K, Manios E, Pamboukas C, Kontogiannis S, Terpos E, Kastritis E (2017) Cardiac and renal complications of carfilzomib in patients with multiple myeloma. *Blood Adv* 1:449–454
 86. Peigne V, Rusinova K, Karlin L, Darmon M, Ferman JP, Schlemmer B, Azoulay E (2009) Continued survival gains in recent years among critically ill myeloma patients. *Intensive Care Med* 35:512–518