

## Erratum to: No impact of severe varus deformity on clinical outcome after posterior stabilized total knee arthroplasty

Bum-Sik Lee · Sang-Jin Lee · Jong-Min Kim ·  
Dae-Hee Lee · Eun-Jong Cha · Seong-II Bin

Published online: 11 January 2011  
© Springer-Verlag 2010

### Erratum to: Knee Surg Sports Traumatol Arthrosc DOI 10.1007/s00167-010-1316-9

Unfortunately, we have given the information in Table 9 of our article incorrectly. Therefore, please find the corrected Table 9 here:

**Table 9** The thicknesses of polyethylene inlays (mm)

	Mild varus group	Severe varus group	<i>P</i> value
Overall <sup>a</sup>	13.9 ± 2.5	14.7 ± 2.8	0.049
Except of the augmentation needed cases <sup>b</sup>	13.8 ± 2.5	15.1 ± 2.7	0.004

<sup>a</sup> The mean difference was 0.8 mm

<sup>b</sup> The mean difference was 1.3 mm, which was thicker than overall comparison with more significant statistical difference

The online version of the original article can be found under doi:[10.1007/s00167-010-1316-9](https://doi.org/10.1007/s00167-010-1316-9).

B.-S. Lee · S.-J. Lee · J.-M. Kim · S.-I. Bin (✉)  
Department of Orthopaedic Surgery, Asan Medical Center,  
College of Medicine, Ulsan University, 388-1, Pungnap-2-dong,  
Songpa-gu, Seoul 138-736, South Korea  
e-mail: sabin@amc.seoul.kr

D.-H. Lee  
Department of Orthopaedic Surgery,  
Korea University Anam Hospital, College of Medicine,  
Korea University, Anam-dong 5-Ga, Seongbuk-gu,  
Seoul 136-705, South Korea

E.-J. Cha  
Department of Biomedical Engineering, College of Medicine,  
Chungbuk National University, Gaesin-dong, Heungdeok-gu,  
Cheongju 361-763, Chungcheongbuk-do, South Korea