Correction to: Alignment in the transverse plane, but not sagittal or coronal plane, affects the risk of recurrent patella dislocation

Shigeru Takagi¹ · Takashi Sato² · Satoshi Watanabe² · Osamu Tanifuji¹ · Tomoharu Mochizuki¹ · Go Omori³ · Naoto Endo¹

Published online: 18 December 2017

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2017

Correction to:

Knee Surgery, Sports Traumatology, Arthroscopy
<https://doi.org/10.1007/s00167-017-4806-1>

The author would like to correct the errors in the publication of the original article. The corrected detail is given below for your reading.

In the Results section, Second paragraph should read as:

The mean values and standard deviation of the alignment parameters are described in Table 2. No significant difference was shown in 3D flexion angle (n.s.) and 3D

hip–knee–ankle angle (n.s.) between the RPD group and the control group. No CID was shown in 3D flexion angle [95% confidence interval (CI) of the difference, -3.1° to 3.8°] and 3D hip–knee–ankle angle (95% CI of the difference, -0.7° to 2.0°) between the two groups (Table 2). The significant and clinically important differences between the two groups were shown in 3D femoral anteversion, 3D external tibial torsion, and 3D knee rotational angle (Table 2).

The original article was corrected.

The original article can be found online at <https://doi.org/10.1007/s00167-017-4806-1>.

✉ Shigeru Takagi
m01a045c-shigeru@hotmail.co.jp

¹ Division of Orthopaedic Surgery, Department of Regenerative and Transplant Medicine, Graduate School of Medicine and Dental Sciences, Niigata University, 1-754 Asahimachidori Cuo-ku, Niigata, Niigata 951-8510, Japan

² Niigata Medical Center, Niigata, Japan

³ Niigata University of Health and Welfare, Niigata, Japan