



Correction to: Does increased femoral antetorsion predispose to cartilage lesions of the patellofemoral joint?

Johannes Oppermann¹ · Jan Bredow¹ · Boris Wissussek² · Christian Karl Spies³ · Christoph Kolja Boese¹ · Shi-Min Chang⁴ · Peer Eysel¹ · Jens Dargel¹

Published online: 31 March 2018

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2018

Correction to:

Knee Surg Sports Traumatol Arthrosc
(2017) 25:2695–2701

<https://doi.org/10.1007/s00167-015-3625-5>

In the original article, one of the co-author's (Boris Wissussek) family name has been published incorrectly. The correct family name should be Wissussek.

The original article can be found online at <https://doi.org/10.1007/s00167-015-3625-5>.

✉ Johannes Oppermann
johannes.oppermann@uk-koeln.de

Jan Bredow
jan.bredow@uk-koeln.de

Boris Wissussek
boris-alexander.wissussek@cellitinnen.de

Christian Karl Spies
dr.christian.spies@vulpiusklinik.de

Christoph Kolja Boese
christoph.boese@uk-koeln.de

Shi-Min Chang
shiminchang11@aliyun.com

Peer Eysel
peer.eysel@uk-koeln.de

Jens Dargel
jens.dargel@uk-koeln.de

¹ Department of Orthopedics and Trauma Surgery, University Hospital of Cologne, Kerpener Strasse 62, 50937 Cologne, Germany

² Department for Orthopedic and Trauma Surgery, St. Vinzenz Hospital, Merheimer Straße 221-223, 50733 Cologne, Germany

³ Department of Hand Surgery, Vulpius Klinik, Vulpiusstraße 29, 74906 Bad Rappenau, Germany

⁴ Department of Orthopedic Surgery, Yangpu Hospital, Tongji University School of Medicine, 450 Tengyue Road, 200090 Shanghai, People's Republic of China