Correction to: Patients with an Achilles tendon re-rupture have long-term functional deficits and worse patient-reported outcome than primary ruptures

Olof Westin^{1,2} · Katarina Nilsson Helander^{1,2,3} · Karin Grävare Silbernagel^{1,5} · Kristian Samuelsson^{1,2} · Annelie Brorsson^{1,4} · Jón Karlsson^{1,2}

Published online: 17 May 2018

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2018

Correction to:

Knee Surgery, Sports Traumatology, Arthroscopy
<https://doi.org/10.1007/s00167-018-4952-0>

Unfortunately, the title of the article contained a mistake in the online publication of the article.

The correct title should be read as follows:

Patients with an Achilles tendon re-rupture have long-term functional deficits and worse patient-reported outcome than primary ruptures.

The original article can be found online at <https://doi.org/10.1007/s00167-018-4952-0>.

✉ Olof Westin
olof.westin@gmail.com

¹ Department of Orthopaedics, Institute of Clinical Sciences, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

² Department of Orthopaedics, Sahlgrenska University Hospital, Mölndal, Sweden

³ Hallands Sjukhus, Kungsbacka, Sweden

⁴ IFK Kliniken Rehab, Gothenburg, Sweden

⁵ Department of Physical Therapy, University of Delaware, Newark, DE, USA