Correction to: Surgeon experience with dynamic intraligamentary stabilization does not influence risk of failure

Philipp Henle¹ · Kathrin S. Bieri² · Janosch Haeberli¹ · Nele Arnout³ · Jan Victor³ · Mirco Herbort⁴ · Clemens Koesters⁴ · Stefan Eggli¹

Published online: 25 August 2018

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2018

Correction to:
Knee Surgery, Sports Traumatology, Arthroscopy
<https://doi.org/10.1007/s00167-018-4847-0>

Tegner score and the OR (Multivariable) value of Injury to surgery have been corrected.

The corrected Table 4 is given below:

The original version of this article unfortunately contains mistake in Table 4. The *p* value of the univariable Age and

Table 4 Uni- and multi-variable regression analysis of factors associated with revision ACL surgery

Variable	Univariable			Multivariable (<i>n</i> = 96)		
	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value
Categoric						
Designer surgeons	1.48	(0.42–6.01)	n.s	2.25	(0.45–14.25)	n.s
Male gender	2.60	(0.70–12.51)	n.s	2.24	(0.40–16.01)	n.s
Proximal tear	0.27	(0.07–1.18)	n.s	0.29	(0.05–1.69)	n.s
Adjuvant procedure	1.19	(0.32–4.25)	n.s	1.23	(0.24–6.01)	n.s
Continuous						
Age (years)	0.88	(0.78–0.97)	0.020	0.92	(0.80–1.02)	n.s
Tegner score (0–10)	1.55	(1.05–2.38)	0.032	1.45	(0.87–2.62)	n.s
Injury to surgery (days)	0.97	(0.84–1.11)	n.s	1.00	(0.85–1.18)	n.s

The original article can be found online at <https://doi.org/10.1007/s00167-018-4847-0>.

✉ Kathrin S. Bieri
kathrin.bieri@ispm.unibe.ch

¹ Department of Knee Surgery and Sports Traumatology, Sonnenhof Orthopaedic Center, Bern, Switzerland

² Swiss RDL, Institute of Social and Preventive Medicine, University of Bern, Finkenhubelweg 11, 3012 Bern, Switzerland

³ Ghent University Hospital, Ghent, Belgium

⁴ University Hospital Muenster, Westphalian-Wilhelms University, Muenster, Germany