Correction to: Femoral and tibial bone bruise volume is not correlated with ALL injury or rotational instability in patients with ACL-deficient knee

Vincent Marot¹ · Boris Corin¹ · Nicolas Reina¹ · Jérôme Murgier⁴ · Emilie Berard² · Etienne Cavaignac^{1,3}

Published online: 28 May 2020

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2020

Correction to:

Knee Surgery, Sports Traumatology, Arthroscopy
<https://doi.org/10.1007/s00167-020-06045-x>

The affiliation of author Jérôme Murgier was published incorrectly in the original article. His correct affiliation is:

Aguiléra Private Clinic, Ramsey Générale de Santé, Biarritz, France

ORCID ID: <https://orcid.org/0000-0002-9237-1372>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00167-020-06045-x>.

✉ Etienne Cavaignac
cavaignac.etienne@gmail.com

¹ Department of Orthopedic Surgery and Trauma, Pierre Paul Riquet University Hospital, Toulouse, France

² Department of Epidemiology, Health Economics and Public Health, UMR1027 INSERM-University of Toulouse III, Toulouse University Hospital, Toulouse, France

³ Department of Orthopedic Surgery, Hôpital Pierre Paul Riquet, CHU Toulouse, Rue Jean Dausset, 3105 Toulouse, France

⁴ Aguiléra Private Clinic, Ramsey Générale de Santé, Biarritz, France