




Correction to: Calculating glenoid bone loss based on glenoid height using ipsilateral three-dimensional computed tomography

Johnny Rayes¹ · Jian Xu¹ · Sara Sparavalo¹ · Jie Ma¹ · Lauren Jonah¹ · Ivan Wong^{1,2} 

Published online: 6 July 2022

© The Author(s) under exclusive licence to European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2022

Correction to:

Knee Surgery, Sports Traumatology, Arthroscopy
<https://doi.org/10.1007/s00167-022-07020-4>

Unfortunately, the figures were incorrectly located in the original publication. Authors would like to update the figures.

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00167-022-07020-4>.

✉ Ivan Wong
research@drivanwong.com

Johnny Rayes
dr_rayes@hotmail.com

Jian Xu
12111220048@fudan.edu.cn

Sara Sparavalo
sara@drivanwong.com

Jie Ma
jie@drivanwong.com

Lauren Jonah
lauren@drivanwong.com

¹ Division of Orthopaedic Surgery, Faculty of Medicine, Dalhousie University, Halifax, NS, Canada

² 5955 Veteran's Memorial Lane, Room 2106 VMB, Halifax, NS B3H 2E1, Canada