Transforming ellipsoidal heights and geoid undulations between different geodetic reference frames

Christopher Kotsakis

Published online: 4 September 2007
© Springer-Verlag 2007

Erratum to: J Geod DOI 10.1007/s00190-007-0174-9

The original version of this article unfortunately contained a few minor mistakes:

- 1) Right after Eq. (6), the symbol ‘M’ should NOT be in italics.
- 2) At the left hand-side of Eqs. (18) and (19), the symbol ‘N’ should be in italics.
- 3) In Table 1 the following corrections occurred:
 - ‘GRF trtranslation terms’ should be changed to ‘GRF translation terms’
 - there should be an empty line before the ‘GRF translation terms’
 - there should not be an empty line between the equations $\delta h(t_x) = \dots$ and $\delta h(t_y) = \dots$
 - there should be an empty line before the ‘GRF orientation terms’
 - there should not be an empty line between the equations $\delta h(\varepsilon_x) = \dots$ and $\delta h(\varepsilon_y) = \dots$
 - there should be an empty line before the ‘GRF scaling term’
 - there should not be an empty line between the equations $\delta h(\delta a) = \dots$ and $\delta h(\delta f) = \dots$

The online version of the original article can be found under doi:[10.1007/s00190-007-0174-9](https://doi.org/10.1007/s00190-007-0174-9).

C. Kotsakis (✉)
Department of Geodesy and Surveying,
Aristotle University of Thessaloniki,
University Box 440, 54124 Thessaloniki, Greece
e-mail: kotsaki@topo.auth.gr