ERRATUM

Erratum to: Effect of vaginal estrogen on pessary use

Sybil G. Dessie^{1,2,3} • Katherine Armstrong^{2,3} • Anna M. Modest² •
Michele R. Hacker^{2,3} • Lekha S. Hota^{1,2,3,4}

Published online: 20 June 2016

© The International Urogynecological Association 2016

Erratum to: Int Urogynecol J
DOI 10.1007/s00192-016-3000-1

In the published version of this article, Figs. 2 and 3 captions were interchanged. These figures are presented again below with the correct captions.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00192-016-3000-1>.

✉ Lekha S. Hota
lhota@mah.harvard.edu

¹ Division of Urogynecology, Department of Obstetrics and Gynecology, Mount Auburn Hospital, Cambridge, MA, USA

² Department of Obstetrics and Gynecology, Beth Israel Deaconess Medical Center, Boston, MA, USA

³ Department of Obstetrics, Gynecology and Reproductive Biology, Harvard Medical School, Boston, MA, USA

⁴ Boston Urogynecology Associates/Mount Auburn Hospital, 725 Concord Ave. Suite #1200, Cambridge, MA 02138, USA

Fig. 2 Time to vaginal bleed among women with pessary use ($n = 199$); log rank test $P = 0.53$

Fig. 3 Increased vaginal discharge among women with pessary use ($n = 199$), log rank test, $P = 0.0001$