

Erratum to: Surgeons' views on sling tensioning during surgery for female stress urinary incontinence

Ali Borazjani¹ · Javier Pizarro-Berdichevsky^{2,3,4} · Jianbo Li⁵ · Howard B. Goldman²

Published online: 12 May 2017
© The International Urogynecological Association 2017

Erratum to: Int Urogynecol J
DOI 10.1007/s00192-017-3298-3

Due to a publisher's error, the original version of this article was published online with an incorrectly transposed Table 1. The corrected table is provided below. Please also note that the original article was corrected.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00192-017-3298-3>

✉ Ali Borazjani
borazjani.ali@gmail.com

- ¹ Lerner Research Institute, Cleveland Clinic, 9500 Euclid Avenue (ND20), Cleveland, OH 44195, USA
- ² Glickman Kidney and Urologic Institute, Cleveland Clinic, Cleveland, OH 44195, USA
- ³ Urogynecology Unit, H. Dr. Sotero del Rio, Santiago, Chile
- ⁴ Division Obstetricia y Ginecología, Pontificia Universidad Católica de Chile, Santiago, Chile
- ⁵ Quantitative Health Sciences, Cleveland Clinic, Cleveland, OH 44195, USA

Table 1 Demographics of the respondents

Parameter	Value	Total ⁿ ^a
Age, mean (SD)	48.1 (9.6)	596
Regions, % (<i>n</i>)		596
North America	37.1 (221)	
Europe	29 (173)	
South America	18.8 (112)	
Asia	2.7 (16)	
Central America	2.7 (16)	
Oceania	5.4 (32)	
Middle East	2.3 (14)	
Africa	2 (12)	
Practice Setting, % (<i>n</i>) ^b		596
University-based academic practice	38.6 (230)	
Multispecialty group practice	9.6 (57)	
Single specialty group practice	15.3 (91)	
Public hospital practice	31.5 (188)	
Private hospital practice	22 (131)	
Primary Specialty, % (<i>n</i>) ^c		596
Gynecology	63.1 (376)	
Urogynecology	54.0 (322)	
Gynecology	23.3 (139)	
Obstetrics & Gynecology	12.1 (72)	
Urology	36.6 (218)	
Female Urology	21.5 (128)	
Urology	20.5 (122)	
Other	0.33 (2)	
Years since residency, % (<i>n</i>)		596
0 to 5 years	18 (107)	
6 to 10 years	17.8 (106)	
11 to 20 years	31.2 (186)	
Greater than 20 years	33.1 (197)	
Fellowship Trained, % (<i>n</i>)	56.5 (337)	596
Percentage of practice devoted to PFDs, median (IQR)	60 (40–80)	596
Annual # of sling operations, % (<i>n</i>)		585
0	0.5 (3)	
1 to 10	6.7 (39)	
11 to 20	15 (88)	
21 to 30	18.5 (108)	
31 to 40	15.9 (93)	
41 to 50	12.3 (72)	
Greater than 50	31.1 (182)	
Annual number of sling revisions performed for obstructive complications, % (N)		584
0	24.5 (143)	
1 to 5	61 (356)	
6 to 10	7.2 (42)	
11 to 15	2.9 (17)	
16 to 20	1.7 (10)	
Greater than 20	2.7 (16)	

Table 1 (continued)

Parameter	Value	Total ⁿ ^a
Annual number of sling revisions performed for other complications, % (N)		585
0	19.5 (1)	
1 to 5	61 (357)	
6 to 10	11.1 (65)	
11 to 20	4.8 (28)	
Greater than 20	3.6 (21)	
Estimated percentage of own patients undergoing sling revisions by outside surgeons, % (<i>n</i>)		585
0 to 5%	85.5 (500)	
6 to 10%	13 (76)	
11 to 25%	1.2 (7)	
Greater than 25%	0.3 (2)	

^a Indicates the total number of respondents that completed the question

^b Respondents were allowed to choose one, or if they practiced in two or more settings equally, two that are most applicable. As such, percentages do not add up to 100% or the specified number of participants

^c Respondents were allowed to select all specialties that applied to them. Those who selected obstetrics & gynecology, urogynecology, and/or gynecology but not urology and/or female urology were considered gynecologists. Similarly, those who selected urology and/or female urology but not obstetrics & gynecology and/or gynecology were considered urologists. Two respondents only selected "other" as their primary specialty