Correction to: Patient attitudes toward pooled surgical waitlists in urogynecology

Rebekah A. Zee¹ · Aisling A. Clancy¹ · Hisham Khalil¹

Published online: 7 December 2019

© The International Urogynecological Association 2019

Correction to: International Urogynecology Journal

<https://doi.org/10.1007/s00192-019-04050-4>

Table 3 in the originally published article contains layout error. Corrected Table 3 shown below.

The online version of the original article can be found at <https://doi.org/10.1007/s00192-019-04050-4>

✉ Hisham Khalil
hkhalil@toh.ca

¹ Division of Urogynecology, Department of Obstetrics and Gynecology, The Ottawa Hospital, 1967 Riverside Dr, Ottawa, ON K1H 1A2, Canada

Table 3 Patient characteristics based on acceptance of a pooled surgical waitlist

	Not accepting of a pooled surgical waitlist (<i>n</i> = 113)	Accepting of a pooled surgical wait list (<i>n</i> = 60)	<i>P</i>
Age, years			0.071
25–44	22 (19.5%)	7 (11.9%)	
45–64	50 (44.2%)	20 (33.9%)	
≥65	41 (36.3%)	32 (54.2%)	
Surgery			0.85
Vag hyst ± prolapse repair ± MUS	30 (26.5%)	14 (23.3%)	
Vaginal prolapse repair ± MUS	44 (38.9%)	25 (41.7%)	
MUS alone	29 (25.7%)	15 (25.0%)	
Sacrocopexy	4 (3.5%)	3 (5.0%)	
Botox bladder injection	2 (1.8%)	2 (3.3%)	
Other	3 (2.7%)	0 (0.0%)	
Self-reported symptom severity			0.84
None	1 (0.9%)	0 (0.0%)	
Mild	2 (1.8%)	1 (1.7%)	
Moderate	52 (46.0%)	26 (43.3%)	
Severe	39 (34.5%)	25 (41.7%)	
Unsure/unknown	19 (16.8%)	8 (13.3%)	
Patient-reported acceptable wait time for surgery			0.041
≤2 months	28 (24.8%)	29 (49.2%)	
2–6 months	69 (61.1%)	25 (42.4%)	
6–12 months	13 (11.5%)	4 (6.8%)	
12–18 months	1 (0.9%)	0 (0.0%)	
18 months–2 years	1 (0.9%)	0 (0.0%)	
Doesn't matter	1 (0.9%)	1 (1.7%)	

MUS mid-urethral sling; *vag hyst* vaginal hysterectomy

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.