



Correction to: Preoperative level II/III MRI measures predicting long-term prolapse recurrence after native tissue repair

Payton Schmidt¹ · Luyun Chen^{1,2} · John O. DeLancey¹ · Carolyn W. Swenson¹

Published online: 22 September 2022

© The International Urogynecological Association 2022

Correction to: International Urogynecology Journal (2022) 33:133–141
<https://doi.org/10.1007/s00192-021-04854-3>

Figure 3 caption of the originally published article had an error. The word “Resting” needs to be changed to “Straining”. The correct Figure 3 caption is presented below:

Fig. 3 Straining preoperative MRI measurements compared to normal range in women with long-term success versus recurrence. The bottom error bar indicates the minimum value to the 25th percentile, and the top error bar indicates the 75th percentile to the maximum value

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00192-021-04854-3>.

✉ Payton Schmidt
payton@med.umich.edu

¹ Department of Obstetrics and Gynecology, University of Michigan, 1500 E. Medical Center Dr., Ann Arbor, MI 48109, USA

² Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI 48109, USA