

Erratum to: Association between fat mass, lean mass, and bone loss: the Dubbo osteoporosis epidemiology study

S. Yang¹ · J. R. Center^{1,2} · J. A. Eisman^{1,2,3} · T. V. Nguyen^{1,4,5}


Published online: 24 March 2015

© International Osteoporosis Foundation and National Osteoporosis Foundation 2015

Erratum to: Osteoporos Int DOI 10.1007/s00198-014-3009-6

Owing to errors in typesetting, this article was published with an incorrect version of Fig. 1; moreover,

Table 3 was poorly formatted. The correct Fig. 1 and Table 3 are given here.



The online version of the original article can be found at <http://dx.doi.org/10.1007/s00198-014-3009-6>.

✉ T. V. Nguyen
t.nguyen@garvan.org.au

¹ Osteoporosis and Bone Biology Research Program, Garvan Institute of Medical Research, 384 Victoria Street, Darlinghurst, NSW 2010, Australia

² Department of Endocrinology, St Vincent's Hospital, Sydney, Australia

³ School of Medicine, The University of Notre Dame Australia, Fremantle, Australia

⁴ School of Public Health & Community Medicine, Faculty of Medicine, University of New South Wales, Sydney, Australia

⁵ University of Technology, Sydney, Australia

Table 3 Association between fat mass, lean mass, and relative rate of femoral neck and lumbar spine BMD change for men and women: results of multiple linear regression analysis

Adjusted variables	Variables in the model	Per unit	Men				Women			
			Femoral neck BMD change (%/year)		Lumbar spine BMD change (%/year)		Femoral neck BMD change (%/year)		Lumbar spine BMD change (%/year)	
			Estimate (SE)	<i>P</i> value	Estimate (SE)	<i>P</i> value	Estimate (SE)	<i>P</i> value	Estimate (SE)	<i>P</i> value
Unadjusted	Fat mass (kg)	5	0.36 (0.20)	0.07	−0.11 (0.19)	0.57	−0.15 (0.11)	0.19	<i>0.42 (0.14)</i>	<i>0.003</i>
Unadjusted	Lean mass (kg)	5	0.36 (0.20)	0.08	0.28 (0.19)	0.14	−0.22 (0.22)	0.32	0.45 (0.28)	0.11
Unadjusted	Fat mass (kg)	5	0.26 (0.22)	0.25	−0.28 (0.21)	0.17	−0.12 (0.13)	0.33	<i>0.40 (0.16)</i>	<i>0.012</i>
	Lean mass (kg)	5	0.25 (0.23)	0.28	0.40 (0.21)	0.06	−0.11 (0.25)	0.66	0.09 (0.31)	0.78
Adjusted for age, height, history of prior fracture, smoking, osteoarthritis and physical activity	Fat mass (kg)	5	0.22 (0.22)	0.33	−0.26 (0.21)	0.21	−0.13 (0.13)	0.34	<i>0.36 (0.17)</i>	<i>0.03</i>
	Lean mass (kg)	5	0.20 (0.30)	0.51	0.47 (0.28)	0.09	−0.17 (0.29)	0.56	0.13 (0.37)	0.73

Statistical significance (at 5 % level) are shown in italics