



## Correction to: The factor-of-risk biomechanical approach predicts hip fracture in men and women: the Framingham Study

A.B. Dufour<sup>1,2</sup> · B. Roberts<sup>3</sup> · K.E. Broe<sup>2</sup> · D.P. Kiel<sup>2,4,5</sup> · M.L. Bouxsein<sup>4,3</sup> · M.T. Hannan<sup>2,4,5</sup>

Published online: 1 July 2019

© International Osteoporosis Foundation and National Osteoporosis Foundation 2019

### Correction to: Osteoporosis International

<https://doi.org/10.1007/s00198-011-1569-2>

The original version of this article, published 23 February 2011, unfortunately contained a mistake. The following correction has therefore been made in the original:

In section “Factor-of-risk,” a sentence was worded incorrectly. The correct wording is: “For women, trochanteric soft tissue thickness (millimeters) is equal to  $3.4795 \times [\text{BMI (kilograms per square meter)}] - 38.015$ . For men, trochanteric soft tissue thickness (millimeters) is equal to  $2.3415 \times [\text{BMI (kilograms per square meter)}] - 33.444$ .”

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00198-011-1569-2>

✉ A.B. Dufour  
[alyssadufour@hsl.harvard.edu](mailto:alyssadufour@hsl.harvard.edu)

B. Roberts  
[bjrobert@bidmc.harvard.edu](mailto:bjrobert@bidmc.harvard.edu)

K.E. Broe  
[broe@hsl.harvard.edu](mailto:broe@hsl.harvard.edu)

D.P. Kiel  
[kiel@hsl.harvard.edu](mailto:kiel@hsl.harvard.edu)

M.L. Bouxsein  
[mbouxsei@bidmc.harvard.edu](mailto:mbouxsei@bidmc.harvard.edu)

M.T. Hannan  
[hannan@hsl.harvard.edu](mailto:hannan@hsl.harvard.edu)

- <sup>1</sup> Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA
- <sup>2</sup> Institute for Aging Research, Hebrew SeniorLife, 1200 Centre St, Boston, MA 02131, USA
- <sup>3</sup> Orthopedic Biomechanics Laboratory, Beth Israel Deaconess Medical Center, Boston, MA, USA
- <sup>4</sup> Harvard Medical School, Boston, MA, USA
- <sup>5</sup> Division of Gerontology, Department of Medicine, Beth Israel Deaconess Medical Center, Boston, MA, USA