



Correction to: The efficacy and safety of menatetrenone in the management of osteoporosis: a systematic review and meta-analysis of randomized controlled trials

S. Su^{1,2} · N. He^{1,2} · P. Men¹ · C. Song³ · S. Zhai¹

Published online: 27 August 2021

© International Osteoporosis Foundation and National Osteoporosis Foundation 2021

Correction to: Osteoporosis International (2019) 30:1175–1186
<https://doi.org/10.1007/s00198-019-04853-7>

The original version of this article, published on February 7, 2019, contained a mistake.

The correct information is given below. The original article has been corrected.

Results - *Efficacy evaluations - Menatetrenone versus placebo/no additional drug:*

“In addition, analysis of **four** studies [22,24,26,38] (N=356) revealed a significantly greater percent increase in lumbar BMD from baseline for menatetrenone compared with placebo or no additional anti-osteoporotic drug (MD=2.02%, 95% CI 1.05 to 2.99%, P <0.0001) (Fig. 3 and Online Resource 3).”

Table 1. *Characteristics of the included RCTs*

In the “Outcome” column, the Orimo 1999 study should be “9,12,13” rather than “5,9,12,13”.

Correction in supplementary material

The online version of the original article can be found at <https://doi.org/10.1007/s00198-019-04853-7>

✉ S. Zhai
zhaisuodi@163.com

¹ Department of Pharmacy, Peking University Third Hospital, 49 Huayuan North Road, Haidian District, Beijing 100191, China

² Department of Pharmacy Administration and Clinical Pharmacy, School of Pharmaceutical Science, Peking University, Beijing, China

³ Department of Orthopaedics, Peking University Third Hospital, Beijing, China




Fig. 3 Summary of results of the meta-analyses for continuous variables (only summary effect estimate of menatetrenone vs. placebo – LBMD change (%) was changed)

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.