Correction to: A randomized, controlled trial of once-weekly teriparatide injection versus alendronate in patients at high risk of osteoporotic fracture: primary results of the Japanese osteoporosis intervention Trial-05

H. Hagino¹ • T. Sugimoto² • S. Tanaka³ • K. Sasaki³ • T. Sone⁴ • T. Nakamura⁵ • S. Soen⁶ • S. Mori⁷

Published online: 27 August 2021

© International Osteoporosis Foundation and National Osteoporosis Foundation 2021

Correction to: Osteoporosis Int

<https://doi.org/10.1007/s00198-021-05996-2>

The original version of this article, published on 17 May 2021, unfortunately contained a mistake.

The article was published with incomplete author list. The Author K. Sasaki was missing due to a typesetting mistake.

The publisher apologizes for this mistake.

The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00198-021-05996-2>

✉ S. Mori
stmori@sis.seirei.or.jp

H. Hagino
hagino@tottori-u.ac.jp

T. Sugimoto
sugimoto@med.shimane-u.ac.jp

S. Tanaka
tanaka.shiro.8n@kyoto-u.ac.jp

K. Sasaki
s.kotaro79@gmail.com

T. Sone
tsone@med.kawasaki-m.ac.jp

T. Nakamura
t-nak@utopia.ocn.ne.jp

S. Soen
nra48207@nifty.com

- ¹ School of Health Science, Tottori University Faculty of Medicine, Yonago, Tottori, Japan
- ² Eikokai Ono Hospital, 973 Tenjin-cho, Ono, Hyogo, Japan
- ³ Department of Clinical Biostatistics, Graduate School of Medicine, Kyoto University, Kyoto, Japan
- ⁴ Department of Nuclear Medicine, Kawasaki Medical School, Kurashiki, Okayama, Japan
- ⁵ Japan Osteoporosis Foundation, Tokyo, Japan
- ⁶ Soen Orthopaedics, Osteoporosis and Rheumatology Clinic, Kobe, Hyogo, Japan
- ⁷ Bone and Joint Surgery, Seirei Hamamatsu General Hospital, Hamamatsu, Shizuoka, Japan