



# Morphological and molecular comparison as a useful tool for identification of the three centric marine diatoms (Bacillariophyceae: Chaetoceros)

Maliwan Kutako<sup>1</sup> · Rachanimuk Hiransuchalert<sup>1,2</sup> · Sarayut Watchasit<sup>3</sup> · Mookthida Kaewduang<sup>1</sup> · Orawan Hanchana<sup>1</sup> · Pakawan Setthamongkol<sup>1</sup> · Parinya Chindudsadeegul<sup>4</sup> · Vichaya Gunbua<sup>5</sup> · Somtawin Jaritkhuan<sup>5</sup>

Received: 13 January 2023 / Revised: 17 March 2023 / Accepted: 28 March 2023 / Published online: 5 April 2023  
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

## Abstract

The objective of this study was to identify morphological and molecular comparison of three marine *Chaetoceros* species using microscopic observations, sequence analysis of 18S rDNA, random amplified polymorphic DNA (RAPD-PCR) barcoding and nuclear magnetic resonance (NMR) spectroscopy. *Chaetoceros* were obtained from three different algae laboratories: Center of Excellence for Marine Biotechnology (CEMB), Chanthaburi Coastal Fisheries Research and Development (CHAN) and Institute of Marine Science, Burapha University (BIM). Genomic DNA for the RAPD-PCR analysis was extracted using the phenol–chloroform method, followed by 18S rDNA amplification. The blast results of 18S rDNA sequence confirmed the significantly matched to *C. gracilis* for *Chaetoceros* BIM and CHAN and *C. muelleri* for *Chaetoceros* CEMB (e-value = 0.0, identity = 99%). The RAPD-PCR results revealed differences in the three *Chaetoceros* isolates with polymorphisms between 30.43% and 60.00%, and *Chaetoceros* CEMB showed high polymorphic bands. Scanning electron microscopy revealed that *Chaetoceros* CEMB were larger and had larger setae compared to the other isolates ( $P < 0.05$ ). The results of the NMR characterization of metabolites were consistent with the results of the sequence and morphological analyses. The concentrations of several metabolites, including chlorophyll  $c_1$ , chlorophyll a, Myo-inositol, fucoxanthin, astaxanthin, lutein and zeaxanthin, were lower in *Chaetoceros* CEMB than in *Chaetoceros* BIM and CHAN. However, high concentrations of fatty acids, such as oleic acid, linoleic acid,  $\alpha$ -linolenic acid and arachidic acid, were observed in all isolates. Generally, the results of this study will aid future studies examining the diversity of *Chaetoceros* in various cultural environments.

**Keywords** Morphology · 18S rDNA sequence · NMR · RAPD-PCR · *Chaetoceros*

---

Communicated by Yusuf Akhter.

✉ Rachanimuk Hiransuchalert  
rachanimuk@buu.ac.th

<sup>1</sup> Marine Biotechnology Research Unit, Faculty of Marine Technology, Chanthaburi Campus, 57 Moo 1, Kamong, Thamai, Chanthaburi 22170, Thailand

<sup>2</sup> Faculty of Science, Burapha University, 169 Long-Hard Bangsaen Road, Saensook, Muang, Chon Buri 20131, Thailand

<sup>3</sup> Nuclear Magnetic Resonance Spectroscopic Laboratory, Science Innovation Facility, Faculty of Science, Burapha University, 169 Long-Hard Bangsaen Road, Saensook, Muang, Chon Buri 20131, Thailand

<sup>4</sup> Faculty of Gems, Burapha University, Chanthaburi Campus, 57 Moo 1, Kamong, Thamai, Chanthaburi 22170, Thailand

<sup>5</sup> Department of Aquatic Science, Faculty of Science, Burapha University, 169 Long-Hard Bangsaen Road, Saensook, Muang, Chon Buri 20131, Thailand

## Introduction

Ocean's diatoms are an important group of eukaryotic phytoplankton that dominate phytoplankton communities in upwelling regions and at high latitudes (Benoiston et al. 2017). Diatoms are commonly fed to the larvae of aquatic animals in the aquaculture industry. The main species of diatoms used as food in the shrimp aquaculture industry are *Chaetoceros*, *Thalassionema*, *Closterium* and *Thalassiosira*. The genus *Chaetoceros* is highly representative in the marine plankton all over the world, in terms of diversity, abundance and distribution. The genus includes more than 225 recognized species and about 376 validly published names (Rines and Theriot 2003; Guiry 2020). The genus *Chaetoceros* is characterized by the presence of setae, which are long and hollow silicate spine-like projections protruding from the valve surface (Assmy et al. 2008; Kooistra et al. 2010). It is an ecologically important genus of marine planktonic diatoms that are found in coastal and upwelling regions (Jensen and Moestrup 1998). Moreover, *Chaetoceros* has been used for biofuel production because of its high growth rates and high lipid yield (Spaulding and Edlund 2008). Their utility stems in part from their small size and their high n-3 polyunsaturated fatty acids (PUFA) content; these properties also make *Chaetoceros* an important source of lipids and fatty acids for marine fish, bivalves and crustaceans (Xu et al. 1993; Zhou et al. 2007). Among *Chaetoceros* species, *C. muelleri* and *C. gracilis* have been cultivated for use as food for *Litopenaeus vannamei* larvae (Sangha et al. 2000). Rotifers fed *C. gracilis* showed increased viability, larger size, and low ciliate contamination (Knu 2004). Freeze-dried *C. muelleri* is commercially available as feed for shrimp, sea cucumbers and oysters and can be used in green water techniques in fish larviculture (e.g., <https://algae.provi ron.com>). Previous research has indicated that *C. muelleri* shows antimicrobial activity against *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans* (Mendiola et al. 2007). *C. gracilis* has the ability to produce high-quality fatty acids for the lipid industry (Rika Partwi et al. 2009).

*Chaetoceros* cells have a silica cell wall in the form of two valves called the frustule. Microscopic identification of diatoms has been traditionally used based on morphological characters such as frustule shape (Rine and Hargraves, 1988). Although there are several detailed taxonomic descriptions of the microstructure of their silica frustules, diatoms are time-consuming and difficult to identify. New approaches (e.g. the use of combined morphological and molecular tools) have been recently used which have revealed a cryptic diversity of the genus (Gaonkar et al. 2018).

DNA barcoding is a fast, accurate and standardized method for species-level identification that uses short DNA sequences. DNA barcoding has become an effective tool for assessing global biodiversity patterns and permits non-taxonomic biologists to diagnose species challenging to identify (Siddall et al. 2009). Diatoms are an ideal model group for developing DNA barcoding methods that provide easy-to-use, standardized and fast identification tools. Nuclear 18S rRNA is the most widely used sequence that has been used in phylogenetic analyses of diatoms (Evans et al. 2007; Gaonkar et al. 2018). Random amplified polymorphic (RAPD) markers are also highly useful for the study of populations within species because of their low cost and the fact that they do not require large sample sizes to generate preliminary results (Godhe et al. 2006).

In this study, the morphological and molecular taxonomic characteristics of *Chaetoceros* isolated from the coast of Eastern Thailand were investigated through microscopic observations, DNA barcoding using RAPD-PCR techniques, and molecular phylogenetic approaches using 18S rDNA. In addition, we present NMR patterns as well as chemical information on *Chaetoceros*.

## Materials and methods

### Diatom propagation and culturing system

*Chaetoceros* were obtained from three different algae laboratories: the Center of Excellence for Marine Biotechnology (CEMB), Faculty of Science, Chulalongkorn University, which originally isolated *Chaetoceros* from natural sea water in Angsila, Chon Buri province (13°20'22.5" N and 100°55'26.5" E) (hereafter referred to as *Chaetoceros* CEMB); Institute of Marine Science, Burapha University (BIM), which originally isolated *Chaetoceros* from natural sea water in Bang Saen, Chon Buri (13°16'10.0" N and 100°55'20.6" E) (hereafter referred to as *Chaetoceros* BIM); and Chanthaburi Coastal Fisheries Research and Development (CHAN), Department of Fisheries, Ministry of Agriculture and cooperatives, which originally isolated *Chaetoceros* from natural sea water in Laemsing, Chanthaburi province (12°31'46.1" N and 102°03'02.0" E) (hereafter referred to as *Chaetoceros* CHAN).

Diatom culture procedures were carried out under sterile conditions to avoid contamination. The diatom samples from the laboratory collections were purified using the centrifugation and streaking method. The samples were washed 12 times by RO water (30 ppt salinity), followed by centrifugation at 2000 × g for 1–2 min, and the supernatants were discarded every round. Each diatom strain was then aseptically streaked onto a sterile plate containing F/2 (Guillard 1975) media with water at 30 ppt salinity and 1.2–1.5% agar. The

plate was incubated at 25 °C until diatom colonies appeared. A single colony was transferred into test tubes containing 10 mL of fresh media. The diatom propagation process was first carried out in a 100-mL flask, followed by 500 mL in F/2 (Guillard 1975) media with water at 30 ppt salinity. The room was maintained at 25 °C and under continuous illumination at a low light intensity (1000  $\mu$  mol m<sup>-2</sup> s<sup>-1</sup>). To accelerate growth, diatom cultures were vigorously aerated until they reached the stationary phase. Diatom cultures were then centrifuged at 12,000  $\times$ g for 1–2 min, and the supernatants were discarded. The cells were kept at – 20 °C until DNA extraction.

### Morphological observations

Monoclonal cultures of diatom strains were identified to the genus or species level by morphological features based on observations using a light and scanning electron microscope. For light microscopy, diatom cultures were treated with concentrated hydrochloric acids (1:1, sample:HCl). A mixture was washed with distilled water 5 times and leave to sediment for 2 h to prepare cleaned frustules. The slides were examined using light microscopy under a 100 $\times$  oil immersion objective lens (model BX53, Olympus). For scanning electron microscopy, diatom cells were placed on glass plates coated with 0.5% Alcian blue, fixed with 2.5% glutaraldehyde in 0.1 M phosphate buffer saline (PBS) solution for 1 h and then washed with distilled water. Cells were further fixed under dark conditions with 1% osmium tetroxide in 0.1 M PBS solution for 1 h and then washed with distilled water. The samples were dehydrated in a graded ethanol series (70%, 80%, 90%, 95% and 100%) and dried using the critical point drying method. Finally, the samples were mounted onto stubs and sputter-coated with 99.99% pure gold. The specimens were examined under a scanning electron microscope (LEO1450VP, Zeiss, Oberkochen, Germany) with a 205-nm resolution at 10 kV. Morphological comparisons followed the procedures described in previous studies. Cell size (length, width and setae length, in  $\mu$ M) measurements of *Chaetoceros* were taken.

### Genomic DNA extraction

Fifty nanograms of diatom cells were placed in a prechilled microcentrifuge tube containing 500  $\mu$ L of the extraction buffer (100 mM Tris–HCl, 100 mM EDTA, and 250 mM NaCl; pH 8.0). High molecular weight genomic DNA (gDNA) was extracted from the washed diatom cells using the phenol–chloroform–proteinase K method (Sambrook and Russell 2001). The DNA pellet was resuspended in 100  $\mu$ L of TE buffer (10 mM Tris–HCl, pH 8.0 and 0.1 mM EDTA). The DNA solution was incubated at 37 °C for 1–2 h and kept at 4 °C until used.

### RAPD-PCR analysis

Polymerase chain reaction (PCR) amplification was performed in 25- $\mu$ L volumes containing 50 ng of genomic DNA; 2.5  $\mu$ L of 10X *Taq* DNA polymerase buffer; 200  $\mu$ M each of dNTP; 0.2  $\mu$ M primer (UBC101, 5'-GCGCCT GGAG-3' and OPB01, 5'-GTTTCGCTCC-3') [the University of British-Columbia pool (University of British-Columbia, Canada) and Operon (Operon Technologies, Inc., USA)]; 2.0 mM MgCl<sub>2</sub>; and 1.0 unit of Dynazyme™ *Taq* DNA polymerase (FINNZYMES, Finland). PCR thermocycling conditions were as follows: initial denaturation at 94 °C for 3 min, followed by 40 cycles of denaturation at 94 °C for 15 s; annealing at 36 °C for 1 min and extension at 72 °C for 1.30 min; and final extension at 72 °C for 7 min. Twelve  $\mu$ L of PCR products were electrophoresed on a 2.0% (w/v) agarose gel using a 1-kb DNA ladder (Promega) for size comparison before being stored at – 20 °C.

### PCR amplification, gene cloning and sequencing

The 18S rDNA locus from the genomic DNA of each diatom was amplified using universal diatom 18S rDNA-specific primers (Ki et al. 2009) (forward AT18F01, 5'-TAC CTGGTTGATCCTGCCAGTAG-3' and reverse AT18R01, 5'-GCTTGATCCTTCTGCAGGTTACC-3'). PCR amplification was carried out in 25- $\mu$ L volumes containing 50 ng of genomic DNA; 2.5  $\mu$ L of 10X *Taq* DNA polymerase buffer; 200  $\mu$ M each of dNTP; 0.2  $\mu$ M of each primer; 2.0 mM MgCl<sub>2</sub>; and 1.0 unit of Dynazyme™ II Hot Start *Taq* DNA polymerase (FINNZYMES, Finland). PCR thermocycling conditions were as follows: initial denaturation at 94 °C for 3 min, followed by 35 cycles of denaturation at 94 °C for 30 s; annealing at 50 °C for 1 min and extension at 72 °C for 2.30 min; and final extension at 72 °C for 7 min. The PCR products were electrophoresed on 1.5% (w/v) agarose gel using a 1-kb DNA ladder (Promega) for size comparison before being stored at – 20 °C.

Successful amplification products were purified, cloned and unidirectionally sequenced (AITBiotech Pte. Ltd., Singapore). Nucleotide sequences were analyzed against molecular reference databases using the BLAST® algorithm (Basic Local Alignment Search Tool) ([www.ncbi.nlm.nih.gov/BLAST](http://www.ncbi.nlm.nih.gov/BLAST)); similarity was considered significant when the probability (E) value recovered was less than 10<sup>-4</sup> (Altschul et al. 1990).

### DNA sequence characteristics and phylogenetic analyses

Intraspecific variation in *Chaetoceros* was investigated by comparing the DNA similarities and genetic distances of 18S rDNA sequences. Multiple alignments were performed with

each dataset using the ClustalW algorithm (Thompson et al. 1994). The aligned sequences were trimmed at each end to the same length, and obvious base errors that were only found in single strands were manually removed. Identical positions of the aligned sequences were used. The corrected pairwise ( $p$ -) genetic distances were calculated using the Kimura 2-parameter model (MEGA 10.0, Tamura et al. 2007).

An 18S rDNA phylogeny was constructed using the unweighted pair group method with arithmetic mean and Maximum Likelihood algorithms (MEGA 10.0, Tamura et al. 2007) based on the Kimura two-parameter distance matrix; 1000 bootstrap replicates were performed to assess the reliability of the topology. The similarity of GenBank sequences and sequences obtained in our study were also compared. A total of 12 rDNA sequences from *Chaetoceros* were used in analyses. The 18S rDNA sequences were retrieved from eight *Chaetoceros* species (Supplementary Fig. 1).

### Sample preparation for NMR analysis

Approximately 10 mg of methanol crude extract was dissolved in 600  $\mu$ L of deuterated acetone, which included the internal standard tetramethylsilane (TMS). The mixture was sonicated for 5 min and centrifuged at  $1400\times g$  for 5 min. Next, 550  $\mu$ L of supernatant was collected with a pipette and placed in a 5-mm NMR tube. All experiments (1D and 2D experiments) were performed using a Bruker Avance III HD 400 MHz NMR spectrometer, equipped with a 5-mm BBFO probe (Double Resonance Broadband Observe with  $^{19}\text{F}$  probe) at 25 °C. The  $^1\text{H}$ -NMR spectra of *Chaetoceros* extract were collected using the following parameters: pulse program zg30; relaxation delay 1 s; pulse width 8.90  $\mu$ s; number of scans 64; sweep width 18 ppm; and center of spectrum 6.50 ppm. The  $^{13}\text{C}$ -NMR spectrum of *Chaetoceros* extract was collected using the following parameters: pulse program zgpg30; relaxation delay 2 s; pulse width 7.50  $\mu$ s; number of scans 8900; sweep width 240 ppm; and center of spectrum 100 ppm. Parameters of the 2D  $J$ -resolved NMR were as follows: pulse program jresgp-prqf; relaxation delay 2 s; 128 increments; 16 transients; sweep width 18 ppm; and center of spectrum for  $^1\text{H}$  was 8.0 ppm in both dimensions. The parameters for HMBC are as follows: pulse program hmbcetgpl3nd; relaxation delay 2 s; 512 increments; 84 transients; sweep width and center of spectrum for  $^1\text{H}$  were 20.0 and 9.0 ppm, respectively. Sweep width and center of spectrum for  $^{13}\text{C}$  were 240.0 and 100.0 ppm, respectively.

## Results and discussion

### Growth pattern of *Chaetoceros*

*Chaetoceros* CHAN had the greatest cell numbers between day 3 ( $183\times 10^4$  cells/mL) and day 6 ( $192\times 10^4$  cells/mL) during the culture period (Supplementary Fig. 2). The growth of *Chaetoceros* CHAN and *Chaetoceros* BIM exponentially increased during days 2–3, and *Chaetoceros* CEMB exponentially increased during days 3–5. At day 6, the numbers of cells of *Chaetoceros* CEMB and *Chaetoceros* BIM were  $168\times 10^4$  and  $150\times 10^4$  cells/mL, respectively.


### Morphological characterization of *Chaetoceros*

The most abundant phytoplankton in the Central Gulf of Thailand are diatoms and blue-green algae (Kajonwatanakul et al. 2008). In this study, *Chaetoceros* from this study were isolated from the Gulf of Thailand: *Chaetoceros* CEMB and *Chaetoceros* BIM were isolated from Chonburi province, and *Chaetoceros* CHAN was isolated from Chanthaburi province.

Light microscope images are shown in Figs. 1a–c. The sizes of the three *Chaetoceros* isolates were small (ca. 5  $\mu$ M). They were delicate and nearly square or rectangular in girdle view, with the perivalvar axis longer than the apical axis. Traditionally, the identification of diatoms at the species level has been based on morphological features determined with the aid of light microscopy, including the morphology of the colonies, the shape and dimensions of cells, the thickness and direction of setae, the number and shape of chloroplasts and the presence and morphology of resting spores. However, other features that can only be resolved with electron microscopy, such as the fine structure of valves and setae and the location and number of rimoportulae, are now considered important (Sunesen et al. 2008). To minimize possible misidentifications, we used both scanning electron microscopy and light microscopy.

Scanning electron microscopy shows that the cells of *Chaetoceros* CEMB, CHAN and BIM were usually solitary with flat or slightly convex valves (Figs. 1d–l). The setae are straight and narrow in diameter and arise from the poles of the cells. The surfaces of the frustules or cell walls were smooth and rectangular in girdle view. *Chaetoceros* CEMB cells were shallow rectangular, whereas those of *Chaetoceros* CHAN and *Chaetoceros* BIM were square to rectangular. Setae size ( $18.37\pm 7.41\ \mu\text{M}$ ) and transapical axis ( $4.66\pm 1.25\ \mu\text{M}$ ) were significantly higher in *Chaetoceros* CEMB than in *Chaetoceros* CHAN and





**Fig. 1** Light (a–c) and scanning electron (d–l) microscope images of *Chaetoceros* CEMB (a, d, g and j), *Chaetoceros* CHAN (b, e, h and k) and *Chaetoceros* BIM (c, f, i and l); the higher magnification shows the shape of entire cells (scale bars 5  $\mu$ M)

**Table 1** Morphological characters for differentiating the *Chaetoceros* isolates in this study

Character	<i>Chaetoceros</i> CEMB	<i>Chaetoceros</i> CHAN	<i>Chaetoceros</i> BIM
Cell shape	Shallow rectangular	Square to rectangular	Square to rectangular
Setae shape	Round	Round	Round
Setae size ( $\mu$ M)	11.56–28.93 (18.37 $\pm$ 7.41)*	10.25–13.58 (12.01 $\pm$ 1.50)	10.27–11.14 (10.62 $\pm$ 0.36)
Transapical axis ( $\mu$ M)	3.24–6.16 (4.66 $\pm$ 1.25)*	3.50–4.05 (3.70 $\pm$ 0.25)	3.27–3.90 (3.53 $\pm$ 0.25)
Apical axis ( $\mu$ M)	4.15–8.26 (6.29 $\pm$ 1.50)	4.77–6.18 (5.33 $\pm$ 0.58)	4.42–5.49 (4.90 $\pm$ 0.49)

\*indicates statistically significant differences between isolates ( $p < 0.05$ )

*Chaetoceros* BIM (Table 1). *Chaetoceros* is a centric diatom with lightly silicified frustules. Each frustule possesses four long, thin spines or setae. The setae link the frustules together to form colonies of several cells. Frustules can usually be seen in girdle view. Distinguishing *Chaetoceros* species is difficult using a light microscope. The form of the chains, the sizes of the apical axis and valve shape are some of the most important morphological characters for recognizing species in this genus (Lee et al. 2014). All *Chaetoceros* isolates were confirmed to be *Chaetoceros* based on observation of their morphological features with a scanning electron microscope (Table 2).

Cells of *C. debilis* are roughly rectangular in girdle view and connected in spiraling chains. The basal part of the setae is distinct, and setae extend outward from the spiral. Valves are flat or slightly convex (although the spines make it appear concave). Apertures are narrowly oval and sometimes slightly constricted in the middle. Their diameter ranges from 8 to 40  $\mu\text{M}$ ; although they are distributed worldwide, they mainly occur in cold waters (Hasle and Syvertsen 1996). Species like *C. debilis* include *C. curvisetus*, which forms spirals wider in diameter; in addition, the apertures are larger and widely oval in *C. curvisetus* (Guiry and Guiry 2012; Tas and Hernández-Becerril 2017). The cells of *C. gracilis* and *C. muelleri* have an apical axis that ranges from 6 to 8 and 8–9  $\mu\text{M}$  and transapical axis that ranges from 3 to 6 and 6–7  $\mu\text{M}$ , respectively, while *C. calcitrans* has a cell diameter that ranges from 2 to 5  $\mu\text{M}$  (Olenina et al. 2006). *C. debilis* is larger than *C. gracilis* and *C. muelleri*. Light and scanning microscope observations showed that *Chaetoceros* CEMB may be a different species compared with *Chaetoceros* CHAN and *Chaetoceros* BIM. Nevertheless, light microscopy is still faster and more reliable for diatom identification in a mixed sample for trained diatomologists.

*C. gracilis* and *C. calcitrans* are extensively used as food sources for rearing prawn larvae (Seraspe et al. 2014). *C. gracilis* is the phytoplankton species most commonly used in bivalve mollusk and fish hatcheries (Helm et al. 2004). Their effectiveness stems in part from their small size and n-3 HUFA content.

## Nucleotide sequences and 18S rDNA phylogeny of *Chaetoceros*


The 18S rDNA sequences of *Chaetoceros* were obtained from gene cloning and unidirectional sequencing. *Chaetoceros* CEMB contained 18S rDNA sequences that were 1794 bp in length (Accession number MW513719.1), which were similar to those of *C. muelleri* (e-value = 0.0, identity = 99%). *Chaetoceros* CHAN had 18S rDNA sequences that were 1788 bp in length (Accession number MW513720.1), which were similar to those of *C. gracilis* (e-value = 0.0, identity = 99%). *Chaetoceros* BIM contained 18S rDNA sequences that were 1789 bp in length (Accession number MW513721.1), which were similar to those of *C. gracilis* (e-value = 0.0, identity = 99%). The 18S rDNA sequences of the three *Chaetoceros* showed high similarity (Supplementary Fig. 3).

The BLAST analysis revealed high similarity between the *Chaetoceros* sequences obtained in our study and GenBank sequences. We characterized partial nuclear 18S rDNA sequences of three *Chaetoceros* and compared them with available DNA sequences (12 sequences) obtained from GenBank ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) (Fig. 2). *Chaetoceros* CHAN and BIM were clustered in the same clade with *C. gracilis*, and *Chaetoceros* CEMB was distinct from the others. This result was consistent with morphological data suggesting that *Chaetoceros* CEMB contained significantly larger setae and apical axes than *Chaetoceros* CHAN and BIM. The lack of complete consistency between molecular and morphological identification may stem from morphological shifts that occur between environmental species and cultured ones. Thus, species identification both before and after culture might be required to ensure the accuracy of identification (Kesici et al. 2013).

*Chaetoceros* is a diverse genus of marine diatoms. Although the morphology of many members of the genus has been well described, molecular taxonomic studies of *Chaetoceros* are scarce. However, the use of new approaches by combination of morphological and molecular tools, have been recently used which have revealed a cryptic diversity

**Table 2** Comparison of the morphological features of *Chaetoceros* CEMB, CHAN and BIM by scanning electron microscopy

Characteristic	<i>Chaetoceros</i> CEMB	<i>Chaetoceros</i> CHAN	<i>Chaetoceros</i> BIM
Cell	Thick and stiff cells that are shallow rectangular	Thick and stiff cells that are rectangular	Thick and stiff cells that are rectangular
Valve face	Oval	Rectangular to oval	Rectangular to oval
Cell Length ( $\mu\text{M}$ )	4–8	4–6	4–6
Apical axis ( $\mu\text{M}$ )	3–6	3–4	3–4
Setae	4 long and thin intercalary setae with rounded terminal parts	4 long and thin intercalary setae with slender terminal parts	4 long and thin intercalary setae with slender terminal parts
Spines	Invisible spine	Invisible spine	Invisible spine



**Fig. 2** Partial sequences 18S rDNA Maximum Likelihood phylogenetic hypothesis inferred for the species of *Chaetoceros* CEMB (Accession number MW513719.1), CHAN (Accession number MW513720.1) and BIM (Accession number MW513721.1) (black

dots) and *Chaetoceros* 18S rDNA sequences (12 diatom taxa) from GenBank; 1000 bootstrap replicates were performed to assess the reliability of the topology. A partial sequence 18S rDNA of *Chlorella vulgaris* (Accession number X13688.1) was used as the outgroup

of the genus (Gaonkar et al. 2018). The 18S and 28S rDNA phylogenies might provide suitable markers for resolving the species-level taxonomy of *Chaetoceros* (Oh et al. 2010).


### RAPD profiles of *Chaetoceros*

Amplified fragments 300–2000 bp in size were obtained using RAPD-PCR analysis with UBC10 and OPB01 primers (Fig. 3). A dendrogram based on the RAPD-PCR band was created. In the study of the interpopulational variability of the three *Chaetoceros* culture populations, the selection of the RAPD primers was based on the quantity, intensity and repetition of the amplified fragments. These amplified fragments ranged in size from 50 to 2200 bp. A total of 80 fragments were identified, and 113 of these fragments (42.5%) were polymorphic. The average number of fragments per primer was relatively high. The percentage of polymorphic bands was 33.33%, 60.00% and 30.43% for *Chaetoceros* CHAN, *Chaetoceros* CEMB and *Chaetoceros* BIM, respectively (Table 3). DNA barcoding requires molecular loci that are variable enough to discriminate species and a molecular reference database for comparison. The

similarity or divergence of the molecular sequence of an unknown organism to a vouchered reference sequence in the database is used for species identification. DNA barcoding of environmental samples involves the extraction of DNA from a pooled sample, PCR amplification of a target locus, cloning of the resulting PCR products, sequencing and analysis. With DNA barcoding techniques, even morphologically similar strains can be identified at the species level. These molecular phylogenetic analyses also enable the rapid, convenient, and accurate classification of diatoms and have thus contributed considerably to studies of diatom diversity.

RAPD-PCR has been used for the molecular characterization and identification of 17 samples of *Sargassum* spp. (Ho et al. 1995). A 450-bp fragment generated using OPA13 was detected in 12 of 17 samples of *Sargassum*. This fragment was present in profiles from *Turbinaria* (Sargassaceae). This study showed that RAPD-PCR is useful for discriminating *Sargassum* samples and developing fingerprints for them. PCR–RFLP analysis has been used to resolve the species-level differences of 18 isolates of *Chaetoceros* Ehrenberg (Bacillariophyceae) by targeting the *rbcL* region of chloroplast DNA, which encodes the Rubisco large subunit

**Fig. 3** A 2.0% agarose gel showing RAPD patterns of *Chaetoceros* using UBC10 (left) and OPB01 (right). Lanes 1 and 2 *Chaetoceros* CHAN, Lanes 3 and 4 *Chaetoceros* CEMB, Lanes 5 and 6 *Chaetoceros* BIM; M is the 100 bp DNA Marker



**Table 3** Number of RAPD fragments and polymorphic products obtained in the analysis of three *Chaetoceros* populations

Primer	Total number of fragments	Number of amplified fragments			Number of polymorphic fragments		
		<i>Chaetoceros</i> CHAN	<i>Chaetoceros</i> CEMB	<i>Chaetoceros</i> BIM	<i>Chaetoceros</i> CHAN	<i>Chaetoceros</i> CEMB	<i>Chaetoceros</i> BIM
UBC10	39	14	14	11	5	6	2
OPB01	41	13	16	12	4	12	5
Total	80	27	30	23	9	18	7
Polymorphism					33.33%	60.00%	30.43%


(Toyoda et al. 2011). RAPD patterns for the species-level differences of *Chaetoceros* have not been reported to date. Molecular identification appears to be relatively effective for diatom identification given the similar efficacies of molecular and morphological identification in this study. However, more work is needed to optimize morphological and molecular approaches for diatom identification.

### Identification of metabolites extracted from *Chaetoceros* by NMR spectroscopy

The  $^1\text{H-NMR}$  spectra of methanol extract from all *Chaetoceros* isolates showed similar characteristic chemical shift peaks. There was a total of 27 metabolites that were clearly identified based on comparison with previous research, a free NMR database (The Human Metabolome Database, HMDB) and a commercial NMR database (Bruker AssureNMR). The  $^1\text{H-NMR}$  spectra shown in Fig. 4 contain different groups of metabolites, including amino acids, sugars, carboxylic acids, fatty acids, vitamins and carotenoids. The peaks corresponding to the structures of each metabolite are summarized in Supplementary data.

The characteristic chemical shifts of eight amino acids and sugars were observed around the region 4.10–1.98 ppm, which correspond to the  $-\text{CH}_2-$  protons of amino acids, and 1.48–0.96 ppm, which correspond to the  $-\text{CH}-$  and  $-\text{CH}_3$  protons of amino acids (Azizan et al. 2018; Ma et al. 2019; Iglesias et al. 2019). The peaks around 5.20–3.82 ppm correspond to the  $-\text{CH}-$  protons of glucose and sucrose, and the peaks around 3.82–3.67 ppm correspond to the  $-\text{CH}_2-$  protons of glucose and sucrose (Richter and Berger 2013). The representative proton signals of fucoxanthin (olefinic- $H$ ), astaxanthin, lutein and zeaxanthin were observed at a chemical shift around 7.01–6.10 ppm (Zailanie and Purnomo 2017; Shumilina et al. 2020; Otaka et al. 2016; Iwai et al. 2008). The identifications of these carotenoids have been confirmed by 2D-NMR (HMBC and JRES); the JRES spectrum showed the singlet signals of fucoxanthin and astaxanthin at 2.01 and 1.98 ppm, respectively, which correspond to the methyl groups (Supplementary Fig. 4) (Subramanian et al. 2015). The correlation between the proton and carbon signals in the HMBC spectrum is consistent with the results of previous studies (Azizan et al. 2018) (Supplementary Figs. 5–6). The signals of chlorophyll a and chlorophyll  $c_1$





**Fig. 4**  $^1\text{H-NMR}$  spectra of *Chaetoceros* methanol extract in acetone- $\text{D}_6$ : (a) *Chaetoceros* CEMB (b) *Chaetoceros* BIM and (c) *Chaetoceros* CHAN where 1, glutamate; 2, proline; 3, alanine; 4, isoleucine; 5, methionine; 6, choline; 7, glycine; 8, cholesterol; 9, palmitic acid; 10, oleic acid; 11, linolenic acid; 12,  $\alpha$ -linolenic acid; 13, arachidic

acid; 14, glucose; 15, sucrose; 16, myo-inositol; 17, fucoxanthin; 18, astaxanthin; 19, lutein; 20, zeaxanthin; 21, violaxanthin; 22, chlorophyll  $c_1$ ; 23, chlorophyll  $a$ ; 24, glutamine; 25, valine; 26, leucine; and 27, steric acid


observed around 9.77–8.20 ppm correspond to the  $-\text{NH}-$  protons of chlorophyll structures.

$^1\text{H-NMR}$  and 2D-NMR spectra of the crude extract revealed signals for six fatty acids, including palmitic acid, oleic acid, linoleic acid,  $\alpha$ -linolenic acid, arachidic acid and stearic acid (Fig. 5). The characteristic peaks were similar to the results of previous studies (Roswanda et al. 2017; Otto et al. 2014; Singer et al. 1996). The correlation between proton and carbon signals observed around 1.73–1.29 ppm corresponded to arachidic acid, palmitic acid and stearic acid. Other valuable metabolites, such as myo-inositol, cholesterol and choline, were detected by 1D and 2D-NMR spectroscopy. The results indicated that further purification was not required for the identification of some major and minor small metabolites by NMR spectroscopy.

Lipids play an important role in larval growth and survival. Eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) are considered essential fatty acids because they are integral components of the plasma membrane and marine fish larvae cannot synthesize them from linoleic acid 18:3 (n-3). *M. rosenbergii* also lacks the ability to synthesize linolenic acid and linoleic acid (D'Abramo and Sheen 1993) and has a limited ability to elongate and desaturate

short-chain n-3 and n-6 polyunsaturated fatty acids (e.g., C18) to long-chain polyunsaturated fatty acids (e.g., C20) (Reigh and Stickney 1989). Thus, marine fish larvae must acquire PUFAs through their diet of zooplankton (e.g., rotifers and crustaceans), which are enriched in these nutrients. Increasing the PUFA content of zooplankton before feeding larval fish and shrimp is a regular practice in the aquaculture industry (Apt and Behrens 1999).

The supply of EPA and DHA traditionally produced by marine fisheries will be insufficient to meet their market demand in food industry. Consequently, a sustainable alternative source is urgently required. Moreover, EPA and DHA as n-3 supplements can potentially be used as an adjuvant for cardiac issues associated with coronavirus disease 2019 (COVID-19) (Oliver et al. 2020). Through adaptive laboratory semi-continuous cultures condition, the smaller *C. gracilis* cells can accumulate relatively higher EPA (41.5% EPA content per total fatty acid) and fucoxanthin (Tachihana et al. 2020). In this study, unsaturated fatty acids and fucoxanthin in *Chaetoceros* were found. The results from this study suggesting that, high biomass production of unsaturated fatty acids and Fucoxanthin contents in *Chaetoceros* CHAN and BIM may achieve in semi-continuous culture at a



**Fig. 5** Expanded  $^1\text{H-NMR}$  spectra of *Chaetoceros* methanol extract with the peak area (ratio) compared with the 0.03% TMS peak area in acetone- $\text{D}_6$ : (a) *Chaetoceros* CEMB (b) *Chaetoceros* BIM and

(c) *Chaetoceros* CHAN where 10, oleic acid; 11, linolenic acid; 12,  $\alpha$ -linolenic acid; 13, arachidic acid; 22, chlorophyll  $c_1$ ; 23, chlorophyll  $a$ ; 24, glutamine; 25, valine; 26, leucine; and 27, steric acid

low dilution and providing sufficient nutrients, to obtain high biomass production and valuable bio-compounds contents.

## Conclusions

This study demonstrates the potential for DNA barcoding, coupled with microscopic observation and NMR characterization, for assessing *Chaetoceros* biodiversity. The RAPD barcodes, 18S rDNA sequences and NMR profiles of the three diatom isolates from this study can be used to identify *Chaetoceros* species when morphological differences are ambiguous.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00203-023-03525-9>.

**Acknowledgements** We thank the Center of Excellence for Marine Biotechnology (CEMB), Faculty of Science, Chulalongkorn University; Institute of Marine Science, Burapha University (BIM); and Chanthaburi Coastal Fisheries Research and Development (CHAN), Department of Fisheries, Ministry of Agriculture for providing *Chaetoceros*. We thank Mrs. Jantane Boonmameepool and the Microscopic

Center, Faculty of Science, Burapha University for help with scanning electron microscopy.

**Author contributions** MK, RH, VK, and SJ designed the experiments. MKa, OH, SW and PC conducted the experiments and carried out the analyses. PS, VK and SJ evaluated the data. RH and MK. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding** This research was financially supported by The Higher Education Commission of Thailand (code 1/2556 Burapha University) to RH.

**Data Availability** The data presented in the study are deposited in the GenBank. The accession number for 18S rRNA gene sequence of *Chaetoceros* CEMB, *Chaetoceros* CHAN and *Chaetoceros* BIM are MW513719.1, MW513720.1 and MW513721.1, respectively.

## Declarations

**Conflict of interest** The authors declare that they have no conflict of interest.

**Ethical approval** This research does not contain any studies with animals performed by any of the authors. This research was approved for biosafety by the Institutional Biosafety Committee of Burapha University Thailand.

## References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- Apt KE, Behrens PW (1999) Commercial developments in microalgal biotechnology. *J Phycol* 35:215–226
- Assmy P, Hernández-Becerril DU, Montresor M (2008) Morphological variability and life cycle traits of the type species of the diatom genus *Chaetoceros*. *C Dichaeta Journal of Phycology* 44(1):152–163
- Azizan A, Bustamam MSA, Maulidiani M, Shaari K, Ismail IS, Nagao N, Abas F (2018) Metabolite profiling of the microalgal diatom *Chaetoceros calcitrans* and correlation with antioxidant and nitric oxide inhibitory activities via <sup>1</sup>H NMR-based metabolomics. *Mar Drugs* 16:154
- Benoiston AS, Ibarbalz FM, Bittner L, Guidi L, Jahn O, Dutkiewicz S, Bowler C (2017) The evolution of diatoms and their biogeochemical functions. *Phil Trans R Soc B* 372:20160397
- D'Abramo, L.R. and Sheen, S.S. (1993). Polyunsaturated fatty acid nutrition in juvenile freshwater prawn *Macrobrachium rosenbergii*. *Aquaculture*. 115: 63e86.
- Evans KM, Wortley AH, Mann DG (2007) An Assessment of potential diatom “Barcode” genes (cox1, rbcL, 18S and ITS rDNA) and their effectiveness in determining relationships in *Sellaphora* (Bacillariophyta). *Protist* 158:349–364
- Gaonkar CC, Piredda R, Minucci C, Mann DG, Montresor M, Sarno D, Kooistra WHCF (2018) Annotated 18S and 28S rDNA reference sequences of taxa in the planktonic diatom family Chaetocerotaceae. *PLoS ONE* 13:e0208929
- Godhe A, McQuoid MR, Karunasagarb I, Karunasagarb I, Rehnstam-Holm NS (2006) Comparison of three common molecular tools for distinguishing among geographically separated clones of the diatom *Skeletonema marinoi* Sarno et Zingone (Bacillariophyceae). *J Phycol* 42:280–291
- Guillard, R. R. L. (1975). Culture of phytoplankton for feeding marine invertebrates. 26–60.
- Guiry, M. D. and Guiry, G. M. 2012. *Chaetoceros debilis* Cleve. [http://www.algaebase.org/search/species/detail/?species\\_id=37416](http://www.algaebase.org/search/species/detail/?species_id=37416). Accessed 18 Mar 2012.
- Guiry, M. D. in Guiry, M. D., and Guiry, G. M. (2020). *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>; searched on 15 May 2020.
- Hasle, G. R. and Syvertsen, E. E. (1996). Marine diatoms. (In: Tomas, C. R. ed.) *Identifying marine diatoms and dinoflagellates*. Academic Press, Inc., San Diego. 5–385.
- Helm, M. M., Bourne, N. and Lovatelli, A. (2004). *Hatchery Culture of Bivalves: A Practical Manual*. FAO Fisheries Technical Paper Number 471. Rome, Italy. 200.
- Ho SC, Lau J, Woo J (1995) Life satisfaction and associated factors in older Hong Kong Chinese. *Ame Geriatrics Soc* 43:252–255
- Iglesias, M.J., Soengas, R., Probert, I., Guilloud, E., Gourvil, P., Mehiri, M., Lópezd, Y., Cepas, V., Gutiérrez-del-Río, I., Redondo-B., S., Villar, C.J., Lombó, F. Soto, S. and Ortiza F.L. (2019). NMR characterization and evaluation of antibacterial and antibiofilm activity of organic extracts from stationary phase batch cultures of five marine microalgae (*Dunaliella* sp., *D. salina*, *Chaetoceros calcitrans*, *C. gracilis* and *Tisochrysis lutea*). *Phytochemistry*. 164: 192–205
- Iwai M, Maoka T, Ikeuchi M, Takaichi S (2008) 2,2'-β-Hydroxylase (CrtG) is involved in Carotenogenesis of Both Nostoxanthin and 2-Hydroxymyxol 2'-Fucoside in *Thermosynechococcus elongatus* Strain BP-1. *Plant Cell Physiol* 49(11):1678–1687
- Jensen KG, Moestrup Ø (1998) The genus *Chaetoceros* (Bacillariophyceae) in inner Danish waters. *Opera Botanica* 133:5–68
- Kajonwattanakul S, Numnual W, Sirichaiseth T, Wannarangsri T (2008) Survey of Marine phytoplankton in Ship's ballast tanks at laem Chabang International Port, Thailand ASEAN. *J Sci and Technology Development* 35(1–2):141–145
- Kesici K, Tüney I, Zeren D, Güden M, Sukata A (2013) Morphological and molecular identification of pennate diatoms isolated from Urla, İzmir, coast of the Aegean Sea. *Turk J Biol* 37:530–537
- Ki JS (2009) Comparative molecular analysis of freshwater centric diatoms with particular emphasis on the nuclear ribosomal DNA of *Stephanodiscus* (Bacillariophyceae). *Algae* 24:129–138
- Knu CV (2004) A comparison of yield and quality of the rotifer (*Brachionus plicatilis*-L-strain) fed different diets under aquaculture conditions. *Vietnam Asian Fisheries Science* 17:357–363
- Kooistra WHCF, Sarno D, Hernández-Becerril DU, Assmy P, Di Prisco C, Montresor M (2010) Comparative molecular and morphological phylogenetic analyses of taxa in the Chaetocerotaceae (Bacillariophyta). *Phycologia* 49(5):471–500
- Lee SD, Park JS, Yun SM, Lee JH (2014) Critical criteria for identification of the genus *Chaetoceros* (Bacillariophyta) based on setae ultrastructure. I Subgenus *Chaetoceros* *Phycologia* 53(2):174–187
- Ma NL, The KY, Ahmad A, Lam SL, Saidon SAB, Husri MN, San CT (2019) Metabolite profiling of *Scenedesmus regularis* using nuclear magnetic resonance (NMR). *Malaysian Appl Biol J* 48(1):117–121
- Mendiola JA, Torres CF, Toré A, Martín-Álvarez PJ, Santoyo S, Arredondo BO, Señoráns FJ, Cifuentes A, Ibáñez E (2007) Use of supercritical CO<sub>2</sub> to obtain extracts with antimicrobial activity from *Chaetoceros muelleri* microalga. a correlation with their lipidic content. *Europ Food Res Technol* 224:505–510
- Oh, H. Y., Cheon I, J. Y., Lee, J. H., Hur, S. B. and Ki, J. S. (2010). Nuclear rDNA characteristics for DNA taxonomy of the centric diatom *Chaetoceros* (Bacillariophyceae). *Algae*. 25(2): 65–70
- Olenina I, Hajdu S, Edler L, Andersson A, Wasmund N, Busch S, Göbel J, Gromisz S, Huseby S, Huttunen M, Jaanus A, Kokkonen P, Ledaine I, Niemkiewicz E (2006) Biovolumes and size-classes of phytoplankton in the Baltic Sea. *BSEP* 106:1–144
- Oliver L, Dietrich T, Marañón I, Villarán MC, Barrio RJ (2020) Producing omega-3 polyunsaturated fatty acids: a review of sustainable sources and future trends for the EPA and DHA market. *Resources* 9:148
- Otaka J, Seo S, Nishimura M (2016) Lutein, a natural carotenoid, induces α-1,3-glucan accumulation on the cell wall surface of fungal plant pathogens. *Molecules* 21:980
- Otsuka S, Suda S, Li R, Matsumoto S, Watanabe M (2000) Morphological variability of colonies of *Microcystis* morphospecies in culture. *J Gen Appl Microbiol* 46:39–50
- Otto JR, Freeman MJ, Malau-Aduli BS, Nichols PD, Lane PA, Malau-Aduli AEO (2014) Reproduction and fertility parameters of dairy cows supplemented with omega 3 fatty acid-rich canola oil. *Annual Rev Rev Biol* 4(10):1611–1636
- Reigh, R. C. and Stickney, R. R. (1989). Effects of purified dietary fatty acids on the fatty composition of freshwater shrimp *Macrobrachium rosenbergii*. *Aquaculture*. 157–174.
- Richter T, Berger S (2013) A NMR method to determine the anomeric specificity of glucose phosphorylation. *Bioorg Med Chem* 21:2710–2714
- Rika Partiwí A, Dahrul S, Linawati H, Lily Maria GP, Maggy TS (2009) Fatty acid synthesis by Indonesian marine diatom, *Chaetoceros gracilis*. *HAYATI J Biosci* 16(4):151–156
- Rines JEB, Hargraves PE (1988) The *Chaetoceros ehrenbergii* (Bacillariophyceae) flora of Narragansett Bay, Rhode Island, U.S.A. *Bibl Phycol* 79:1–196
- Rines JEB, Theriot EC (2003) Systematics of Chaetocerotaceae (Bacillariophyceae) I. a phylogenetic analysis of the family. *Phycol Res* 51:83–98

- Roswanda R, Putra IA, Mujahidin D (2017) Dimethyl 9-Octadecenoate and 9-olefin metathesis reaction. *Inter J Tech* 8(8):1422
- Sambrook, J. and Russell, D.W. (2001). *Molecular Cloning a Laboratory manual*. New York: Cold Spring Harbor.
- Sangha MK, Gupta PK, Thapar VK, Verma SR, Bal AS, Dixit A (2000) Agricultural mechanisation in Asia. Africa and Latin America (AMA). 13(1):54
- Seraspe EB, Gabotero S, Milagros R, Pahila IG, Amar EC (2014) Evaluation of dietary freeze-dried *Chaetoceros calcitrans* supplementation to control *Vibrio harveyi* infection on *Penaeus monodon* juvenile. *Aquaculture* 432:212–216
- Shumilina E, Ciampa A, Ytrestøl T, Dikiy A (2020) Assessment of astaxanthin accumulation in hepatocytes of atlantic salmon fed different diets using NMR spectroscopy. *Molecules* 25:1769
- Siddall ME, Fontanella FM, Watson SC, Kvist S, Erséus, C. (2009) Barcoding bamboozled by bacteria: convergence to metazoan mitochondrial primer targets by marine microbes. *Syst Biol* 58(4):445–451
- Singer S, Sivaraja M, Souza K, Millis K, Corson JM (1996) <sup>1</sup>H-NMR detectable fatty acyl chain unsaturation in excised Leiomyosarcoma correlate with grade and mitotic activity. *J Clin Investig* 98(2):244–250
- Spaulding, S. and Edlund, M. (2008). *Chaetoceros*. In *Diatoms of the United States*. Retrieved October. 19: 2017.
- Subramanian B, Thibault M-H, Djaoued Y, Pelletier C, Touaibia M, Tchoukanova N (2015) Chromatographic, NMR and vibrational spectroscopic investigations of astaxanthin esters: application to “Astaxanthin-rich shrimp oil” obtained from processing of Nordic shrimps. *Analyst* 140:7423–7433
- Sunesen I, Hernández-Becerril DU, Sar EA (2008) Marine diatoms from Buenos Aires coastal waters (Argentina). V. species of the genus *Chaetoceros*. *Rev Biol Mar Oceanogr* 43(2):303–326
- Tachihana S, Nagao N, Katayama T, Hirahara M, Yusoff FM, Banerjee S, Shariff M, Kurosawa N, Toda T, Furuya K (2020) High Productivity of Eicosapentaenoic acid and fucoxanthin by a marine diatom *Chaetoceros gracilis* in a Semi-continuous culture. *Front Bioeng Biotechnol* 8:602721
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599
- Taş S, Hernández-Becerril DU (2017) Diversity and distribution of the planktonic diatom genus *Chaetoceros* (Bacillariophyceae) in the Golden Horn Estuary (Sea of Marmara). *Diatom Res* 32:309–323
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22:4673–4680
- Toyoda K, Nakasaki K, Williams DM, Tomaru Y (2011) PCR-RFLP Analysis for Species-Level Distinction of The Genus *Chaetoceros* Ehrenberg (Bacillariophyceae). *Nat Sci* 50:21–29
- Xu, X., Ji, W., Castell, J.D. and O'Dor, R., (1993). The nutritional value of dietary n-3 and n-6 fatty acids for the Chinese prawn (*Penaeus chinensis*). *Aquaculture*. 118 (3e4): 277–285.
- Zailanie K, Purnomo H (2017) Identification of fucoxanthin from brown algae (*Sargassum filipendula*) from Padike village, Talango district, Sumenep regency, Madura islands, using nuclear magnetic resonance (NMR). *Int Food Res J* 24(1):372–378
- Zhou QC, Li CC, Liu CW, Chi SY, Yang QH (2007) Effects of dietary lipid sources on growth and fatty acid composition of juvenile shrimp, *Litopenaeus vannamei*. *Aquac Nutr* 13:222–229

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.