



Correction to: Polymyxin B causes DNA damage in HK-2 cells and mice

B. Yun¹ · T. Zhang² · M. A. K. Azad¹ · J. Wang¹ · C. J. Nowell¹ · P. Kalitsis² · T. Velkov^{1,3} · D. F. Hudson² · J. Li^{1,4} 

Published online: 18 June 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: Archives of Toxicology

<https://doi.org/10.1007/s00204-018-2192-1>

In the original publication of the article, part of Fig. 6 is missing. The missing subpanels, Fig. 6c, d are given below.

The original article can be found online at <https://doi.org/10.1007/s00204-018-2192-1>.

✉ D. F. Hudson
damien.hudson@mcri.edu.au

✉ J. Li
jian.li@monash.edu

¹ Monash Institute of Pharmaceutical Sciences, Monash University, Melbourne, Australia

² Murdoch Children's Research Institute, Royal Children's Hospital, Melbourne, Australia

³ Department of Pharmacology and Therapeutics, University of Melbourne, Melbourne, Australia

⁴ Infection & Immunity Program, Department of Microbiology, Biomedicine Discovery Institute, Monash University, Melbourne, Australia

Fig. 6 c Time-course of PMB-induced nephrotoxicity in mice showed mean increase in foci to cell percentile of total population as time post-treatment increased, with representative examples of undamaged and damaged mouse kidney sections showing γ H2AX foci staining. All data are presented as mean \pm SD. **d** Foci percentage of individual mice which showed no obvious histological damage ($n = 8$)

