



On two mod p period maps: Ekedahl–Oort and fine Deligne–Lusztig stratifications

Fabrizio Andreatta¹

Received: 22 October 2020 / Revised: 3 November 2021 / Accepted: 21 December 2021 /
Published online: 13 January 2022

© The Author(s) 2022

Abstract

Consider a Shimura variety of Hodge type admitting a smooth integral model S at an odd prime $p \geq 5$. Consider its perfectoid cover $S^{\text{ad}}(p^\infty)$ and the Hodge–Tate period map introduced by Caraiani and Scholze. We compare the pull-back to $S^{\text{ad}}(p^\infty)$ of the Ekedahl–Oort stratification on the mod p special fiber of a toroidal compactification of S and the pull back to $S^{\text{ad}}(p^\infty)$ of the fine Deligne–Lusztig stratification on the mod p special fiber of the flag variety which is the target of the Hodge–Tate period map. An application to the non-emptiness of Ekedahl–Oort strata is provided.

Keywords Ekedahl–Oort stratifications · Deligne–Lusztig varieties · Period maps

Mathematics Subject Classification 14G35 · 11G18

Contents

1	Introduction	512
2	Algebraic zip data	515
2.1	Fine Deligne–Lusztig varieties	517
2.1.1	A variant	519
2.2	H -zips	520
2.3	The Ekedahl–Oort stratification	521
3	Smoothness of ζ^Σ	523
3.1	The structure at the boundary	524
3.2	The underlying algebraic groups	525
3.3	The proof	527
3.4	Ekedahl–Oort strata at the boundary	530

Communicated by Wei Zhang.

✉ Fabrizio Andreatta
Fabrizio.Andreatta@unimi.it

¹ Dipartimento di Matematica “F. Enriques”, Università degli Studi di Milano, via C. Saldini, 50, Milano 20133, Italy

3.5 A result on zips in mixed characteristics 530
 4 On the mod p Hodge–Tate period map 532
 5 Proof of Theorem 1.1 536
 5.1 The setup 537
 5.2 A description of $\zeta'_1(x_0)$ and $\zeta'_2(x_0)$ 538
 5.3 Reduction steps 539
 5.4 Conclusion 542
 6 Appendix I: Fontaine’s rings 543
 6.1 An approximation result 546
 7 Appendix II: some facts about Weyl groups 547
 References 549

1 Introduction

Let (G, X) be a Shimura datum; in particular, G is a connected reductive group over \mathbb{Q} and X is a $G(\mathbb{R})$ -conjugacy class of a homomorphism of algebraic groups $h : \text{Res}_{\mathbb{C}/\mathbb{R}}(\mathbb{G}_m) \rightarrow G_{\mathbb{R}}$. Denote by $\text{Sh}_K(G, X)$ the associated Shimura variety over \mathbb{C} , for some compact open subgroup $K \subset G(\mathbb{A}^f)$ of the finite adelic points of G . We fix an odd prime p and we assume that:

- (i) K is hyperspecial at p i.e., $K = K_p K^p$ with $K^p \subset G(\mathbb{A}^{f,(p)})$ a compact open subgroup of the prime-to- p , finite adelic points of G and $K_p = G_{\mathbb{Z}_p}(\mathbb{Z}_p)$, the group of \mathbb{Z}_p -points of some reductive group $G_{\mathbb{Z}_p}$ over \mathbb{Z}_p with generic fiber $G_{\mathbb{Q}_p}$;
- (ii) K is neat;
- (iii) (G, X) is of Hodge type, that is, there is an embedding of Shimura data $(G, X) \subset (\text{GSp}_{2g}, \mathbb{H}_g)$ with GSp_{2g} the group of symplectic similitudes on a \mathbb{Q} -vector space of dimension $2g$, endowed with a non-degenerate symplectic pairing and with \mathbb{H}_g the double Siegel space.

Results of Shimura and Deligne guarantee that $\text{Sh}_K(G, X)$ admits a canonical model over the reflex field E which is a number field, defined as the minimal subfield of \mathbb{C} of definition of the conjugacy class Z of the Hodge cocharacter

$$\mu_h : \mathbb{G}_{m,\mathbb{C}} \xrightarrow{\text{Id} \times 1} \mathbb{G}_{m,\mathbb{C}} \times \mathbb{G}_{m,\mathbb{C}} \cong \text{Res}_{\mathbb{C}/\mathbb{R}}(\mathbb{G}_m)_{\mathbb{C}} \xrightarrow{h_{\mathbb{C}}} G_{\mathbb{C}}.$$

Fix a prime v of E over the prime p ; let κ be the residue field of \mathcal{O}_E at v . Thanks to our assumptions v is unramified over p so that $\widehat{\mathcal{O}}_{E,v} \cong \mathbb{W}(\kappa)$ and Z admits a $\mathbb{W}(\kappa)$ -valued point that we denote μ .

Denote by S the integral canonical model of $\text{Sh}_K(G, X)$ over $\mathcal{O}_{E,v}$ constructed by Kisin [11]. Given a finite and admissible rational cone decomposition Σ for G , we denote by S^{Σ} the associated toroidal compactification constructed in [16]. We fix one and we suppose that S^{Σ} is smooth and projective over $\mathcal{O}_{E,v}$ (and certain other technical assumptions for Σ hold; see (2.12)). We let S_0^{Σ} be the mod p special fiber. Write S^{Σ} for the p -adic formal scheme defined by completing S^{Σ} along S_0^{Σ} and $S^{\Sigma,\text{ad}}$ for its adic generic fiber (in the sense of Huber). We denote by $S^{\text{ad}} \subset S^{\Sigma,\text{ad}}$ the open, adic subspace defined by the scheme $S_{\mathbb{Q}_p}$.


We also let $\text{Fl}_{G,\mu^{-1}}$ be the flag variety over $\mathbb{W}(\kappa)$ classifying parabolic subgroups of $G_{\mathbb{Z}_p}$ of type μ^{-1} . Set $\text{Fl}_{G,\mu^{-1},0}$ to be the special fiber over κ , $\mathcal{F}\ell_{G,\mu^{-1}}$ to be the

formal scheme defined by the completion of $\mathrm{Fl}_{G,\mu^{-1}}$ along $\mathrm{Fl}_{G,\mu^{-1},0}$ and $\mathcal{F}\ell_{G,\mu^{-1}}^{\mathrm{ad}}$ to be the generic adic fiber of $\mathcal{F}\ell_{G,\mu^{-1}}$.

Consider the perfectoid tower $\pi : S^{\mathrm{ad}}(p^\infty) \rightarrow S^{\mathrm{ad}}$ and the Hodge–Tate period map

$$\pi_{\mathrm{HT}} : S^{\mathrm{ad}}(p^\infty) \longrightarrow \mathcal{F}\ell_{G,\mu^{-1}},$$

defined by [5, Thm. 2.1.3]. We have the following diagram of topological spaces:



where the bottom vertical arrows are the specialization maps $\mathrm{sp} : S^{\Sigma,\mathrm{ad}} \rightarrow S_0^\Sigma$ and $\mathrm{sp} : \mathcal{F}\ell_{G,\mu^{-1}}^{\mathrm{ad}} \rightarrow \mathrm{Fl}_{G,\mu^{-1},0}$. We write $q_1 : S^{\mathrm{ad}}(p^\infty) \rightarrow S_0^\Sigma$ and $q_2 : S^{\mathrm{ad}}(p^\infty) \rightarrow \mathrm{Fl}_{G,\mu^{-1},0}$ for the two composite maps. These are the two mod p period maps of the title.

The varieties S_0^Σ and $\mathrm{Fl}_{G,\mu^{-1},0}$ are endowed with natural stratifications. On S_0^Σ we have the so called *Ekedahl–Oort stratification* introduced by Ekedahl and Oort in the Siegel case, by Zhang [26] in the general setting of the special fiber of Shimura varieties of Hodge type, such as S_0 , and then extended to toroidal compactifications, such as S_0^Σ , by Goldring and Koskivirta [7]. On the other hand $\mathrm{Fl}_{G,\mu^{-1},0}$ is endowed with the so called *fine Deligne–Lusztig stratification*, introduced by Lusztig in [14] and [15]. One knows that:

- the strata of these two stratifications are classified by subsets ${}^{I_+}W$ (in the EO case) and ${}^{I_-}W$ (in the DL case) of the Weyl group W of G . Here, I_+ and I_- are the types of the parabolic subgroups P_{μ^\pm} of G defined by μ and μ^{-1} respectively and ${}^{I_\pm}W$ is the set of Kostant representatives of the quotient sets W/W_{I_\pm} of W by the Weyl group of the standard Levi subgroups of P_{I_\pm} ;
- there exist explicit order relations \preceq_{EO} on ${}^{I_+}W$ and \preceq_{DL} on ${}^{I_-}W$ corresponding to the closure relations on the associated Ekedahl–Oort stratification on S_0^Σ and Deligne–Lusztig stratification on $\mathrm{Fl}_{G,\mu^{-1},0}$ respectively.
- the stratum associated to $w \in {}^{I_\pm}W$ is smooth and equidimensional of dimension equal to the length $\ell(w)$ of w in the Weyl group W , if non-empty.

For $w \in {}^{I_\pm}W$ we let $C_{\mathrm{EO}}(w)$, resp. $C_{\mathrm{DL}}(w)$ be the corresponding stratum on S_0^Σ , resp. on $\mathrm{Fl}_{G,\mu^{-1},0}$. The natural question we address in this paper is the relation between these two stratifications when pulled-back to $S^{\mathrm{ad}}(p^\infty)$. Our main result is the following. In Lemma 7.2 we prove that the map

$$\sigma_0 : {}^{I_+}W \rightarrow {}^{I_-}W, \quad w \mapsto w_0^{I_+}w,$$

given by left multiplication by a suitable element $w_0^{I^+} \in W$, is an order reversing bijection and $\ell(\sigma_0(w)) = \ell(w_0^{I^+}) - \ell(w)$ for every $w \in I^+W$. Here the order relations are \preceq_{EO} and \preceq_{DL} respectively. Then:

Theorem 1.1 *Assume that $p \geq 5$ and that $q_1^{-1}(C_{EO}(w')) \cap q_2^{-1}(C_{DL}(w))$ is non-empty for some $w' \in I^+W$ and $w \in I^-W$. Then $w \preceq_{DL} \sigma_0(w')$ (equivalently $w' \preceq_{EO} \sigma_0(w)$).*

This should be seen as the discrete analogue of the phenomenon pointed out in [5], where a stratification (in a loose sense) of $\mathcal{F}\ell_{G,\mu^{-1}}^{\text{ad}}$ is defined so that its inverse image via π_{HT} coincides with the inverse image of the Newton polygon stratification on S^{ad} but the closure relation is opposite to the one of the Newton stratification. In loc. cit., the classification of vector bundles on the Fargues–Fontaine curve $\mathcal{X}_{\mathbb{C}_p}$ plays a crucial role. Recall that in Fontaine’s theory we have a classical period ring A_{inf} and a map $A_{\text{inf}} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ with kernel a principal ideal, of which we fix a generator ω . The Fargues–Fontaine curve is the quotient of $\text{Spa}(A_{\text{inf}}[p^{-1}\omega^{-1}], A_{\text{inf}})$ with respect to Frobenius. In our case we need to understand what happens at the non-analytic point of $\text{Spa}(A_{\text{inf}}, A_{\text{inf}})$ defined by the vanishing of the ideal (p, ω) . The assumption that $p \geq 5$ comes from the technical requirement that a certain open of $\mathcal{X}_{\mathbb{C}_p}$ is non-empty.

The EO stratification is constructed by Goldring and Koskivirta using a map of stacks

$$\zeta^\Sigma : S_0^\Sigma \longrightarrow H_{\mathbb{F}_p} - \text{Zip}^\chi$$

to the stack of so called H -Zips, where H is the mod p reduction of a model of G over \mathbb{Z}_p and χ is the cocharacter of H defined by the mod p reduction of the Hodge cocharacter μ . The notion of H -Zips is due Pink et al. [21] and is recalled in Sect. 2.2. It fits in the language of Algebraic Zip Data introduced by Pink et al. [20] and reviewed in Sect. 2. This provides a common framework to prove at the same time the properties of the EO and the DL stratifications recalled above (see Sects. 2.1 and 2.3 respectively). We stress, though, that historically these properties were already known in the DL case. Namely, Bédard has proven in [1] that DL strata are indexed by the elements of I^-W . The partial order \preceq_{DL} on I^-W has been introduced by He [8] and it is shown in [9] that it describes the closure relation of the DL stratification.

A second result, conjectured by Wedhorn and Ziegler [23], is Theorem 3.1 stating that

Theorem 1.2 *Assume that $p \geq 3$. Then, the map ζ^Σ is a smooth morphism of stacks.*

Remark 1.3 The proof of 1.2 uses as an input that the restriction of the maps $\zeta^\Sigma|_{S_0}$ to S_0 is smooth for the special fiber of the integral canonical model of Shimura varieties of Hodge type with hyperspecial level at p . This is proved in [26, Cor. 3.12] under the assumption that $p \geq 3$ as loc. cit. uses the integral canonical models defined in [11] only for p odd. Notice that also ζ^Σ is constructed in [7] for $p \geq 3$.

Nowadays, integral canonical models for Shimura varieties of abelian type with hyperspecial level at primes above 2 are available thanks to work of Kim and Madapusi Pera, see [10]. Their toroidal compactifications have been constructed in [16]. As

remarked by one of the referees, assuming that the construction of ζ^Σ of [7] goes through also in this case and admits the local description given in Lemma 2.13 and assuming that [26, Cor. 3.12] holds true also in this 2-adic setting, the proof of Theorem 1.2 works for $p = 2$ as well.

Theorems 1.1 and 1.2 allow to transfer questions on one stratification to questions on the other stratification. For example, we prove the following

Corollary 1.4 *Assume that $p \geq 5$. Then, for every $w' \in {}^L W$ and every $w \in {}^L W$, the sets $q_1^{-1}(C_{EO}(w'))$ and $q_2^{-1}(C_{DL}(w))$ are non-empty. In fact, $q_1^{-1}(C_{EO}(w') \cap S_0)$ is also non-empty.*

The result in the EO case has been announced by Yu [25] based on work of Lee [13] and Kisin [12]. We prove this result differently by showing first in Proposition 4.3 the stronger statement that the map q_2 is surjective, and hence that every DL stratum is non-empty. This is elementary projective geometry. It implies, in particular, that $q_2^{-1}(C_{DL}({}^L w_0))$ is non-empty, where the element $w_0^{L+} = {}^L w_0 \in {}^L W$ is the element of maximal length. Due to Theorem 1.1 this implies that the zero dimensional stratum $C_{EO}(\mathbf{1})$, corresponding to the identity element $\mathbf{1}$, is non-empty. We deduce from Theorem 1.2 that the EO stratification is closed under generalization for the order \preceq_{EO} and, hence, all the EO strata are non-empty as $C_{EO}(\mathbf{1})$ is non-empty. As the map q_1 is the composite of the surjective maps $\pi : S^{\text{ad}}(p^\infty) \rightarrow S^{\text{ad}}$ and $\text{sp} : S^{\text{ad}} \rightarrow S_0^\Sigma$, it is also surjective and the first part of the Corollary follows.

In fact, we show in Corollary 3.7 that the stratum $C_{EO}(\mathbf{1})$ is fully contained in S_0 verifying Condition 6.4.2 in [7]. This implies the second claim of the Corollary.

2 Algebraic zip data

We recall the notion of Algebraic Zip Data introduced in [20]. We follow the notation and conventions of [7, Sect. 1.2]. Let H be a connected reductive group defined over the prime field \mathbb{F}_p . Fix an algebraic closure k of \mathbb{F}_p and a cocharacter $\chi : \mathbb{G}_{m,k} \rightarrow H_k$ over k . The pair (H, χ) is called a cocharacter pair in loc. cit. We denote by $P = P_\chi$ the parabolic subgroup of H_k defined by χ : it is characterized by the fact that $\text{Lie} P \subset \text{Lie} H_k$ is the sum of the non-negative weight spaces for the adjoint action of $\mathbb{G}_{m,k}$ via χ .

As H is defined over a finite field, there exists a Borel subgroup $B \subset H$, with maximal torus T , defined over \mathbb{F}_p . We assume that $B_k \subset P$; this is Assumption (1.2.3) of [7]. Possibly after conjugating χ by an element of $H(k)$, this holds. We let $W = W(H_k, T_k)$ be the Weyl group of H_k with respect to T_k and we let $\Delta \subset W$ be the subset of simple reflections defined by B_k . Let $I \subset \Delta$ be the type of P . It is the subset of simple reflections generating the Weyl group W_L of the standard Levi subgroup $L \subset P$, the Levi subgroup containing T_k . We then have two cosets representatives ${}^L W$ and W^I in W of the quotient set $W_I \backslash W$, resp. W/W_I : for every coset $W_I w \in W_I \backslash W$ the corresponding element $w' \in {}^L W$ is the unique element $w' \in W_I w$ of shortest length (and similarly for W^I and W/W_I).

With the notation and assumptions above, we recall the following definition [20, Def. 3.1].

Definition 2.1 An algebraic zip datum for the cocharacter pair (H, χ) is a triple $\mathcal{Z} := (P, Q, \varphi)$ where P is the parabolic subgroup of H_k defined by χ , Q is another parabolic subgroup of H_k and $\varphi: P/R_u P \rightarrow Q/R_u Q$ is an isogeny of the reductive quotients. The group

$$E_{\mathcal{Z}} := \{(p, q) \in P \times Q \mid \varphi(\pi_P(p)) = \pi_Q(q)\}$$

is called the zip group associated to \mathcal{Z} . It acts on H_k on the left as follows

$$E_{\mathcal{Z}} \times H_k \rightarrow H_k, \quad (p, q) \cdot g = pqg^{-1}$$

Thanks to [20, Prop. 3.7] there exists $g \in H(k)$ such that

- (a) ${}^s B_k \subset Q$,
- (b) $\varphi(\pi_P(B_k)) = \pi_Q({}^s B_k)$,
- (c) $\varphi(\pi_P(T_k)) = \pi_Q({}^s T_k)$.

Let $L \subset P$ and $M \subset Q$ be the standard Levi subgroups defined by $T_k \subset P$ and ${}^s T_k \subset Q$ respectively. The isogeny φ defines an isogeny $\varphi: L \rightarrow M$ such that $\varphi(B_k \cap L) = {}^s B_k \cap M$ and $\varphi(T_k) = {}^s T_k$. If J is the type of Q this induces an isomorphism

$$\psi: (W_I, I) \xrightarrow{\cong} (W_J, J) \tag{1}$$

of the Weyl groups $W_I \rightarrow W_J$ such that $\Psi(I) = J$.

We also fix representatives $\tilde{w} \in N_{H_k}(T_k)$ for $w \in W = N_{H_k}(T_k)/C_{H_k}(T_k)$ such that $\widetilde{w_1 w_2} = \tilde{w}_1 \tilde{w}_2$ if $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$. This can be done as explained in [20, Sect. 2.3].

We assume that \mathcal{Z} is *orbitally finite*, that is the number of orbits for the action of $E_{\mathcal{Z}}$ on H_k is finite. This holds, for example, if the differential of the isogeny φ at 1 vanishes. For every $w \in W$ we write $H_w := E_{\mathcal{Z}} \cdot (\tilde{w}g^{-1}) \subset H_k$ for the $E_{\mathcal{Z}}$ -orbit defined by $\tilde{w}g^{-1}$. Then, [20, Thm. 7.5] states that:

Theorem 2.2 *The map*

$${}^I W \longrightarrow \{E_{\mathcal{Z}}\text{-orbits in } H_k\}, \quad w \mapsto H_w$$

is a bijection. Moreover,

1. for any $w \in {}^I W$ the corresponding orbit H_w is a locally closed, smooth subvariety of H_k of dimension $\dim P + \ell(w)$.
2. for any $w, w' \in {}^I W$ we have that $H_{w'} \subset \overline{H_w}$, and we write $w' \preceq w$, if and only if there exists $y \in W_I$ such that $yw'\psi(y)^{-1} \leq w$ for the Bruhat order \leq on W .

Remark 2.3 The Assumption that $B \subset P$ is related to the concept of frame in [20, Def. 3.6] for \mathcal{Z} . If we set (B', T', g') with $B' := g^{-1} B_k$, $T' := g^{-1} T_k$ and $g' = g$, the triple (B', T', g') is a frame for \mathcal{Z} . We prefer to use the conventions of [7] fixing a Borel $B_k \subset P$ instead of a Borel in Q . This has the effect, for example, that the statement of [20, Thm. 7.5] is slightly changed.

There is a dual presentation of the orbit space for the action of $E_{\mathcal{Z}}$ on H_k , described in [20, Thm. 11.2], in terms of the subset $W^J \subset W$. As explained in [20, Prop. 9.13, 9.14] there exists a length preserving bijection

$$\sigma : {}^I W \rightarrow W^J$$

characterized by the property that for any $w \in {}^I W$ there exists a $y \in W_I$ such that $\sigma(w) = yw\psi(y)^{-1}$. Then [20, Thm. 11.2, 11.5] state that:

Theorem 2.4 *The map $W^J \rightarrow \{E_{\mathcal{Z}} - \text{orbits in } H_k\}$, given by $w \mapsto H_w := E_{\mathcal{Z}} \cdot (\tilde{w}g^{-1})$, is a bijection. Moreover,*

1. *for any $w \in {}^I W$ we have $H_w = H_{\sigma(w)}$. In particular, for any $u \in W^J$ the corresponding orbit H_u is a locally closed, smooth subvariety of H_k of dimension $\dim P + \ell(u)$.*
2. *for any $w, w' \in W^J$ we have that $H_{w'} \subset \overline{H}_w$, and we write $w' \preceq w$, if and only if there exists $y \in W_I$ such that $yw'\psi(y)^{-1} \leq w$ for the Bruhat order \leq on W .*

In fact, we have a richer structure of quotient stacks; see for example [21, Sect. 2.2]. Recall that if X is a k -scheme and Q is a linear algebraic k -group scheme acting on the left on X the quotient stack $[Q \backslash X]$ is the stack fibered over the category of k -schemes whose S -valued points consist of pairs (I, α) with (1) I a left Q -torsor over S (2) $\alpha : I \rightarrow X$ a morphism commuting with the left Q -actions on I and on X . There is a notion of topological space associated to $[Q \backslash X]$ that we denote $|[Q \backslash X]|$

If the number of orbits $Q \backslash X$ in X for the action of H is finite, then such space has a unique topology compatible with the closure relation among orbits; see [21, Prop. 2.1]. Furthermore, [21, Prop. 2.2] guarantees that the natural map $X \rightarrow [Q \backslash X]$ induces an homeomorphism $(Q(k) \backslash X(k)) \cong |[Q \backslash X]|$. So Theorems 2.2 and 2.4 describe the topological space underlying the quotient stack $[E_{\mathcal{Z}} \backslash H_k]$.

2.1 Fine Deligne–Lusztig varieties

We use the notation of the previous section. We start with a connected reductive group H over \mathbb{F}_p and a cocharacter $\chi : \mathbb{G}_{m,k} \rightarrow H_k$, over $k = \overline{\mathbb{F}}_p$. This defines a parabolic subgroup $P \subset H_k$ that we assume to contain a Borel subgroup B_k , defined over \mathbb{F}_p . Write $\text{Fl}_{H_k, \chi} := P \backslash H_k$ for the corresponding flag variety. It is endowed with a stratification by *fine Deligne–Lusztig varieties*, introduced by Lusztig [14] and [15]. In this section we review its definition and we recast its properties, known thanks to results of Bédard [1] and He [8], [9], using the language of Algebraic Zip Data.

Let $Q := P^{(p)}$ be the base change of $P \subset H_k$ via Frobenius on k . It is a parabolic subgroup of $(H_k)^{(p)} = H_k$ (as H is defined over \mathbb{F}_p). It contains the Borel subgroup $(B_k)^{(p)} = B_k$ (as B_k is also defined over \mathbb{F}_p). The Frobenius map $\varphi : P \rightarrow P^{(p)} = Q$ defines an isogeny $\varphi : P/R_u P \rightarrow Q/R_u Q$.

Then $\mathcal{Z} := \mathcal{Z}_{\text{DL}} := (P, Q, \varphi)$ is a zip datum as in Definition 2.1. Note that $Q/R_u Q \cong (P/R_u P)^{(p)}$ and then φ is Frobenius. In particular, the differential of φ at 1 is zero and \mathcal{Z}_{DL} is orbitally finite.

The projection map $\rho: H_k \rightarrow \text{Fl}_{H_k, \chi}$ is a P -torsor (for the étale topology) so that $\text{Fl}_{H_k, \chi}$ represents the quotient stack $[P \backslash H_k]$. The inverse of Lang isogeny $\gamma: H_k \rightarrow H_k, h \mapsto h\varphi(h)^{-1}$ defines a map of quotient stack

$$\bar{\gamma}: [P \backslash H_k] \longrightarrow [E_{\mathcal{Z}} \backslash H_k].$$

Proposition 2.5 *The map $\bar{\gamma}$ is smooth. In particular, the induced map on underlying topological spaces is surjective, continuous and open.*

Proof Recall that to give a morphism $S \rightarrow [E_{\mathcal{Z}} \backslash H_k]$ is equivalent to give an $E_{\mathcal{Z}}$ -torsor Y over S and a $E_{\mathcal{Z}}$ -equivariant morphism $Y \rightarrow H_k$. In our case, using that $\text{Fl}_{H_k, \chi}$ represents $[P \backslash H_k]$, the morphism $\bar{\gamma}$ is defined by

- (1) the $E_{\mathcal{Z}}$ -torsor $E_{\mathcal{Z}} \times^P H_k$ over $\text{Fl}_{H_k, \chi}$ defined by pushing-forward $P \backslash (E_{\mathcal{Z}} \times_k H_k)$ of the inclusion $P \subset H_k$ via the group homomorphism $P \hookrightarrow E_{\mathcal{Z}}, h \mapsto (h, \varphi(h))$;
- (2) the $E_{\mathcal{Z}}$ -equivariant map $\nu: E_{\mathcal{Z}} \times^P H_k \rightarrow H_k$ defined by $((p, q), h) \mapsto (p, q) \cdot \gamma(h) = p\gamma(h)q^{-1}$.

Thus the diagram

$$\begin{array}{ccc} E_{\mathcal{Z}} \times^P H_k & \xrightarrow{\nu} & H_k \\ \downarrow & & \downarrow \\ \text{Fl}_{H_k, \chi} & \xrightarrow{\bar{\gamma}} & [E_{\mathcal{Z}} \backslash H_k], \end{array}$$

where the vertical maps are the natural projection maps, is cartesian and in order to prove the claim it suffices to show that ν is smooth. The scheme $E_{\mathcal{Z}} \times^P H_k$ is a smooth over k , being a torsor under the smooth algebraic group $E_{\mathcal{Z}}$ over the smooth scheme $\text{Fl}_{H_k, \chi}$. As also H_k is smooth, in order to prove that ν is smooth it suffices to show that it is surjective on the tangent space at every k -valued point.

The map γ is finite and étale so that inclusion $H_k \subset E_{\mathcal{Z}} \times^P H_k \rightarrow H_k$ is already surjective on tangent spaces and the claim follows. □

Recall from Theorems 2.2 and 2.4 that $[[E_{\mathcal{Z}} \backslash H_k]]$ is homeomorphic to ${}^I W$ (resp. W^J), where W is the Weyl group of H_k , I is the type of the parabolic P , J is the type of Q , we can take $g = \text{Id}$ in loc. cit. and the topological structure on ${}^I W$ (resp. W^J) is defined by the order relation $w' \preceq w$ described as follows.

Let I be the type of P and let $J = \varphi(I)$ be the type of $Q = P^{(p)}$. Frobenius on H_k defines an isomorphism of Weyl groups $\varphi: W \rightarrow W$ and a bijection on simple reflections, as the Weyl group and the simple reflections are defined with respect to a Borel subgroup and a torus defined over \mathbb{F}_p . Then φ defines an isomorphism

$$\psi_{\text{DL}}: W_I \longrightarrow W_J, \quad w \mapsto \varphi(w) \tag{2}$$

and, thanks to Theorems 2.2 and 2.4, for w and $w' \in {}^I W$ (or in W^J) we have $w' \preceq w$ iff there exists $y \in W_I$ such that $yw'\psi_{\text{DL}}(y)^{-1} \leq w$.

Definition 2.6 For every $w \in {}^I W$ we let $\text{Fl}_{H_k, \chi, w}$ be the inverse image of $w \in {}^I W$ via the map of topological space $\text{Fl}_{H_k, \chi} \rightarrow |[E_{\mathcal{Z}} \setminus H_k]| \cong {}^I W$ with reduced induced scheme structure.

Similarly for $u \in W^J$ we let $\text{Fl}_{H_k, \chi}^u$ be the inverse image of $u \in W^J$ via the map of topological space $\text{Fl}_{H_k, \chi} \rightarrow |[E_{\mathcal{Z}} \setminus H_k]| \cong W^J$.

By Theorem 2.4 there is a bijection $\sigma : {}^I W \cong W^J$ and for every $w \in {}^I W$ we have $\text{Fl}_{H_k, \chi, w} = \text{Fl}_{H_k, \chi}^{\sigma(w)}$.

Corollary 2.7 Each $\text{Fl}_{H_k, \chi, w}$ and each $\text{Fl}_{H_k, \chi}^u$ is smooth and equidimensional of dimension $\ell(w)$, resp. $\ell(u)$. Moreover $\text{Fl}_{H_k, \chi, w'}$ is in the closure of $\text{Fl}_{H_k, \chi, w}$ (respectively $\text{Fl}_{H_k, \chi}^{u'}$ is in the closure of $\text{Fl}_{H_k, \chi}^u$) if and only if $w' \preceq w$ (respectively $u' \preceq u$).

Proof We provide a proof for the strata $\text{Fl}_{H_k, \chi, w}$. For the strata $\text{Fl}_{H_k, \chi}^u$ the argument is the same. The closure relation and the fact that each $\text{Fl}_{H_k, \chi, w}$ is non-empty follows from the fact that the map of topological spaces induced by $\bar{\gamma}$ is continuous, open and surjective. This also implies that such map respects codimensions. Since by Theorem 2.2 the codimension of $w \in {}^I W$ is $\dim H_k - (\dim P + \ell(w))$ it follows that the codimension of $\text{Fl}_{H_k, \chi, w}$ in $\text{Fl}_{H_k, \chi}$ is also $\dim H_k - (\dim P + \ell(w))$. Hence its dimension is $\ell(w)$ as wanted. The smoothness assertion follows from the smoothness of $\bar{\gamma}$. \square

In particular, we get a stratification on $\text{Fl}_{H_k, \chi}$ with strata $\text{Fl}_{H_k, \chi, w}$, for $w \in {}^I W$. Thanks to results of He [9] it coincides with the stratification by fine Deligne–Lusztig varieties introduced by Lusztig.

2.1.1 A variant

We consider the flag variety $\text{Fl}_{H_k, \chi^{-1}} := P_- \setminus H_k$. We describe an isomorphism

$$\text{Fl}_{H_k, \chi^{-1}} \cong \text{Fl}_{H_k, w_0 \chi^{-1}},$$

where $w_0 \in W$ the longest element in the Weyl group W , compatible with stratifications. Later in the paper we will be lead to work with the first space but we will need the dictionary provided by this isomorphism to get a description of the stratification in terms of the Weyl group W and the fixed set of simple reflections Δ .

Recall that I is the type of P and I_- is the type of the parabolic P_- defined by the cocharacter χ^{-1} . As P_- is the parabolic opposite to P we have $P_{I_-} = {}^{w_0} P_-$ so that $I_- = {}^{w_0} I$. Recall also that Frobenius defines an automorphism φ of the Weyl group W and that $J := \varphi(I)$ is the type of $P^{(p)}$ and $K = \varphi(I_-) = {}^{w_0} J$ is the type of $Q = P_-^{(p)}$. Consider the elements $y := {}^I w_0 = w_0^{-I}$ and $z = \varphi(y) = {}^J w_0 = w_0^K$ of W defined in Lemma 7.1. Then $P_- = {}^y P_{I_-}$ and conjugation by \tilde{y}^{-1} provides the identification $L \cong {}^{w_0} L$ of the Levi subgroup $L = P \cap P_-$ of P_- and the Levi subgroup of P_{I_-} so that $B \cap L \cong B \cap {}^{w_0} L$. Similarly, conjugation by \tilde{z}^{-1} provides the identification $L^{(p)} \cong {}^{w_0} L^{(p)}$ of the Levi subgroup $Q = P_-^{(p)}$ and the Levi subgroup of $P_K = P_{I_-}^{(p)}$ so that $B \cap L^{(p)} \cong B \cap {}^{w_0} L^{(p)}$. Consider the zip data $\mathcal{Z}_{\text{opp}} := (P_-, Q, \varphi)$ and $\mathcal{Z}_- := (P_{I_-}, P_K, \varphi)$.

Lemma 2.8 *The map $E_{Z_{opp}} \rightarrow E_{Z_-}, (p, q) \mapsto (\tilde{y}^{-1}p\tilde{y}, \tilde{z}^{-1}q\tilde{z})$, defines an isomorphism of algebraic groups.*

Similarly, the map $H_\kappa \rightarrow H_\kappa, h \mapsto \tilde{y}^{-1}h\tilde{z}$, defines an isomorphism which is equivariant for the action of $E_{Z_{opp}}$ and E_{Z_-} respectively and an isomorphism $\text{Fl}_{H_\kappa, \chi^{-1}} \rightarrow \text{Fl}_{H_\kappa, w_0\chi^{-1}}$ making the following diagram (see Proposition 2.5)

$$\begin{CD} \text{Fl}_{H_\kappa, \mu^{-1}} @>>> [E_{Z_{opp}} \setminus H_\kappa] \\ @VVV @VVV \\ \text{Fl}_{H_\kappa, w_0\mu^{-1}} @>>> [E_{Z_-} \setminus H_\kappa] \end{CD}$$

commute. In particular, the topological space underlying $[E_{Z_-} \setminus H_\kappa]$ admits a description in terms of ${}^L\text{-}W$ with the order \preccurlyeq as in Theorem 2.2 and the scheme $\text{Fl}_{H_\kappa, \mu^{-1}}$ admits an induced stratification labeled by ${}^L\text{-}W$.

2.2 H-zips

Recall the notation. We have a connected reductive group H over \mathbb{F}_p and a cocharacter $\chi : \mathbb{G}_{m,k} \rightarrow H_k$, over $k = \overline{\mathbb{F}}_p$, that defines a parabolic subgroup $P = P_+ \subset H_k$. We assume it contains a Borel subgroup B_k , defined over \mathbb{F}_p .

Denote by P_- be the associated opposite parabolic subgroup of H_k defined by the cocharater χ^{-1} . It is characterized by the property that $\text{Lie}P_- \subset \text{Lie}H_k$ is the sum of the non-positive weight spaces for the adjoint action of $\mathbb{G}_{m,k}$ via χ . Let U and V be the unipotent radicals of P and P_- respectively. The quotients P/U and P_-/V are isomorphic reductive groups and we denote such algebraic group by L . Set $Q := P_-^{(p)} \subset H_k^{(p)} = H_k$. It is a parabolic subgroup of H_k with unipotent radical $V^{(p)}$. The Frobenius map $\varphi : P \rightarrow P^{(p)}$ defines an isogeny $\varphi : P/U \cong L \rightarrow L^{(p)} \cong Q/V^{(p)}$.

Also in this case $\mathcal{Z} := \mathcal{Z}_{EO} := (P, Q, \varphi)$ is a zip datum as in Definition 2.1. It is orbitally finite as φ has trivial differential at 1. We consider the quotient stack $[E_{\mathcal{Z}_{EO}} \setminus H_k]$. Following [21, Def. 1.4] we give the following:

Definition 2.9 We define the category $H\text{-Zip}^\chi$ fibred over the category Sch_k of schemes over k as follows.

For a scheme S over k the objects are quadruples (I, I_+, I_-, ι) where I is a right H_k -torsor, $I_+ \subset I$ is a right P -torsor, $I_- \subset I$ is a right Q -torsor and $\iota : I_+^{(p)}/U^{(p)} \rightarrow I_-/V^{(p)}$ is an isomorphism of $L^{(p)}$ -torsors.

A morphism of two objects $(I, I_+, I_-, \iota) \rightarrow (I', I'_+, I'_-, \iota')$ is a triple of equivariant morphisms $I \rightarrow I', I_\pm \rightarrow I'_\pm$, which are compatible with the inclusions $I_\pm \subset I$ and $I'_\pm \subset I'$ and with ι and ι' .

It is easily verified that $H\text{-Zip}^\chi$ is a stack over Sch_k . There is a morphism $H_k \rightarrow H\text{-Zip}^\chi$, given by [21, Construction 3.4]. To a k -scheme S and an element $g \in H_k(S)$ we associate $I_g = S \times_k H_k, I_{g,+} := S \times_k P, I_{g,-} := g \cdot (S \times_k Q) \subset S \times_k H_k,$

$$\iota_g : S \times_k P/U = S \times_k L \xrightarrow{1 \times \varphi} S \times_k L^{(p)} = (S \times_k Q/V^{(p)}) \xrightarrow{g \cdot} I_{-,g}/V^{(p)}.$$

Then $I_g := (I_g, I_{g,+}, I_{g,-}, \iota_g) \in H\text{-Zip}^\chi(S)$. By [21, Prop. 3.11]:

Proposition 2.10 *The map $H_k \rightarrow H - \text{Zip}^\lambda$, $g \mapsto I_g$, induces an isomorphism of stacks $[E_{Z_{EO}} \setminus H_k] \cong H - \text{Zip}^\lambda$.*

Let I be the type of P , let $J = \varphi(I)$ be the type of $P^{(p)}$ and let $K = {}^{w_0}J$ be the type of Q which is the parabolic opposite to $P^{(p)}$. Here we use that Frobenius on H_k defines an isomorphism of Weyl groups $\varphi: W \rightarrow W$ and a bijection on simple reflections, as the Weyl group and the simple reflections are defined with respect to a Borel and a torus defined over \mathbb{F}_p . In particular, by Theorems 2.2 and 2.4, we have

Corollary 2.11 *The map ${}^I W \rightarrow H - \text{Zip}^\lambda$, $w \mapsto [\tilde{w}z^{-1}]$ with $z = w_0^K$, defines a homeomorphism of underlying topological spaces, and equivalently for W^K instead of ${}^I W$, with the topological structure defined by the partial order \preceq .*

We can describe the order relation as follows. In the notation of Theorem 2.2, we have $g = \tilde{z}$ with $z = w_0^K$ and ψ is the map

$$\psi_{EO}: (W_I, I) \cong (W_K, K), \quad w \mapsto z\varphi(w)z^{-1}. \tag{3}$$

Thanks to Theorems 2.2 and 2.4, for w and $w' \in {}^I W$ or w and $w' \in W^K$, we have $w' \preceq w$ iff there exists $y \in W_I$ such that $yw'\psi_{EO}(y)^{-1} \leq w$.

2.3 The Ekedahl–Oort stratification

Recall from the introduction that we have a scheme $S = S_K$, the integral canonical model of the Shimura variety $\text{Sh}_K(G, X)$ over $\mathcal{O}_{E,v}$, and its sepcial fiber S_0 over the residue field κ of $\mathcal{O}_{E,v}$.

By [11, Sect. 2.3.1] we may assume that the inclusion $G \subset \text{GSp}_{2g}$ extends to a morphism of algebraic groups $G_{\mathbb{Z}_p} \subset \text{GSp}_{2g, \mathbb{Z}_p}$ where $G_{\mathbb{Z}_p}$ is the given model of G over \mathbb{Z}_p and $\text{GSp}_{2g, \mathbb{Z}_p}$ is the base change from $\mathbb{Z}_{(p)} \rightarrow \mathbb{Z}_p$ of the group of symplectic similitudes of free $\mathbb{Z}_{(p)}$ -module Λ of rank $2g$ endowed with a symplectic perfect pairing ψ . The subgroup $G_{\mathbb{Z}_p}$ can be realized as the subgroup fixing a finite set of elements $(s_\alpha)_\alpha$ in the tensor algebra of Λ .

Furthermore, possibly after refining the prime-to- p level subgroup $K^p \subset G(\hat{A}^{f,(p)})$ we may assume that there exists an hyperspecial level $K_g := K_{g,p}K_g^p \subset \text{GSp}_{2g, \mathbb{Z}_p}(\hat{A}^f)$ inducing a morphism of Shimura varieties $\text{Sh}_K(G, X) \rightarrow \text{Sh}_{K_g}(\text{GSp}_{2g}, \mathbb{H}_g)$. This provides a finite morphism of the integral models

$$\xi: S_K \longrightarrow S_{K_g} \otimes_{\mathbb{Z}_{(p)}} \mathcal{O}_{E,v},$$

where S_{K_g} is a Siegel modular variety. In particular, we may assume that it is a fine moduli space so that it carries a universal abelian scheme.

We set $H := (G_{\mathbb{Z}_p})_{\mathbb{F}_p}$. As explained in [7, Sect. 4.1.7] there exists a Borel subgroup $B_{\mathbb{Z}_p} \subset G_{\mathbb{Z}_p}$. The Hodge cocharacter admits a representative $\tilde{\mu}$ in its conjugacy class defined over $\hat{\mathcal{O}}_{E,v} (\cong \mathbb{W}(\kappa))$. Let $\chi: \mathbb{G}_{m,\kappa} \rightarrow H_\kappa$ be the reduction of $\tilde{\mu}$ modulo p . The composite of $\tilde{\mu}$ with the inclusion $G^{\mathbb{W}(\kappa)} \subset \text{GSp}_{2g, \mathbb{W}(\kappa)}$ defines a splitting

$\Lambda \otimes \mathbb{W}(\kappa) = \Lambda_0 \oplus \Lambda_{-1}$ where \mathbb{G}_m acts on Λ_i via the character $z \mapsto z^{-i}$ on Λ_{-i} for $i = 0, -1$.

The main result of [26, Thm. 3.1.2] provides a smooth map

$$\zeta : S_0 \longrightarrow H - \text{Zip}^\chi.$$

This is refined in [7, Thm. 6.2.1], using the work [16] on toridal compactifications, to a morphism

$$\zeta^\Sigma : S_0^\Sigma \longrightarrow H - \text{Zip}^\chi,$$

where S_0^Σ is the special fiber of the smooth and projective toroidal compactification S^Σ of S associated to a finite and admissible rational cone decomposition Σ . We make the following Assumption, following [7], in order to define a well behaved compactification of the universal abelian scheme over $S_{K_g}^{\Sigma_g}$ and hence over S^Σ :

Assumption 2.12 Σ is supposed to refine the pull-back via ξ of a finite and admissible rational cone decomposition Σ_g for the Siegel case such that Σ_g is log-smooth (see [7, Sect. 6.1.2]).

In loc. cit. the weaker condition of Σ_g being log-integral is used. For the purposes of this paper log-smoothness is sufficient. We denote by $\xi^\Sigma : S^\Sigma \rightarrow S_{K_g}^{\Sigma_g}$ the induced morphism on toroidal compactifications.

For the rest of this section we recall the construction of the map ζ^Σ . Let \mathcal{S}^Σ be the p -adic formal scheme defined by completing S^Σ along its special fiber S_0^Σ . The first de Rham cohomology of the universal abelian scheme over S , obtained by pull-back from S_{K_g} , extends to a locally free $\mathcal{O}_{\mathcal{S}^\Sigma}$ -module H_{dR}^1 of rank $2g$ endowed with

- (i) the Hodge filtration $\Omega \subset H_{\text{dR}}^1$ which is a locally free sheaf of rank g , with locally free quotient;
- (ii) a connection ∇ with log poles along the complement of S in S^Σ , extending the Gauss-Manin connection. The base change \mathcal{H} of H_{dR}^1 to \mathcal{S}^Σ , together with the connection, is the evaluation along the DP-thickening $S_0^\Sigma \subset \mathcal{S}^\Sigma$ of the logarithmic crystal over S_0^Σ defines as the dual of the covariant log Dieudonné module of the mod p reduction of the degenerating universal abelian scheme over S_0^Σ (see [16, Prop. 1.3.5] and the discussion in [16, Sect. 4.3.1]);
- (iii) if $\text{Spf}(A) \subset \mathcal{S}^\Sigma$ is an affine open and we choose a lift $\varphi : A \rightarrow A$ of Frobenius on $A_0 = A/pA$, then the evaluation \mathcal{H}_A of the crystal \mathcal{H} at A is endowed with a φ -linear map, the Frobenius,

$$F : \mathcal{H}_A^{(p)} \longrightarrow \mathcal{H}_A.$$

- (iv) elements $(s_{\text{dR},\alpha})_\alpha$ in the tensor algebra of H_{dR}^1 such that

$$J := \text{Isom} \left(((\Lambda, (s_\alpha)_\alpha) \otimes \mathcal{O}_{\mathcal{S}^\Sigma}, H_{\text{dR}}^1, (s_{\text{dR}}, \alpha)_\alpha) \right)$$

is a right $G_{\mathbb{Z}_p}$ -torsor over S^Σ . If we further consider $g \in J$ such that g sends Ω isomorphically onto $\Lambda_0 \otimes \mathcal{O}_{S^\Sigma}$ we have a torsor J_+ under the parabolic P_μ^\pm defined by $\tilde{\mu}$.

Choose A and φ as in (iii). Then the restriction of F to Ω is divisible by p and it is proven in [7, Lemma 6.3.4] (see especially the first paragraph on p. 938) that $\frac{F}{p}(\Omega) + \text{Im}F = \mathcal{H}_A$. Take A small enough so that $J_+(A)$ is non-empty and choose $t \in J_+(A)$. We deduce that the A -linear map

$$g_t := t^{-1} \circ \left(\frac{F}{p}|_{t^{(p)}(\Lambda_0 \otimes A)} \oplus F|_{t^{(p)}(\Lambda_{-1} \otimes A)} \right) \circ t^{(p)} : \Lambda \otimes A \rightarrow \Lambda \otimes A$$

defines an element of $G(A)$. Let $g_{t,0} \in G(A_0)$ be its reduction modulo p .

Definition of the H -zip $\zeta^\Sigma(A_0)$. The element $g^{-1}t = (t^{-1}g)^{-1}$ defines an isomorphism $\mathcal{H}_A \cong \Lambda \otimes A$. Denote by J_- the $P_-^{(p)}$ -torsor of all its trivializations. Let I, I_+, I_- be the reduction modulo p of J, J_+ and J_- . They are right H_k , resp. P , resp. Q -torsors. The map $\gamma := t \circ g_t \circ (t^{(p)})^{-1} = \left(\frac{F}{p}|_{t^{(p)}(\Lambda_0 \otimes A)} \oplus F|_{t^{(p)}(\Lambda_{-1} \otimes A)} \right) : \mathcal{H}_A^{(p)} \rightarrow \mathcal{H}_A$ defines an isomorphism, as explained above. Let γ_0 be its reduction modulo p . Composing with γ_0 we get an isomorphism $\iota : I_+^{(p)}/U^{(p)} \rightarrow I_-/V^{(p)}$ of $L^{(p)}$ -torsors. Then, (I, I_+, I_-, ι) defines an H -zip in the sense of Definition 2.9 and

$$\zeta^\Sigma(A_0) := (I, I_+, I_-, \iota).$$

Then, in the notation of Proposition 2.10 (cf. [26, Rmk. 3.2.8] and [24, Prop. 6.7]),

Lemma 2.13 *We have $\zeta^\Sigma(A_0) = I_{g_{t,0}}$.*

Furthermore, if $z \in \tilde{P}(A)$, we consider the element $t \circ z \in J_+(A)$ and we write $z = \ell u$ with $\ell \in \tilde{L}(A)$ and $u \in R_\mu \tilde{P}(A)$ then

$$g_{t \circ z, 0} \equiv z_0^{-1} g_{t, 0} \ell_0^{(p)}.$$

Proof The first claim is a direct verification. The last formula follows as Frobenius is divisible by p on $t^{(p)}(\Lambda_0 \otimes A)$. □

3 Smoothness of ζ^Σ

As conjectured by Wedhorn and Ziegler in [23, Rmk. 6.4], we have:

Theorem 3.1 *The map ζ^Σ is smooth. In particular, it is open.*

The rest of this section is devoted to the proof of the Theorem and of some of its consequences. By construction S_0^Σ is a smooth scheme and $H - \text{Zip}^X$ is a smooth stack by [21, Cor. 3.12]. Hence, it suffices to prove that ζ^Σ is surjective on tangent spaces at every closed point $x_0 \in S_0^\Sigma$. If x_0 lies in S_0 this is the content of [26,

Thm. 4.1.2]. Assume that $x_0 \in S_0^\Sigma \setminus S_0$. Let A be the complete local ring of S^Σ at x_0 . Denote by \mathcal{H}_A the Frobenius logarithmic crystal relative to the DP-Thickening $A_0 = A/pA \subset A$ obtained by pull-back to A of the logarithmic crystal \mathcal{H} relative to the DP-thickening $S_0^\Sigma \subset S^\Sigma$ defined in Sect. 2.3.

3.1 The structure at the boundary

As explained in [16, Thm. 4.1.5] there exist

- (i) a smooth integral model S' over $\mathbb{W}(\kappa)$ associated to a Shimura datum (G', X') of Hodge type with hyperspecial level K' ;
- (ii) a torsor $\overline{S}' \rightarrow S'$ under an abelian scheme \mathcal{A} over S' ;
- (iii) a torsor $\overline{S} \rightarrow \overline{S}'$ under a torus \mathbf{E} over $\mathbb{W}(\kappa)$;
- (iv) a smooth torus embedding $\overline{S} \subset \widetilde{S}^\Sigma$ relative to \overline{S}'

such that the complete local ring A of S^Σ at x_0 is isomorphic to the completion of \widetilde{S}^Σ at a closed point s_0 .

Over S' one has a universal principally polarized abelian scheme B . Define the open $U \subset \text{Spec}(A)$ as the inverse image of $\overline{S} \subset \widetilde{S}^\Sigma$. By loc. cit. one has a degenerating 1-motive M_U . Write $M_U^\vee = [Y \rightarrow D_U]$ for the Cartier dual 1-motive; then D is a semiabelian scheme over \overline{S}' , extension of the principally polarized abelian scheme B over S' by a torus T which has constant rank. By the discussion in [16, Sect. 4.3.1] the Frobenius logarithmic crystal \mathcal{H}_A is identified with the contravariant log Dieudonné crystal of the mod p reduction of M_U .

In particular, as an A -module \mathcal{H}_A is endowed with a logarithmic connecton ∇ , with a two step Hodge filtration

$$0 = F^2\mathcal{H}_A \subset F^1\mathcal{H}_A \subset F^0\mathcal{H}_A = \mathcal{H}_A$$

and a three step weight filtration

$$0 \subset W_0\mathcal{H}_A \subset W_1\mathcal{H}_A \subset W_2\mathcal{H}_A = \mathcal{H}_A$$

consisting of Frobenius subcrystals such that

- (a) $W_1\mathcal{H}_A$ is identified with the covariant de Rham realization [16, Sect. 1.1.3] of D over A , i.e., the Lie algebra of the universal extension of the pull back of D to A and the Frobenius is obtained from its identification with the covariant crystalline realization of the base change of D to $A_0 = A/pA$;
- (b) $F^1\mathcal{H}_A$ is identified with the non-trivial step of the Hodge filtration of the covariant de Rham realization of D over A ;
- (c) $W_0\mathcal{H}_A$ is identified with the Lie algebra of the toric part T of D and the Frobenius structure is defined by the identification with the covariant Dieudonné module of the base change of T to A_0 .
- (d) the A module $\text{Gr}_1^W \mathcal{H}_A$ is identified with de Rham homology of B and its Frobenius structure arises from the identification with the covariant Dieudonné module of its base change B_0 to A_0 ;

- (e) $\text{Gr}_2^W \mathcal{H}_A$ is identified with the Lie algebra of the universal extension of Y and the Frobenius structure is defined by the identification with the covariant Dieudonné module of the étale p -divisible group $Y \otimes \mathbb{Q}_p$ over A_0 .

3.2 The underlying algebraic groups

We summarize the algebraic group counterpart of the description in Sect. 3.1; see the discussion in [16, Sect. 4.3.4]. The algebraic group $G' = N/U$, appearing in the Shimura datum (G', X') , is the quotient of a normal subgroup N of a parabolic subgroup R of G by the unipotent radical U of N and U is also the unipotent radical of R :

$$U \subset N \subset R \subset G.$$

All these groups are defined over \mathbb{Q} and they extend to subgroups

$$U_{\mathbb{Z}_p} \subset N_{\mathbb{Z}_p} \subset R_{\mathbb{Z}_p} \subset G_{\mathbb{Z}_p}$$

with $R_{\mathbb{Z}_p}$ a parabolic subgroup of the model $G_{\mathbb{Z}_p}$ of G over \mathbb{Z}_p , $N_{\mathbb{Z}_p}$ a normal subgroup of $R_{\mathbb{Z}_p}$ and $U_{\mathbb{Z}_p}$ the unipotent radical of $R_{\mathbb{Z}_p}$ (equivalently of $N_{\mathbb{Z}_p}$). The quotient $N_{\mathbb{Z}_p}/U_{\mathbb{Z}_p}$ is a reductive group $G'_{\mathbb{Z}_p}$. It is a normal subgroup of the reductive group $G''_{\mathbb{Z}_p} := R_{\mathbb{Z}_p}/U_{\mathbb{Z}_p}$ and defines a model of G' over \mathbb{Z}_p . In fact, the parabolic R is defined by a cocharacter, called the weight cocharacter, $w: \mathbb{G}_{m, \mathbb{Q}} \rightarrow G$ that factors via N . By [26, Prop. 3.2.1 and Sect. 3.2.2], after possibly conjugating w by N , we may assume that its base change to \mathbb{Q}_p extends to a cocharacter $w_{\mathbb{Z}_p}: \mathbb{G}_{m, \mathbb{Z}_p} \rightarrow N_{\mathbb{Z}_p}$. Using the embedding $G_{\mathbb{Z}_p} \subset \text{GSp}(\Lambda \otimes \mathbb{Z}_p)$ the cocharacter $w_{\mathbb{Z}_p}$ defines a grading on $\Lambda \otimes \mathbb{Z}_p$ and, hence, an ascending filtration $W_\bullet(\Lambda \otimes \mathbb{Z}_p)$, called the weight filtration:

$$\{0\} = W_{-1}(\Lambda \otimes \mathbb{Z}_p) \subset W_0(\Lambda \otimes \mathbb{Z}_p) \subset W_1(\Lambda \otimes \mathbb{Z}_p) \subset W_2(\Lambda \otimes \mathbb{Z}_p) = \Lambda \otimes \mathbb{Z}_p.$$

Thanks to [19, Prop. 12.1, Lemma 12.2] the Hodge cocharacter μ of G can be chosen to factor through N_E , where E is the reflex field of $\text{Sh}_K(G, X)$, and such that it acts trivially on U_E (via the adjoint action).

Lemma 3.2 *There exists a cocharacter $\tilde{\mu}$ of $G_{\mathbb{W}(\kappa)}$ in the conjugacy class of μ that factors through $N_{\mathbb{W}(\kappa)}$.*

Proof First consider the cocharacter μ' of the reductive group G'_E induced by μ . Due to [26, Prop. 3.2.1, S 3.2.2] there exists a conjugate cocharacter $\tilde{\mu}'$ that is defined for $G'_{\mathbb{W}(\kappa)}$. As $N_{\mathbb{W}(\kappa)} \rightarrow G'_{\mathbb{W}(\kappa)}$ has unipotent kernel the cocharacter $\tilde{\mu}'$ lifts uniquely to the sought for cocharacter $\tilde{\mu}$. \square

The cocharacter $\tilde{\mu}$ defines an increasing filtration, called the Hodge filtration, $F^\bullet(\Lambda \otimes \mathbb{W}(\kappa))$:

$$\{0\} = F^2(\Lambda \otimes \mathbb{W}(\kappa)) \subset F^1(\Lambda \otimes \mathbb{W}(\kappa)) \subset F^0(\Lambda \otimes \mathbb{W}(\kappa)) = \Lambda \otimes \mathbb{W}(\kappa)$$

such that $W_0(\Lambda \otimes \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} \mathbb{W}(\kappa) \subset F^1(\Lambda \otimes \mathbb{W}(\kappa)) \subset W_1(\Lambda \otimes \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} \mathbb{W}(\kappa)$.

We let $\tilde{P} \subset G_{\mathbb{W}(\kappa)}$, resp. $\tilde{P}' \subset G'_{\mathbb{W}(\kappa)}$, resp. $\tilde{P}'' \subset G''_{\mathbb{W}(\kappa)}$ be the parabolic subgroups defined by the cocharacter $\tilde{\mu}$ of $G_{\mathbb{W}(\kappa)}$ and the induced cocharacters $\tilde{\mu}'$ and $\tilde{\mu}''$ of $G'_{\mathbb{W}(\kappa)}$ and $G''_{\mathbb{W}(\kappa)}$ respectively. The intersection $\bar{P} := \tilde{P} \cap R_{\mathbb{W}(\kappa)}$ is a parabolic subgroup of $G_{\mathbb{W}(\kappa)}$; it is the parabolic subgroup containing $U_{\mathbb{W}(\kappa)}$ and with image \tilde{P}'' in $G''_{\mathbb{W}(\kappa)} = R_{\mathbb{W}(\kappa)}/U_{\mathbb{W}(\kappa)}$.

Here is the relation between the two descriptions of the boundary, one via degenerating 1-motives and the other via algebraic groups. Recall that the subgroup $G_{\mathbb{Z}_p}$ can be realized as the subgroup fixing a finite set of elements $(s_\alpha)_\alpha$ in the tensor algebra of Λ . By [16, Prop. 4.3.7] one has canonically defined tensors $(s_{\alpha, \text{dR}})_\alpha$ in the tensor algebra \mathcal{H}_A .

Proposition 3.3 *The set of isomorphisms $\mathcal{H}_A \cong \Lambda \otimes A$, sending the tensors $(s_{\alpha, \text{dR}})_\alpha$ to the tensors $(s_\alpha \otimes 1)_\alpha$, the weight filtration $W_\bullet \mathcal{H}_A$ to the weight filtration $W_\bullet \Lambda \otimes A$ and the Hodge filtration $F^\bullet \mathcal{H}_A$ to the Hodge filtration $F^\bullet(\Lambda \otimes \mathbb{W}(\kappa)) \otimes_{\mathbb{W}(\kappa)} A$, defines a \bar{P} -torsor for the étale topology on $\text{Spec}(A)$.*

Proof If we forget the compatibility with the weight filtrations, this is the content of [16, Prop. 4.3.9]. The isomorphisms preserving tensors and Hodge filtrations are proven in loc. cit. to be a \tilde{P}_A -torsor I over $\text{Spec}(A)$. Let $J \subset I$ be the subscheme of isomorphisms preserving also the weight filtration. Take the base change of \tilde{S}^Σ via $\mathbb{W}(\kappa) \rightarrow \mathbb{W}(k)$ with k an algebraic closure of κ and replace A with the complete local ring \bar{A} of this base change at a point over s_0 . Since $A \rightarrow \bar{A}$ is faithfully flat, it suffices to show that J is a \bar{P} -torsor over an fpqc cover of $\text{Spec}(\bar{A})$. In order to simplify the notation we simply write A in place of \bar{A} . As now A is complete local with algebraically closed residue field, $I(A)$ is non empty and I is the trivial \tilde{P} -torsor. It suffices to show that $J(A)$ is non-empty as, then, J is the trivial \bar{P} -torsor over $\text{Spec}(\bar{A})$.

The tensors $(s_{\alpha, \text{dR}})_\alpha$ are horizontal for the logarithmic connection ∇ on \mathcal{H}_A . The residue of the connection defines the monodromy operator N_A on \mathcal{H}_A . This is an A -linear operator such that $N_A^2 = 0$, the kernel is $W_1 \mathcal{H}_A$ and the image is $W_0 \mathcal{H}_A$. Given an isomorphism $j: \mathcal{H}_A \cong \Lambda \otimes A$ in $I(A)$ the operator $N := j \circ N_A \circ j^{-1}$ annihilates the tensors $s_\alpha \otimes 1$ and, hence, lies in $\text{Lie } G(A)$. The weight filtration $j(W_\bullet \mathcal{H}_A)$, defined by the image of $W_\bullet \mathcal{H}_A$ via j , coincides with $0 \subset \text{Im}(N) \subset \text{Ker}(N) \subset \Lambda \otimes A$ and, in particular, it is split by a cocharacter $w_A: \mathbb{G}_{m,A} \rightarrow G_A$; c.f. the proof of [16, Lemma 3.5.3].

Lift s_0 to a $\mathbb{W}(k)[1/p]$ -point s of \tilde{S}^Σ lying in the same boundary component as s_0 . Let \hat{A}_s be the complete local ring of \tilde{S}^Σ at s . We claim that $J|_{\hat{A}_s}$ is a torsor over $\text{Spec}(\hat{A}_s)$. Take an embedding $\mathbb{W}(k)[1/p] \subset \mathbb{C}$. Let \hat{B}_s be the complete local ring of $\tilde{S}^\Sigma \otimes_{\mathbb{W}(\kappa)} \mathbb{C}$ at the \mathbb{C} -valued point associated to s . By construction we have a variation of mixed Hodge structures over $\tilde{S}^\Sigma \otimes_{\mathbb{W}(\kappa)} \mathbb{C}$ whose de Rham realization over \hat{B}_s is the base change of \mathcal{H}_A and whose underlying \mathbb{Q} -vector space is $\Lambda \otimes \mathbb{Q}$ so that $J(\hat{B}_s)$ is non-empty; see the discussion in [16, Sect. 2.1]. The extension $\hat{A}_s \rightarrow \hat{B}_s$ is faithfully flat as it is local and \hat{B}_s is obtained from \hat{A}_s by a flat base change $\mathbb{W}(k) \rightarrow \mathbb{C}$, localization and completion. This proves that the cocharacters w_A and w of G are conjugate over A_s . It follows from [26, Prop. 3.2.1] that the conjugacy classes of cocharacters of G_A

are the disjoint union of copies of $\text{Spec}(A)$. We deduce that w_A and w are conjugate over an fpqc cover of A , i.e., the filtrations $j(W_\bullet \mathcal{H}_A)$ and $W_\bullet \Lambda$ are conjugate under G_A over an fpqc cover of A . Over this J becomes a \overline{P} -torsor as wanted. \square

3.3 The proof

Consider the parabolic subgroups $\overline{P}_\kappa \subset P \subset H = G_\kappa$ which are the reduction of $\overline{P} \subset \tilde{P} \subset G_{\mathbb{W}(\kappa)}$. Fix a Borel subgroup $B \subset \overline{P}_\kappa$ and a maximal torus T . Let $H'' := G''_{\mathbb{F}_p}$, resp. $H' = G'_{\mathbb{F}_p}$ be the reductive quotient of the mod p -reduction $R_{\mathbb{F}_p}$ of R , resp. of the mod p -reduction $N_{\mathbb{F}_p}$ of N . Recall that $H' \subset H''$ is a normal subgroup.

Let χ' be the mod p reduction of $\tilde{\mu}'$. It is a cocharacter of $H' := G'_{\mathbb{F}_p}$ and, hence, also of H'' defined over κ . By functoriality we get a map

$$\gamma: H' - \text{Zip}^{\chi'} \longrightarrow H'' - \text{Zip}^{\chi''}.$$

The cocharacter χ' defines a pair of opposite parabolic subgroups P'_\pm and P''_\pm of H'_κ and H''_κ respectively. Let $\mathcal{Z}' = (P'_+, (P'_-)^{(p)}, \varphi)$ and $\mathcal{Z}'' = (P''_+, (P''_-)^{(p)}, \varphi)$ be the two associated zip data. Recall that $H' - \text{Zip}^{\chi'} \cong [E_{\mathcal{Z}'} \setminus H'_\kappa]$ and $H'' - \text{Zip}^{\chi''} \cong [E_{\mathcal{Z}''} \setminus H''_\kappa]$. The inclusion $H' \subset H''$ identifies $P'_\pm = P''_\pm \cap H'_\kappa$ and induces a map of zip data $\mathcal{Z}' \rightarrow \mathcal{Z}''$ and a map of quotient stacks

$$\gamma: [E_{\mathcal{Z}'} \setminus H'_\kappa] \longrightarrow [E_{\mathcal{Z}''} \setminus H''_\kappa]$$

which provides a different realization of γ .

We define other zip data for H_k that will be relevant for our discussion. The first is $\mathcal{Z} := (P, P_-^{(p)}, \varphi)$. The second is $\overline{\mathcal{Z}} := (\overline{P}_\kappa, \overline{P}_{\kappa,-}^{(p)}, \varphi)$. Note that $E_{\overline{\mathcal{Z}}} \subset E_{\mathcal{Z}}$ so that we get a map

$$\epsilon: [E_{\overline{\mathcal{Z}}} \setminus H_k] \longrightarrow [E_{\mathcal{Z}} \setminus H_k].$$

Identifying H''_k with the standard Levi factor of $R_{\mathbb{F}_p}$ (for our choices of the Borel B and the torus T), we have an inclusion $H''_k \subset R_\kappa \subset H_k$ and $P''_+ = \overline{P}_\kappa \cap H''_k$ and $P''_- = \overline{P}_{-\kappa} \cap H''_k$. Then we get a map of stacks

$$\nu: [E_{\mathcal{Z}''} \setminus H''_k] \longrightarrow [E_{\overline{\mathcal{Z}}} \setminus H_k].$$

Lemma 3.4 *The morphisms γ, ϵ, ν and, hence, the composite $\tau := \epsilon \circ \nu \circ \gamma$ are smooth.*

Proof We start with γ . Consider the push forward $F := E_{\mathcal{Z}'} \setminus (E_{\mathcal{Z}''} \times_k H'_k)$ of the inclusion $E_{\mathcal{Z}'} \rightarrow E_{\mathcal{Z}''}$ via the group homomorphism $E_{\mathcal{Z}'} \rightarrow H'_k$. Here $E_{\mathcal{Z}'}$ acts on $E_{\mathcal{Z}''} \times_k H'_k$ via $(\gamma, (\alpha, \beta)) \mapsto (\alpha\gamma^{-1}, \gamma\beta)$. The projection $E_{\mathcal{Z}''} \times_k H'_k \rightarrow H'_k$ composed with $H'_k \rightarrow [E_{\mathcal{Z}'} \setminus H'_k]$ defines a map $F \rightarrow [E_{\mathcal{Z}'} \setminus H'_k]$. The inclusion $H'_k \subset H''_k$ induces

a $E_{\mathcal{Z}''}$ -equivariant map $\delta: F \rightarrow H''_k$ where $E_{\mathcal{Z}''}$ act on F through left multiplication on the first factor of $E_{\mathcal{Z}''} \times_k H'_k$. We then have a commutative diagram

$$\begin{array}{ccc} F & \xrightarrow{\delta} & H''_k \\ \downarrow & & \downarrow \\ [E_{\mathcal{Z}'} \setminus H'_k] & \xrightarrow{\gamma} & [E_{\mathcal{Z}''} \setminus H''_k]. \end{array}$$

For any scheme S , any S -valued point of the fibre product is, locally for the fppf topology, defined by the trivial $E_{\mathcal{Z}'}$ -torsor together with the map $E_{\mathcal{Z}'} \rightarrow H'_k$ given by right multiplication by $\beta \in H'_k(S)$, the trivial $E_{\mathcal{Z}''}$ -torsor together with the map $E_{\mathcal{Z}''} \rightarrow H''_k$ given by right multiplication by $\epsilon \in H''_k(S)$ and an element $\alpha \in E_{\mathcal{Z}''}(T)$ such that $\epsilon = \alpha\beta \in H''_k(S)$. Hence, F is the fiber-product.

In order to prove that γ is smooth, it suffices to show that the map δ is smooth. Recall that H' is normal in H'' with reductive quotient H''' . Hence $P'_\pm = P''_\pm \cap H'_k$ is normal in P''_\pm with quotient H'''_κ as χ' defines the trivial character of H''' . Similarly $(P'_-)^{(p)}$ is normal in $(P''_-)^{(p)}$ with quotient $(H'''_\kappa)^{(p)}$. We deduce that F , via the right multiplication by H'_k , is an H'_k -torsor over $E_{\mathcal{Z}'} \setminus E_{\mathcal{Z}''}$ which is identified with H'''_κ (embedded in $H'''_\kappa \times_k (H'''_\kappa)^{(p)}$ via the graph of Frobenius). Both F and H''_k are right H'_k -torsors over H'''_κ . In particular, also F is a smooth k -scheme and the smoothness of δ is equivalent to the surjectivity of δ is on tangent spaces at closed points. The map $H'''_\kappa \cong F/H'_k \rightarrow H''_k/H'_k \cong H'''_\kappa$, induced by δ , is Lang isogeny $h \mapsto h(h^p)^{-1}$ which is surjective and separable. Hence δ induces an isomorphism on tangent spaces as wanted.

As the morphism ϵ is a quotient morphism of smooth stacks, it is smooth.

We are left to prove the smoothness of ν . Let $M := E_{\mathcal{Z}''} \setminus (E_{\overline{\mathcal{Z}}} \times_k H''_k)$ be the push forward of the inclusion $E_{\mathcal{Z}''} \rightarrow E_{\overline{\mathcal{Z}}}$ via the group homomorphism $E_{\mathcal{Z}''} \rightarrow H''_\kappa$. The inclusion $H'' \subset H$ defines a $E_{\overline{\mathcal{Z}}}$ -equivariant map $\tau: M \rightarrow H_k$ and arguing as before we get that M is the fiber product

$$\begin{array}{ccc} M & \xrightarrow{\tau} & H_k \\ \downarrow & & \downarrow \\ [E_{\mathcal{Z}''} \setminus H''_k] & \xrightarrow{\nu} & [E_{\overline{\mathcal{Z}}} \setminus H_k]. \end{array}$$

The smoothness of ν is then equivalent to the smoothness of τ . As τ is a morphism of schemes smooth over k , it suffices to show that it is surjective on tangent spaces at k -valued points and, by right translation by $E_{\overline{\mathcal{Z}}} \times_k H''_k$, it suffices to show this at the 0-sections. Recall that the parabolic subgroup R of $G_{\mathbb{Z}_p}$ is defined over \mathbb{Z}_p and it has unipotent radical $U_{\mathbb{Z}_p}$ also defined over \mathbb{Z}_p . Let $U_{\mathbb{F}_p}$ be its reduction modulo p . Then $R_{\mathbb{F}_p} \cap R_{\mathbb{F}_p,-} = H''$ and it follows that $\text{Lie}H = \text{Lie}H'' \oplus \text{Lie}U_{\mathbb{F}_p} \oplus \text{Lie}U_{\mathbb{F}_p,-}$. Notice that $U_{\mathbb{F}_p}$ is contained in the unipotent radical of \overline{P}_κ . Similarly $U_{\mathbb{F}_p,-}$ is the unipotent radical of $(R_{\mathbb{F}_p,-})^{(p)}$ and is contained in the unipotent radical of $\overline{P}_{\kappa,-}^{(p)}$. Hence, $\text{Lie}U_{\mathbb{F}_p} \oplus \text{Lie}U_{\mathbb{F}_p,-}$ is contained in the image of $\text{Lie}E_{\overline{\mathcal{Z}}}$. Thus τ induces a surjective map on tangent spaces at the 0-sections as wanted. \square

Remark 3.5 In the proof of Lemma 3.4 the fact that the weight cocharacter is defined over \mathbb{Q} , so that $R_{\mathbb{Z}_p}$ and its unipotent radical are defined over \mathbb{Z}_p , is the key point of the proof.

Over S' one has an abelian scheme B . The covariant Dieudonné module of its mod p reduction B_0 defines an associated map

$$\zeta' : S'_0 \rightarrow H' - \text{Zip}^{\chi'}$$

which is smooth by [26, Cor. 3.12]. As $\widetilde{S}_0^\Sigma \rightarrow S'_0$ is smooth, the composite map $\zeta'^\Sigma : \widetilde{S}_0^\Sigma \rightarrow S'_0 \rightarrow H' - \text{Zip}^{\chi'}$ is smooth as well.

Proposition 3.6 *The map $\zeta^\Sigma : \text{Spec}(A_0) \rightarrow H - \text{Zip}^\chi$ of Lemma 2.13 is the restriction to $\text{Spec}(A_0)$ of the morphism $\tau \circ \zeta'$ with $\tau := \gamma \circ \epsilon \circ \nu$. In particular, it is surjective on tangent spaces at the closed point.*

Proof The last statement follows from the fact that $\gamma \circ \epsilon \circ \nu \circ \zeta'$ is smooth as it is the composite of smooth morphisms by Lemma 3.4.

As remarked in Sect. 3.1 the weight filtration on \mathcal{H}_A arises from a filtration by Frobenius crystals, that is, Frobenius preserves the weight filtration. We conclude from Proposition 3.3 that the element $g_{t,0}$ of Lemma 2.13 lies in $R_{\mathbb{F}_p}(A_0)$. Let $g''_{t,0}$ be the image of $g_{t,0}$ in the quotient H'' of $R_{\mathbb{F}_p}$. The image of the class of $g''_{t,0}$ via ν coincides with the class of $g_{t,0}$ as the kernel $U_{\mathbb{F}_p}$ of the quotient map $R_{\mathbb{F}_p} \rightarrow H''$ is contained in the unipotent radical of \overline{P}_κ and hence in $E_{\overline{\mathbb{Z}}}$ (as explained in the proof of Lemma 3.4). Let

$$\zeta'' : \text{Spec}(A_0) \longrightarrow [E_{\overline{\mathbb{Z}}''} \setminus H''_k]$$

be the map defined by $g''_{t,0} \in H''(A_0)$. Then, the map $\zeta^\Sigma : \text{Spec}(A_0) \rightarrow H - \text{Zip}^\chi$ factors as

$$\zeta^\Sigma = \epsilon \circ \nu \circ \zeta''.$$

Possibly extending the residue field of A we may assume that the torus T and the lattice Y of the degenerating 1-motive M_U^\vee over A are trivial (see Sect. 3.1 for the notation). Taking generators of $\text{Lie}(T)$ coming from a basis of cocharacters of T and a \mathbb{Z} -basis of Y , that defines a basis of the universal extension of Y , Frobenius on $W_0\mathcal{H}_A$ becomes multiplication by p and Frobenius on $\text{Gr}_2\mathcal{H}_A$ is the identity matrix. We conclude that, after a finite étale extension of A , we can assume that the projection of $g''_{t,0}$ onto $\text{Aut}(W_0\mathcal{H}_{A_0}) \times \text{Aut}(\text{Gr}_2\mathcal{H}_{A_0})$ is the identity so that it lies in $\text{Aut}(\text{Gr}_2\mathcal{H}_{A_0})$, the automorphism group of the Dieudonné module of B_0 base changed to $\text{Spec}(A_0)$. Hence, the morphism ζ'' factors via ζ' restricted to $\text{Spec}(A_0)$ composed with γ as wanted. \square

3.4 Ekedahl–Oort strata at the boundary

It follows from [16, Thm. 5.3.1] that there exists a minimal compactification S_0^* and a map $\pi_0: S_0^\Sigma \rightarrow S_0^*$ such that the structure at the boundary is induced by the fibration described in Sect. 3.1 (here the index 0 in S_0^Σ denotes as usual the mod p special fiber). More precisely, the boundary components of S_0^* are a disjoint union of the special fiber of integral models of lower dimensional smooth Shimura varieties of the type S'_0 , as considered in Sect. 3.1, indexed by so called cusp label representatives. For any such component the map $\pi_0^{-1}(S'_0) \rightarrow S'_0$ is isomorphic to the map $\partial\tilde{S}_0^\Sigma \rightarrow S_0$ defined by restricting the morphism $\tilde{S}_0^\Sigma \rightarrow S_0$ of Sect. 3.1 to a closed subscheme $\partial\tilde{S}_0^\Sigma \subset \tilde{S}_0^\Sigma$. Then, we have the following description of the Ekedahl–Oort stratification on the boundary of S_0^Σ :

Corollary 3.7 *For every boundary component S'_0 of S_0^* as above, the map ζ^Σ restricted to $\pi_0^{-1}(S'_0)$ is the composite of*

- (i) *the projection $\pi_0^{-1}(S'_0) \rightarrow S'_0$;*
- (ii) *the map $\zeta': S'_0 \rightarrow H' - \text{Zip}^{X'}$ defined in [26];*
- (iii) *the map $\tau: H' - \text{Zip}^{X'} \rightarrow H - \text{Zip}^X$ of Lemma 3.4.*

In particular, ζ^Σ restricted to $\pi_0^{-1}(S'_0)$ is constant on the geometric fibers of π_0 and the induced map on underlying topological spaces

$$|\zeta^\Sigma|: |\pi_0^{-1}(S'_0)| \rightarrow |H - \text{Zip}^X|$$

has open image and does not contain the closed point of $|H - \text{Zip}^X|$.

Proof Statements (i)–(iii) follow from Proposition 3.6. The map τ is smooth and, hence, its image on the underlying topological spaces is open, i.e., it is closed under generalization. If it contained the closed point, it would be surjective and any maximal chain of points of $|H - \text{Zip}^X|$ would be the image of a maximal chain of points of $|H' - \text{Zip}^{X'}|$. This can not be as any maximal chain of points in $|H' - \text{Zip}^{X'}|$ has length equal to the dimension of S'_0 which is strictly smaller than the dimension of S_0 which is equal to the length of a maximal chain of points of $|H - \text{Zip}^X|$. This implies the last statement. □

3.5 A result on zips in mixed characteristics

The cocharacter $\tilde{\mu}$ of $G_{\mathbb{W}(k)}$, whose reduction is χ , defines two parabolic subgroups $\tilde{P} = \tilde{P}_+$ and \tilde{P}_- of $G_{\mathbb{W}(k)}$. The reduction of \tilde{P}_\pm is P_\pm . Let $\tilde{T} \subset \tilde{B} \subset G_{\mathbb{Z}_p}$ be a Borel subgroup and a maximal torus, defined over \mathbb{Z}_p . By replacing $\tilde{\mu}$ by a conjugate cocharacter, that might be defined over an extension $\mathbb{W}(k) \subset \mathbb{W}(k')$, we may and will assume that \tilde{P}_+ and \tilde{P}_- contain \tilde{B} . There is a common Levi subgroup $\tilde{L} = \tilde{P}_+ \cap \tilde{P}_-$. Finally we let $\tilde{Q} \subset G_{\mathbb{W}(k)}$ be the parabolic subgroup $\tilde{P}_-^{(p)}$ defined by base change of $\tilde{P}_- \subset G_{\mathbb{W}(k)}$ via Frobenius using that $G_{\mathbb{W}(k)}^{(p)} = G_{\mathbb{W}(k)}$ as $G_{\mathbb{W}(k)}$ is defined over \mathbb{Z}_p .

Suppose we are given a $\mathbb{W}(k)$ -algebra D endowed with a lift of Frobenius φ . Let $\Phi: G_{\mathbb{W}(k)}(D) \rightarrow G_{\mathbb{W}(k)}(D)$ be the Frobenius map $g \mapsto g^{(p)}$; using the embedding

$G_{\mathbb{Z}_p} \subset \mathrm{GSp}_\Lambda \cong \mathrm{GSp}_{2g}$ we view $g \in G_{\mathbb{W}(k)}(D)$ as a $(2g) \times (2g)$ -matrix with coefficients in D and we let $\Phi(g) = g^{(p)}$ be the matrix where we apply φ to the coefficients. It is still an element of $G_{\mathbb{W}(k)}(D)$ since the equations defining the closed subgroup scheme $G_{\mathbb{Z}_p} \subset \mathrm{GSp}_{2g, \mathbb{Z}_p}$ have coefficients in \mathbb{Z}_p . Define $\tilde{E}_{\mathcal{Z}_{\mathrm{EO}}}(D)$ to be the subgroup of pairs $(g, h) \in G_{\mathbb{W}(k)}(D) \times G_{\mathbb{W}(k)}(D)$ with $g \in \tilde{P}(D)$, $h \in \tilde{Q}(D)$ and such that the map

$$\bar{\Phi}: (\tilde{P}(D)/R_u \tilde{P}(D)) = \tilde{L}(D) \longrightarrow (\tilde{L})^{(p)}(D) = (\tilde{Q}(D)/(R_u \tilde{Q}(D)))$$

sends the class of g to the class of h . It acts on the left on $G_{\mathbb{W}(k)}(D)$ via $(a, b) \cdot g = agb^{-1}$.

Recall the notation of Sect. 3.1; A is the complete local ring of S^Σ at a closed point x_0 . Frobenius on \mathcal{H}_A defines an element $g_t \in G_{\mathbb{W}(k)}(A)$. Assume that $\zeta^\Sigma(x_0)$ is the class of $\tilde{w}\tilde{z}^{-1}$ for $w \in {}^I W$. See Lemma 2.11. For every $\mathbb{W}(k)$ -point x of A we let \mathcal{H}_x be the pull-back of \mathcal{H}_A . It is a $\mathbb{W}(k)$ -module, endowed with a weight filtration, Hodge filtration and a Frobenius linear map F . Proceeding as in the discussion before Lemma 2.13 Frobenius and a splitting of the Hodge filtration produce an element $g_x \in G_{\mathbb{W}(k)}(\mathbb{W}(k))$. Here and in what follows, abusing notation, we identify the Weyl group of $G_{\mathbb{W}(k)}(\mathbb{W}(k))$ and of $G_{\mathbb{F}_p}(k) = H(k)$ with respect to the torus $\tilde{T}(\mathbb{W}(k))$, resp. $T(k)$. Then,

Proposition 3.8 *There exists a $\mathbb{W}(k)$ -point x of A , that reduces to x_0 modulo p and generically lies in $S \subset S^\Sigma$, such that the class of g_x in $\tilde{E}_{\mathcal{Z}_{\mathrm{EO}}}(\mathbb{W}(k)) \setminus G_{\mathbb{W}(k)}(\mathbb{W}(k))$ is $\tilde{w}\tilde{z}^{-1}$.*

Proof First of all we claim that it suffices to construct x without assuming that it generically lies in S . For any $\mathbb{W}(k)$ -valued point y of A , Proposition 3.3 and the fact that Frobenius on \mathcal{H}_y preserves the weight filtration imply that $g_y \in R_{\mathbb{Z}_p}(\mathbb{W}(k))$. The unipotent radical $U_{\mathbb{Z}_p}$ of $R_{\mathbb{Z}_p}$ is contained in the unipotent radical of P so that the class of g_y depends only on the class \bar{g}_y in the reductive quotient $G'_{\mathbb{Z}_p}(\mathbb{W}(k)) = R_{\mathbb{Z}_p}(\mathbb{W}(k))/U_{\mathbb{Z}_p}(\mathbb{W}(k))$. Since Frobenius on $W_0\mathcal{H}_y$ with respect to a basis coming from a basis of cocharacters of the torus $y^*(T)$ can be taken to be multiplication by p and Frobenius on $\mathrm{Gr}_2\mathcal{H}_y$ with respect to a basis coming from a basis of $y^*(Y)$ is the identity matrix, in fact \bar{g}_y can be taken in $G'_{\mathbb{Z}_p}(\mathbb{W}(k))$ and this depends only on Frobenius defined on the factor $\mathrm{Gr}_1\mathcal{H}_y$ which is the Frobenius crystal over $\mathbb{W}(k)$ defined by pulling back the abelian scheme B via y . In particular, if we let $y' \in S'(\mathbb{W}(k))$ be the image of y in the fibration of Sect. 3.1 and if we take any $\mathbb{W}(k)$ -valued point v of A in the fiber of y' then the class of g_y and of g_v in $\tilde{E}_{\mathcal{Z}_{\mathrm{EO}}}(\mathbb{W}(k)) \setminus G(\mathbb{W}(k))$ are the same. As we have v 's in the fiber over y' that generically lie in S , the claim is proven.

We are left to prove the Proposition dropping the requirement that x lies generically in S . We construct inductively $\mathbb{W}(k)$ -points $(y_n)_{n \in \mathbb{N}}$ such that $y_0 \equiv x_0$ modulo p and $y_{n+1} \equiv y_n$ modulo p^{n+1} for every n and $g_{y_n} \equiv \tilde{w}\tilde{z}^{-1}$ in $\tilde{E}_{\mathcal{Z}_{\mathrm{EO}}}(\mathbb{W}_{n+1}(k)) \setminus G_{\mathbb{W}(k)}(\mathbb{W}_{n+1}(k))$. As the maps $\alpha_n: \tilde{E}_{\mathcal{Z}_{\mathrm{EO}}}(\mathbb{W}_{n+1}(k)) \rightarrow E_{\mathcal{Z}_{\mathrm{EO}}}(\mathbb{W}_n(k))$ and $\beta_n: G_{\mathbb{W}(k)}(\mathbb{W}_{n+1}(k)) \rightarrow G_{\mathbb{W}(k)}(\mathbb{W}_n(k))$ are surjective, the limit point $x = \lim_{n \rightarrow \infty} y_n$ satisfies the requirements.

Take y_0 to be any lift of x_0 . Assume that y_n is constructed. Then, possibly after multiplying $g_n := g_{y_n}$ by an element of $\tilde{E}_{\mathcal{Z}_{EO}}(\mathbb{W}_{n+1}(k))$ we can assume that $g_n \tilde{w}^{-1}$ lies in the kernel of β_n which is isomorphic to $\text{Lie}H_k$ (recall that H is the reduction of $G_{\mathbb{Z}_p}$ modulo p). Thus such class defines an obstruction element $\text{Obs}(y_n) \in \text{Lie}H_k$

Let $\text{Lie}(E_{\mathcal{Z}_{EO}})$ be the subset of pairs $(a, b) \in \text{Lie}(P) \otimes k \oplus \text{Lie}(Q) \otimes k$ such that the map $1 \otimes \varphi: \text{Lie}L \otimes k \rightarrow \text{Lie}(L)^{(p)} \otimes k$ sends the class of a modulo $\text{Lie}R_u P \otimes k$ to the class of b modulo $\text{Lie}R_u Q \otimes k$. The kernel of α_n is isomorphic to $\text{Lie}(E_{\mathcal{Z}_{EO}})$. Write $\text{Lie}(P) = \text{Lie}(L) \oplus \text{Lie}(R_u P)$ and $\text{Lie}(Q) = \text{Lie}(L^{(p)}) \oplus \text{Lie}(R_u Q)$. For $a \in \text{Lie}(P)$ write $a = a_0 + a_1$ for the corresponding decomposition. Similarly for $b \in \text{Lie}(Q)$ write $b = b_0 + b_1$. Then $(a, b) \in \text{Lie}(E_{\mathcal{Z}_{EO}}(\mathbb{W}(k)))$ if and only if $\varphi(a_0) = b_0$. As Frobenius on Lie algebras is the zero map, this amounts to require that $b_0 = 0$, that is $\text{Lie}(E_{\mathcal{Z}_{EO}}(k))$ consist of pairs (a, b) such that $b_0 = 0$. Consider the map of Lie algebras over k :

$$\nu: \text{Lie}(E_{\mathcal{Z}_{EO}}) = \text{Lie}(L) \oplus (R_u P) \oplus \text{Lie}(R_u Q) \rightarrow \text{Lie}G_k, \quad (m, n, z) \mapsto m + n - z.$$

The element $\text{Obs}(y_n)$ lies in the image of ν if and only if after multiplying $g_n := g_{y_n}$ by an element of $\tilde{E}_{\mathcal{Z}_{EO}}(\mathbb{W}_{n+1}(k))$ we have $g_{y_n} \equiv \tilde{w}z^{-1}$ in $\tilde{E}_{\mathcal{Z}_{EO}}(\mathbb{W}_{n+1}(k)) \setminus G_{\mathbb{W}(k)}(\mathbb{W}_{n+1}(k))$. In this case we can take $y_{n+1} = y_n$.

Recall that by construction we have a lift $\text{Spec}(A_0) \rightarrow H_k$ of the map $\zeta^\Sigma: S_0^\Sigma \rightarrow H - \text{Zip}^\lambda$. It induces a map from the tangent space T_{x_0} of S_0^Σ at x_0 to the tangent space T_{H, c_0} of H_k at the image c_0 of the closed point of A_0 , that we identify with $\text{Lie}H_k$ by translating by c_0^{-1} . The smoothness of ζ^Σ in Theorem 3.1 implies that the induced map

$$\theta: \text{Lie}(E_{\mathcal{Z}_{EO}}) \times T_{x_0} \rightarrow \text{Lie}H_k$$

is surjective. The possible lifts of y_n to a $\mathbb{W}_{n+1}(k)$ -valued point of A is a principal homogeneous space under T_{x_0} as we assume that our toroidal compactification S^Σ is smooth. Let y'_n be a lift corresponding to an element $s \in T_{x_0}$. Then $\text{Obs}(y'_n) = \text{Obs}(y_n) + \theta((0, s))$. By the surjectivity of θ we can then find an s , and hence a corresponding lift $y_{n+1} = y'_n$, and an element $b \in \text{Lie}(E_{\mathcal{Z}_{EO}})$ such that $\text{Obs}(y'_n) = \theta((b, 0))$. This implies that after multiplying $g_{n+1} := g_{y_{n+1}}$ by an element in the kernel of α_n we have that $g_{y_{n+1}} \equiv \tilde{w}z^{-1}$ in $\tilde{E}_{\mathcal{Z}_{EO}}(\mathbb{W}_{n+1}(k)) \setminus G_{\mathbb{W}(k)}(\mathbb{W}_{n+1}(k))$. This concludes the proof of the inductive step and of the Proposition. \square

4 On the mod p Hodge–Tate period map

We describe the map $q_2: \mathcal{S}(p^\infty) \rightarrow \text{Fl}_{G, \mu^{-1}, 0}$ using certain formal models introduced in [18].

Thanks to [6, Thm. IV.6.7] there exists a smooth toroidal compactification $S_{K_g, \mathbb{Q}_p}^{\Sigma_g}(p^n)$ of the full level p^n Siegel modular scheme $S_{K_g, \mathbb{Q}_p}(p^n) \rightarrow S_{K_g, \mathbb{Q}_p}$, associated to the finite admissible cone decomposition Σ_g fixed in (2.12). It defines a

$\mathrm{Sp}_{2g}(\mathbb{Z}/p^n\mathbb{Z})$ -cover

$$S_{K_g, \mathbb{Q}_p}^{\Sigma_g}(p^n) \rightarrow S_{K_g, \mathbb{Q}_p}^{\Sigma_g}.$$

Let T be the p -adic Tate module of the universal abelian scheme over $S_{\mathbb{Q}_p}$. One has Tate tensors $(s_{\mathrm{et}, \alpha})_\alpha$ in the tensor algebra of T . Over the fiber product $S_{\mathbb{Q}_p} \times_{S_{K_g, \mathbb{Q}_p}^{\Sigma_g}} S_{K_g, \mathbb{Q}_p}^{\Sigma_g}(p^n)$ the étale sheaf $T/p^n T \cong \Lambda/p^n \Lambda$ is trivialized and it extends to a constant sheaf $T_n \cong \Lambda/p^n \Lambda$ over $S_{\mathbb{Q}_p}^{\Sigma} \times_{S_{K_g, \mathbb{Q}_p}^{\Sigma_g}} S_{K_g, \mathbb{Q}_p}^{\Sigma_g}(p^n)$. Let $S_{\mathbb{Q}_p}^{\Sigma}(p^n)$ be the closed subscheme

$$S_{\mathbb{Q}_p}^{\Sigma}(p^n) \subset S_{\mathbb{Q}_p}^{\Sigma} \times_{S_{K_g, \mathbb{Q}_p}^{\Sigma_g}} S_{K_g, \mathbb{Q}_p}^{\Sigma_g}(p^n)$$

such that the isomorphism $t_n : T_n \cong \Lambda/p^n \Lambda$ sends the mod p^n -reduction of the tensors $(s_{\mathrm{et}, \alpha})_\alpha$ in the tensor algebra of T_n to the tensors $(s_\alpha)_\alpha$ of $\Lambda/p^n \Lambda$.

Let $S_{K_g}^{\Sigma_g}(p^n)$ be the normalization of the toroidal compactification $S_{K_g}^{\Sigma_g}$ of the Siegel modular scheme in $S_{K_g, \mathbb{Q}_p}^{\Sigma_g}(p^n)$. Similarly let $S(p^n)$, resp. $S^\Sigma(p^n)$ be the normalization of S , resp. S^Σ in $S_{\mathbb{Q}_p}^{\Sigma}(p^n)$. It is finite over S , resp. S^Σ by construction. By [18, Prop. 1.5] we have a map

$$d \log_n : T_n^\vee(1) \rightarrow \Omega \otimes (\mathcal{O}_{S^\Sigma(p^n)}/p^n \mathcal{O}_{S^\Sigma(p^n)})$$

given as follows. First of all the Cartier dual of the p^n -torsion of the universal semiabelian scheme G over $S_{\mathbb{Q}_p}^{\Sigma_g}(p^n)$ is a quotient of $T_n^\vee(1)$ through which $d \log_n$ factors. Any such element extends uniquely to an element $\gamma : G[p^n] \rightarrow \mathbb{G}_{m, S^\Sigma(p^n)}$, defined over the fppf topology of $S^\Sigma(p^n)$. The pull-back of the standard invariant differential on \mathbb{G}_m defines an element of the invariant differentials of $G[p^n]$ that coincide with the mod p^n -reduction of the invariant differentials of G . Finally we use [6, Thm. 1.1] to identify the sheaf of invariant differentials Ω_G of the universal semiabelian scheme G over S^Σ , used in [18], with the first piece of the Hodge filtration $\Omega \subset H_{\mathrm{dR}}^1$ of the canonical extension H_{dR}^1 described in Sect. 2.3 and used in [7]. We can now state [18, Prop. 1.13] and [18, Sect. 2.3.1].

Denote by $S_0^\Sigma(p^n)$ and $S_{K_g, 0}^{\Sigma_g}(p^n)$ the mod p -reductions of $S^\Sigma(p^n)$ and $S_{K_g}^{\Sigma_g}(p^n)$ respectively. Set $\overline{S}^\Sigma(p^n)$ and $\overline{S}_{K_g}^{\Sigma_g}(p^n)$ to be the formal schemes defined by completing $S^\Sigma(p^n)$ along $S_0^\Sigma(p^n)$, resp. of $S_{K_g}^{\Sigma_g}(p^n)$ along $S_{K_g, 0}^{\Sigma_g}(p^n)$. Set $\tau : S^\Sigma(p^n) \rightarrow S_{K_g}^{\Sigma_g}(p^n)$ be the morphism induced by the map $S^\Sigma(p^n) \rightarrow S_{K_g}^{\Sigma_g}(p^n)$. Let ζ_p a primitive p -th root of unity.

Proposition 4.1 [18]. *Given integers $n \geq \frac{g}{p-1} + 1$ and $1 \leq m \leq n - 1$ there exist:*

- (i) *admissible normal formal schemes $\overline{S}^\Sigma(p^n)$ and $\overline{S}_{K_g}^{\Sigma_g}(p^n)$, projective morphisms $\overline{S}^\Sigma(p^n) \rightarrow S^\Sigma(p^n)$ and $\overline{S}_{K_g}^{\Sigma_g}(p^n) \rightarrow S_{K_g}^{\Sigma_g}(p^n)$ inducing an isomorphism on the*

adic generic fiber and a morphism $\bar{\tau}: \bar{S}^\Sigma(p^n) \rightarrow \bar{S}_{K_g}^{\Sigma_g}(p^n)$ making the following diagram commute:

$$\begin{CD} \bar{S}^\Sigma(p^n) @>\bar{\tau}>> \bar{S}_{K_g}^{\Sigma_g}(p^n) \\ @VVV @VVV \\ S^\Sigma(p^n) @>\tau>> S_{K_g}^{\Sigma_g}(p^n); \end{CD}$$

(ii) a $\mathcal{O}_{\bar{S}^\Sigma(p^n)}$ -submodule $\Omega^{mod} \subset \Omega$, locally free of rank g , such $(\zeta_p - 1)\Omega \subset \Omega^{mod}$.

There is a similar statement over $\bar{S}_{K_g}^{\Sigma_g}(p^n)$ and the sheaves over $\bar{S}^\Sigma(p^n)$ are obtained from those on $\bar{S}_{K_g}^{\Sigma_g}(p^n)$ by pull-back via the morphism $\bar{S}^\Sigma(p^n) \rightarrow \bar{S}_{K_g}^{\Sigma_g}(p^n)$ in (i);

(iii) a surjective map

$$HT_m: (\Lambda^\vee / p^m \Lambda^\vee) \otimes \mathcal{O}_{\bar{S}^\Sigma(p^n)}(1) \rightarrow \Omega^{mod} / p^m \Omega^{mod}$$

induced by the map $d \log_n$ and the identification $t_n: T_n \cong \Lambda / p^n \Lambda$. There is a similar statement over $\bar{S}_{K_g}^{\Sigma_g}(p^n)$ and the map HT_m over $\bar{S}^\Sigma(p^n)$ is obtained from the one on $\bar{S}_{K_g}^{\Sigma_g}(p^n)$ by base change via the morphism $\bar{S}^\Sigma(p^n) \rightarrow \bar{S}_{K_g}^{\Sigma_g}(p^n)$ in (i).

The sheaf Ω^{mod} is independent of n and m and for $1 \leq m' \leq m \leq n - 1$ the map $HT_{m'}$ is the reduction of HT_m modulo $p^{m'}$.

Let $\bar{S}_0^\Sigma(p^n)$ be the scheme over κ given by the mod p special fiber of $\bar{S}^\Sigma(p^n)$. Denote by $(\bar{S}_0^\Sigma(p^n))^{red}$ the reduced scheme associated to $\bar{S}_0^\Sigma(p^n)$

Recall that, using the Hodge cocharacter, we have defined a cocharacter χ of $(G_{\mathbb{Z}_p})_\kappa = H_\kappa$ and hence a decomposition $\Lambda \otimes \kappa = \Lambda_0 \otimes \kappa \oplus \Lambda_{-1} \otimes \kappa$. Taking the sequence $\Lambda_{-1} \otimes \kappa \subset \Lambda \otimes \kappa$ we get a base point in $Fl_{G, \mu^{-1}, 0}$, i.e., an identification as schemes over κ :

$$Fl_{G, \mu^{-1}, 0} \cong Fl_{H, \chi^{-1}} \cong P_- \setminus H_\kappa.$$

Corollary 4.2 *The subsheaf $Ker(HT_1)$ provided by Proposition 4.1 defines a map of schemes over κ :*

$$\pi_{HT,0}: (\bar{S}_0^\Sigma(p^n))^{red} \rightarrow P_- \setminus H_\kappa.$$

Proof Let P_g^\pm be the parabolic of $GSp_{2g, \kappa}$ induced by the cocharacter χ^\pm via the inclusion $G_\kappa \subset GSp_{2g, \kappa}$. Then $Ker(HT_1)$ provides a map $\bar{S}_0^\Sigma(p^n) \rightarrow P_g^- \setminus GSp_{2g, \kappa}$. The Hodge–Tate period map π_{HT} is functorial with respect to morphisms of Shimura data $(G, X) \rightarrow (GSp_{2g}, \mathbb{H}_g)$ by [5, Thm. 2.1.2, Thm. 2.1.3]. By [18] the integral

Hodge–Tate period map for the Siegel variety $\overline{S}_{K_g^\Sigma}(p^\infty) \rightarrow \text{Fl}_{\text{GSp}_{2g, \mu^{-1}}}$ is compatible with the one constructed by Scholze in [22]. Hence the image of $\overline{S}^\Sigma(p^\infty)$ is contained in $\text{Fl}_{G, \mu^{-1}}$. By the compatibilities of HT_1 over $\overline{S}^\Sigma(p^n)$ and $\overline{S}_{K_g^\Sigma}(p^n)$ respectively (see Proposition 4.1 (iii)) the composite morphism on special fibers

$$\overline{S}_0^\Sigma(p^n) \rightarrow \overline{S}_{K_g, 0}^{\Sigma_g}(p^n) \rightarrow \text{Fl}_{\text{GSp}_{2g, \mu^{-1}, 0}},$$

provided by $\text{Ker}(\text{HT}_1)$, factors through the image of the composite $\text{Fl}_{G, \mu^{-1}, 0} \rightarrow \text{Fl}_{\text{GSp}_{2g, \mu^{-1}, 0}} \cong P_g^- \backslash \text{GSp}_{2g, \kappa}$. Such image coincides with $P_- \backslash H_k$ as $P_- = H_\kappa \cap P_g^-$ and the map $P_- \backslash H_k \rightarrow P_g^- \backslash \text{GSp}_{2g, \kappa}$ is a closed immersion. The claim follows. \square

We need to restate the Corollary using Sect. 2.1.1 in order to express the fine Deligne–Lusztig stratification on $\text{Fl}_{H, \chi^{-1}, 0}$ in terms of the Weyl group W . Let I_- be the type opposite to the type I of P . Thanks to Lemma 2.8 we have an isomorphism as schemes over k

$$P_- \backslash H_k \cong P_{I_-} \backslash H_k, \quad h \mapsto \tilde{y}^{-1} h \tilde{z}.$$

Via the projection map

$$P_{I_-} \backslash H_k \rightarrow [E_{\mathcal{Z}_-} \backslash H_k]$$

we get a stratification on $P_- \backslash H_k$ indexed by the elements of $I_- W$ with closure relation described by the order relation \preceq ; given two elements w and $w' \in I_- W$ we write $w \preceq w'$ if and only if there exists $x \in W_{I_-}$ such that $xw\varphi(x)^{-1} \leq w'$ (for the Bruhat order on W). We let

$$\bar{\gamma}: \text{Fl}_{H_k, \chi^{-1}, 0} \cong P_- \backslash H_k \cong P_{I_-} \backslash H_k \longrightarrow [E_{\mathcal{Z}_-} \backslash H_k] \tag{4}$$

be the composite map.

For later purposes we prove the following result about the modulo p Hodge–Tate period map.

Proposition 4.3 *The map $\pi_{\text{HT}, 0}$ is surjective.*

Proof Let $\overline{S}^\Sigma(p^n) \rightarrow S^\Sigma(p^n)$ be the projective scheme over $\mathbb{W}(\kappa)$ and the morphism to $S^\Sigma(p^n)$ algebraizing the morphism of formal scheme $\overline{S}^\Sigma(p^n) \rightarrow S^\Sigma(p^n)$ of Proposition 4.1(i). Such morphism is projective and it an isomorphism over \mathbb{Q}_p . Consider the invertible sheaf L on $\overline{S}^\Sigma(p^n)$ algebraizing the sheaf $\det \Omega^{\text{mod}}$ on $\overline{S}^\Sigma(p^n)$ and let L_0 be the induced invertible sheaf on $\overline{S}_0^\Sigma(p^n)$. The sheaf L is base point free and after inverting p it coincides with the Hodge bundle by Proposition 4.1(ii). In particular, its restriction to the $S(p^n)_{\mathbb{Q}_p}$, identified with an open subscheme of $\overline{S}^\Sigma(p^n)$, is ample. Take a large enough power L^h which is very ample on $S(p^n)_{\mathbb{Q}_p}$. Choosing a

$\mathbb{W}(\kappa)$ -basis of global sections we get a map

$$\Psi : \overline{S}^\Sigma(p^n) \longrightarrow \mathbb{P}_{\mathbb{W}(\kappa)}^N.$$

As $\overline{S}^\Sigma(p^n)$ is projective over $\mathbb{W}(\kappa)$, the morphism Ψ is projective and its image $D^\Sigma(p^n)$ is closed. After inverting p , it is an immersion on $S(p^n)_{\mathbb{Q}_p}$ so that the image of Ψ , after inverting p , has dimension equal to the dimension d of $S_{\mathbb{Q}_p}$. In particular, $D_0^\Sigma(p^n) := D^\Sigma(p^n) \cap \mathbb{P}_\kappa^N$ is equidimensional of dimension d . Denote by Ψ_0 the morphism induced by Ψ on the mod p special fibers. Consider the Stein factorization

$$\pi_{\text{HT},0} : \overline{S}_0^\Sigma(p^n)^{\text{red}} \rightarrow Y \rightarrow P_- \setminus H_\kappa$$

with $\overline{S}_0^\Sigma(p^n)^{\text{red}} \rightarrow Y$ surjective with connected fibers and $Y \rightarrow P_- \setminus H_\kappa$ finite. Over $\overline{S}_0^\Sigma(p^n)^{\text{red}}$ the sheaf L_0 is the pull-back of the tautological invertible sheaf L^{taut} on $P_- \setminus H_\kappa$ via $\pi_{\text{HT},0}$. Hence the restriction of Ψ_0 to $\overline{S}_0^\Sigma(p^n)^{\text{red}}$ factors via Y . Let $Y' \subset Y$ be an irreducible component dominating an irreducible component of $D_0^\Sigma(p^n)$; it must have dimension $\geq d$.

The scheme $P_- \setminus H_\kappa$ is irreducible and has dimension equal to the dimension of the unipotent radical of P , which is also d . As the map $Y' \rightarrow P_- \setminus H_\kappa$ is finite, it must be surjective. Hence the map $\pi_{\text{HT},0} : \overline{S}_0^\Sigma(p^n)^{\text{red}} \rightarrow P_- \setminus H_\kappa$ is surjective and the claim follows. □

5 Proof of Theorem 1.1

Thanks to Corollary 4.2 the two maps of topological spaces $q_1 : S^{\text{ad}}(p^\infty) \rightarrow S_0^\Sigma$ and $q_2 : S^{\text{ad}}(p^\infty) \rightarrow \text{Fl}_{G,\mu^{-1},0}$ factor through a morphism $S^{\text{ad}}(p^\infty) \rightarrow \overline{S}_0^\Sigma(p^n)$, the projective morphism $u : \overline{S}_0^\Sigma(p^n) \rightarrow S_0^\Sigma$ and the map $\pi_{\text{HT},0} : (\overline{S}_0^\Sigma(p^n))^{\text{red}} \rightarrow \text{Fl}_{G,\mu^{-1},0} \cong P_- \setminus H_\kappa$. Hence it suffices to prove the result for $\overline{\mathbb{F}}_p$ -points of $\overline{S}_0^\Sigma(p^n)$. Consider the two maps

$$\zeta'_1 : (\overline{S}_k^\Sigma(p^n))^{\text{red}} \xrightarrow{u} S_k^\Sigma \xrightarrow{\zeta^\Sigma} H - \text{Zip}^\chi \cong [E_{Z_{\text{EO}}^\chi} \setminus H_k]$$

and

$$\zeta'_2 : (\overline{S}_k^\Sigma(p^n))^{\text{red}} \xrightarrow{\pi_{\text{HT},0}} P_- \setminus H_k \xrightarrow{\overline{\gamma}} [E_{Z_{\text{DL}}^{\chi^{-1}}} \setminus H_k],$$

where $\overline{\gamma}$ is defined in (4) and the index k in $\overline{S}_k^\Sigma(p^n)$ denotes the base change to k .

5.1 The setup

Take a closed point x_0 of $S_0^\Sigma(p^n)$. Suppose it is defined over a finite extension \mathbb{F} of κ . First lift the image x'_0 of x_0 in S_0^Σ to a $\mathbb{W}(\mathbb{F})$ -valued point x' of S^Σ such that the induced $\mathbb{W}(\mathbb{F})[p^{-1}]$ -point lies in S : this can be done as S^Σ is log-smooth over $\mathbb{W}(\kappa)$. Then lift x' , viewed as a $\mathbb{W}(\mathbb{F})$ -point of S^Σ , to a \mathcal{O}_K -valued point x of $\bar{S}^\Sigma(p^n)$ for some finite field extension $\mathbb{W}(\kappa)[1/p] \subset K$. Let $\mathbb{W}(\kappa)[1/p] \subset K_0 \subset K$ be the maximal unramified extension. We will assume that \mathbb{F} is the residue field of K_0 . This gives, in particular,

- (i) an abelian variety A over K_0 , with semistable model \tilde{A} over $\mathbb{W}(\mathbb{F})$ and special fiber $\tilde{A}_{\mathbb{F}}$ such that the Galois group of K acts trivially on $T_p A/p^n T_p A$;
- (ii) elements $(s_{\text{et},\alpha})_\alpha$ in the tensor algebra of étale cohomology of A , identified with the dual of the p -adic Tate module $T_p(A)$ of A , fixed by the Galois group of K_0 ;
- (iii) a trivlization $d: T_p(A)^\vee \cong \Lambda$, sending the tensors $(s_{\text{et},\alpha})_\alpha$ to the elements $(s_\alpha)_\alpha$;
- (iv) a finite and free $\mathbb{W}(\mathbb{F})$ -module $H_0 := (x')^*(\mathcal{H})$, pull back of the log crystal \mathcal{H} defined in Sect. 3.1 and identified with the contravariant log Dieudonné module $H_0(\tilde{A}_{\mathbb{F}})$ of $\tilde{A}_{\mathbb{F}}$ via [16, Prop. 1.3.5], endowed with a Frobenius linear operator φ and elements $(s_{\text{cris},\alpha})_\alpha$ in the tensor algebra of H_0 , fixed by φ ;
- (v) a saturated $\mathbb{W}(\mathbb{F})$ -submodule $\omega_0 \subset H_0$ such that $\varphi(\omega_0) \subset p\omega_0$ and a lattice $\omega^{\text{mod}} := (x')^*(\Omega^{\text{mod}}) \subset \omega_0 \otimes_{\mathbb{W}(\mathbb{F})} \mathcal{O}_K$, pull-back of the sheaf Ω^{mod} of Proposition 4.1, such that the Hodge–Tate map $T_p(A^\vee) \otimes \mathcal{O}_{\mathbb{C}_p} \rightarrow \omega_0 \otimes_{\mathbb{W}(\mathbb{F})} \mathcal{O}_{\mathbb{C}_p}$ factors via a surjective map

$$\text{HT}: T_p(A^\vee) \otimes \mathcal{O}_{\mathbb{C}_p} \longrightarrow \omega^{\text{mod}} \otimes_{\mathcal{O}_K} \mathcal{O}_{\mathbb{C}_p};$$

- (vi) a trivlization $q: H_0 \cong \Lambda \otimes \mathbb{W}(\mathbb{F})$ such that ω_0 is mapped isomorphically onto $\Lambda_0 \otimes_{\mathbb{W}(\kappa)} \mathbb{W}(\mathbb{F})$ and it sends the tensors $(s_{\text{cris},\alpha})_\alpha$ to the elements $(s_\alpha)_\alpha$;

Thanks to [16, Prop. A.2.2] we have a comparison morphism

$$\alpha_{\text{st}}: T_p(A^\vee) \otimes_{\mathbb{Z}_p} \widehat{A}_{\text{st}} \longrightarrow H_0 \otimes_{\mathbb{W}(\mathbb{F})} \widehat{A}_{\text{st}} \cdot \frac{1}{t}.$$

The ring \widehat{A}_{st} is a variant of Fontaine period rings introduced in [3]: it is the p -adic completion of the DP- A_{cris} -algebra $A_{\text{cris}}\langle X \rangle$, it is endowed with an action of the Galois group of K_0 , with an A_{cris} -linear derivation N and with a Frobenius φ , it admits a Galois equivariant map $\vartheta: \widehat{A}_{\text{st}} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ restricting to the classical map of Fontaine on A_{cris} and sending $X \mapsto 0$. The element $t \in A_{\text{cris}}$ is the classical period of Fontaine. The map α_{st} is a map of \widehat{A}_{st} -modules, compatible with all these extra structures. By [16, Prop 1.4.10, Cor. 3.3.9] it is an isomorphism after inverting t and it sends $(s_{\text{et},\alpha})_\alpha$ to $(s_{\text{cris},\alpha})_\alpha$.

Write $\widehat{B}_{\text{st}} := \widehat{A}_{\text{st}}[t^{-1}]$. There is a Galois equivariant map $\widehat{A}_{\text{st}} \rightarrow B_{\text{dR}}^+$ (see [3, Sect. 7]). Consider the base change $\alpha_{\text{dR}}^+(A)$ of $\alpha_{\text{st}}(A)$ to B_{dR}^+ . By loc. cit., it coincides with the classical de Rham comparison isomorphism of Fontaine after inverting t . In particular,

Corollary 5.1 *The reduction of α_{st} via $\widehat{A}_{st} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ is the map $d \log: T_p A^\vee \otimes_{\mathbb{Z}_p} \mathcal{O}_{\mathbb{C}_p} \rightarrow \Omega_{\widehat{A}_{st}} \otimes_{\mathbb{W}(\mathbb{F})} \mathcal{O}_{\mathbb{C}_p}$ of Proposition 4.1. Moreover, $H_0 \otimes_{\mathbb{W}(\mathbb{F})} \widehat{A}_{st}$ is contained in the image of α_{st} .*

Proof Consider the map $\alpha_{st}(A^\vee)$ for the dual abelian variety,

$$\alpha_{st}(A^\vee): T_p A \otimes_{\mathbb{Z}_p} \widehat{A}_{st} \longrightarrow H_0(\widetilde{A}_{\mathbb{F}}^\vee) \otimes_{\mathbb{W}(\mathbb{F})} \widehat{A}_{st} \cdot \frac{1}{t}.$$

Then $H_0(\widetilde{A}_{\mathbb{F}}^\vee) \cong H_0(\widetilde{A}_{\mathbb{F}})^\vee = H_0^\vee$ by [16, Prop. 1.3.5] and $(T_p A)^\vee \cdot t \cong T_p(A^\vee)$. Taking the \widehat{A}_{st} -dual of $\alpha_{st}(A^\vee)$ and using these identifications we get a map

$$\beta_{st}: H_0 \otimes_{\mathbb{W}(\mathbb{F})} \widehat{A}_{st} \longrightarrow T_p(A^\vee) \otimes_{\mathbb{Z}_p} \widehat{A}_{st}. \tag{5}$$

We claim that $\beta_{st} \circ \alpha_{st}$ is multiplication by t . This can be proved after inverting t . Notice that by [16, Prop. A.2.7] the base change of the maps α_{st} and β_{st} to \widehat{B}_{st} are already defined over the subring $B_{st} \subset \widehat{B}_{st}$. Then the fact that $\beta_{st} \circ \alpha_{st}$ is multiplication by t can be checked after base change to B_{dR} , as the induced map $B_{st} \rightarrow B_{dR}$ is injective. The base change of α_{st} and β_{st} to B_{dR} are Fontaine’s classical de Rham comparison isomorphisms. The claim follows then from the fact that the de Rham comparison isomorphisms are compatible with the Poincaré duality in étale and de Rham cohomology respectively. \square

5.2 A description of $\zeta'_1(x_0)$ and $\zeta'_2(x_0)$

Fix a basis \mathcal{C} of Λ as \mathbb{Z}_p -module. Define $g \in G_{\mathbb{Z}_p}(\mathbb{W}(\mathbb{F}))$ to be

$$g := q \circ \frac{\varphi}{p}|_{\omega_0} \oplus \varphi|_{q^{-1}(\Lambda_{-1} \otimes_{\mathbb{W}(k)} \mathbb{W}(\mathbb{F}))} \circ q^{-1}: \Lambda \otimes_{\mathbb{W}(k)} \mathbb{W}(\mathbb{F}) \longrightarrow \Lambda \otimes_{\mathbb{W}(k)} \mathbb{W}(\mathbb{F}).$$

Here q is the trivialization in Sect. 5.1(vi). Let $g_0 \in H(\mathbb{F})$ be the reduction of g modulo p ; recall that $H = (G_{\mathbb{Z}_p})_{\mathbb{F}_p}$. As explained in Sect. 2.3

$$\zeta'_1(x_0) = [g_0] \in E_{\mathcal{Z}_{EO}}(k) \setminus H(k).$$

Assume $\zeta'_1(x_0) \in H_{w\widetilde{z}^{-1}}$ for some $w \in {}^I W$ and with $z := w_0^K$, see Lemma 7.1 for the notation. Thanks to Proposition 3.8 there exists a lift x' in S^Σ such that, possibly after multiplying g by an element $\gamma = (a, b^{(p)})$ of $\widetilde{E}_{\mathcal{Z}_{EO}}(\mathbb{W}(k))$ with $a \in \widetilde{P}(\mathbb{W}(k))$ and $b \in \widetilde{P}_-(\mathbb{W}(k))$, we have $w\widetilde{z}^{-1} = ag(b^{(p)})^{-1}$.

We denote by \mathcal{F} the basis $q^{-1}(\mathcal{C})$ of H_0 . Write $a = u_+ \ell$ and $b^{(p)} = u_- \ell^{(p)}$ for their decomposition using $\widetilde{P} = R_u \widetilde{P} \cdot \widetilde{L}$ and $\widetilde{Q} = R_u \widetilde{Q} \cdot \widetilde{L}^{(p)}$. We let \mathcal{A} be the basis of H_0 such that the matrix of change of basis is ${}_{\mathcal{A}}M_{\mathcal{F}}(\text{Id}) = a^{-1}$. Thanks to Lemma 2.13 we have

$${}_{\mathcal{A}}M_{\mathcal{A}^{(p)}}(\varphi) = ag \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} (a^{(p)})^{-1} = ag(\ell^{(p)})^{-1} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} (u_+^{(p)})^{-1} = w\widetilde{z}^{-1} u_- \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} (u_+^{(p)})^{-1},$$

as $\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}$ commutes with $\tilde{L}^{(p)}$, and $w\tilde{z}^{-1} = ag(\ell^{(p)})^{-1}u_-^{-1}$. We remark that $r_- := u_- \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \in \tilde{z}\tilde{B}(\mathbb{W}(k)[p^{-1}])$ and that $u_+ \in R_u\tilde{P}(\mathbb{W}(k))$ as $\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \in \tilde{T}(\mathbb{W}(k)[p^{-1}])$ and $u_- \in R_u\tilde{P}_-^{(p)}(\mathbb{W}(k)) \subset \tilde{z}R_u\tilde{B}(\mathbb{W}(k))$. In particular, we have the following formula:

$$\mathcal{A}M_{\mathcal{A}^{(p)}}(\varphi) = w\tilde{z}^{-1} \cdot r_- \cdot u_+^{(p)}, \quad r_- \in \tilde{z}\tilde{B}(\widehat{A}_{\text{st}}[p^{-1}]), \quad u_+ \in R_u\tilde{P}(\widehat{A}_{\text{st}}). \tag{6}$$

On the other hand we have a surjective map

$$\text{HT} \circ d^{-1}: \Lambda \otimes \mathcal{O}_{\mathbb{C}_p}(1) \longrightarrow T_pA^\vee \otimes \mathcal{O}_{\mathbb{C}_p} \longrightarrow \omega^{\text{mod}} \otimes_{\mathcal{O}_K} \mathcal{O}_{\mathbb{C}_p}$$

defining the point $\pi_{\text{HT}}(x)$, where d is the trivialization in 5.1(iii). Take $h \in G_{\mathbb{Z}_p}(\mathcal{O}_{\mathbb{C}_p})$ to be an element such that $\text{Ker}(\text{HT} \circ d^{-1})$ is the filtration $h(\Lambda_{-1} \otimes_{\mathbb{W}(\mathbb{F})} \mathcal{O}_{\mathbb{C}_p})$. Then h_0 , which is h modulo p , represents $\pi_{\text{HT},0}(x_0)$ and

$$\zeta'_2(x_0) = [h_0(h_0^{(p)})^{-1}] \in E_{\mathcal{Z}_{\text{opp}}}(k) \setminus H(k).$$

We let $v \in {}^L W$ be such that $\zeta'_2(x_0)$ lies in the v -stratum as defined in Corollary 4.2 and the discussion that follows the Corollary, that is,

$$v = \tilde{y}^{-1}h_0(h_0^{(p)})^{-1}\tilde{z} \in E_{\mathcal{Z}_-}(k) \setminus H(k) \cong {}^L W.$$

Recall that $y := {}^L w_0 = w_0^L$ and, denoting by φ the automorphism of W defined by Frobenius on G_k , we have $\varphi(y) = z$.

Denote by $\mathcal{E} = d^{-1}(\mathcal{C})$ the basis of $T_p(A)^\vee$ given by pull back via d of the given basis \mathcal{C} of Λ . Lift $h \in G(\mathcal{O}_{\mathbb{C}_p})$ to an element $\tilde{h} \in G(\widehat{A}_{\text{st}})$ via the surjective map $\vartheta: \widehat{A}_{\text{st}} \rightarrow \mathcal{O}_{\mathbb{C}_p}$. Then $\tilde{h}(\tilde{h}^{(p)})^{-1} \in G(\widehat{A}_{\text{st}})$ lifts $h_0(h_0^{(p)})^{-1}$. Here $\tilde{h}^{(p)}$ is defined using the Frobenius φ on \widehat{A}_{st} . Let \mathcal{D} be the basis of $T_p(A)^\vee \otimes \widehat{A}_{\text{st}}$ defined by $\tilde{h}^{-1} \cdot d$. Then the matrix of change of basis expressing \mathcal{E} as a combination of \mathcal{D} is ${}_{\mathcal{D}}M_{\mathcal{E}}(\text{Id}) = \tilde{h}$. Let Frobenius φ on $T_p(A) \otimes \widehat{A}_{\text{st}}$ be $1 \otimes \varphi$. Its matrix with respect to the basis \mathcal{E} is the identity so that

$${}_{\mathcal{D}}M_{\mathcal{D}^{(p)}}(\varphi) = {}_{\mathcal{D}}M_{\mathcal{E}}(\text{Id})_{\mathcal{E}}M_{\mathcal{E}^{(p)}}(\varphi)_{\mathcal{E}^{(p)}}M_{\mathcal{D}^{(p)}}(\text{Id}) = \tilde{h}(\tilde{h}^{(p)})^{-1}. \tag{7}$$

5.3 Reduction steps

We let $\tau \in G(\widehat{B}_{\text{st}})$ be the element $\tau = \mathcal{A}M_{\mathcal{D}}(\alpha_{\text{st}})$. It is the matrix of

$$\begin{aligned} ((aq) \otimes 1) \circ \alpha_{\text{st}} \circ ((\tilde{h}^{-1}d) \otimes 1)^{-1}: \Lambda \otimes \widehat{A}_{\text{st}} &\longrightarrow T_p(A^\vee) \otimes_{\mathbb{Z}_p} \widehat{A}_{\text{st}} \\ &\longrightarrow H_0 \otimes_{\mathbb{W}(\mathbb{F})} \widehat{A}_{\text{st}} \longrightarrow \Lambda \otimes \otimes_{\mathbb{Z}_p} \widehat{A}_{\text{st}} \end{aligned}$$

with respect to the given basis \mathcal{C} of Λ . Then,

Proposition 5.2 *Assume that $p > 3$. There exists a \widehat{A}_{st} -algebra R^+ with the following properties:*

- (i) R^+ is t -torsion free and local with maximal ideal \mathfrak{m} and residue field k and the quotient map $j: R^+ \rightarrow k$ is compatible with the map $\widehat{A}_{st} \rightarrow k$ given by composing $\vartheta_{st}: \widehat{A}_{st} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ and the residue field map $\mathcal{O}_{\mathbb{C}_p} \rightarrow k$. We write $R := R^+[t^{-1}]$;
- (ii) τ and $\tau^{(p)}$, viewed in $G(R)$, admit decompositions $\tau = \beta \cdot \ell$ and $\tau^{(p)} = \beta' \cdot \ell'$ such that $\beta \in \widetilde{B}(R)$ and $\beta' \in \widetilde{B}(R)$ and ℓ and ℓ' are in $G(R^+)$ and their reduction $\bar{\ell}$ and $\bar{\ell}'$ modulo \mathfrak{m} lie in $L(k)$ and satisfy $\bar{\ell}' = \bar{\ell}^{(p)}$.
- (iii) we have a decomposition as in (ii) with the same $\bar{\ell}$ of (ii), if we replace τ and $\tau^{(p)}$ with $s \cdot \tau$, resp. $(s \cdot \tau)^{(p)}$, for some $s \in \mathbf{R}_u \widehat{P}(\widehat{A}_{st})$.

Proof We write τ as

$$\tau = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{End}(\Lambda) \otimes \widehat{A}_{st}$$

with $a \in \text{End}(\Lambda_0) \otimes_{\mathbb{W}(\kappa)} \widehat{A}_{st}$, $b \in \text{Hom}(\Lambda_1, \Lambda_0) \otimes_{\mathbb{W}(\kappa)} \widehat{A}_{st}$ etc.

Due to Corollary 5.1 we have that b, c, d are 0 modulo $\text{Ker}(\vartheta_{st})$ and if we write $\delta = \begin{pmatrix} m & n \\ u & v \end{pmatrix} \in \text{End}(\Lambda \otimes \widehat{A}_{st})$ for the matrix ${}_{\mathcal{D}}M_{\mathcal{A}}(\beta_{st})$ of the morphism β_{st} of (5) we have

$$\delta \cdot \tau = \tau \cdot \delta = t \mathbf{1}_{\Lambda}.$$

Then also m, n and u lie in $\text{Ker}(\vartheta_{st})$.

Consider the ring $A_{0,p-1,\omega}^{st}$ defined in Sect. 6. It is a \widehat{A}_{st} -algebra, the map ϑ_{st} extends to a map $\vartheta_{0,p-1,\omega}: A_{0,p-1,\omega} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ and $\text{Ker}(\vartheta_{st}) \subset \omega A_{0,p-1,\omega}^{st}$. The element $t \in A_{\text{cris}}$ is equal to $[\varepsilon] - 1$ times a unit of A_{cris} (see for example [4, Lemme 6.2.13]). As $[\varepsilon] - 1 = ([\varepsilon^{\frac{1}{p}}] - 1)\omega$ times a unit of A_{inf} we conclude that $t = ([\varepsilon^{\frac{1}{p}}] - 1)\omega g$ with g a unit of A_{cris} . Hence

$$\tau' := \widetilde{\mu}(\omega)^{-1} \tau = \begin{pmatrix} a & b \\ c\omega^{-1} & d\omega^{-1} \end{pmatrix}, \quad \delta' := \delta \cdot (\widetilde{\mu}(\omega)\omega^{-1}) = \begin{pmatrix} m\omega^{-1} & n\omega^{-1} \\ u & v \end{pmatrix}$$

are both elements of $\text{End}(\Lambda) \otimes A_{0,p-1,\omega}^{st}$ and

$$\delta' \cdot \tau' \equiv [p^{\frac{1}{p-1}}] \mathbf{1}_{2g}$$

modulo $\text{Ker}(\vartheta_{0,p-1,\omega}^{st})$.

As $\vartheta_{0,p-1,\omega}(b) = 0$, the reduction $\bar{\tau}'$ of τ' modulo $\text{Ker}(\vartheta_{0,p-1,\omega})$ lies in $P_-(\mathbb{C}_p)$. Thanks to Lemma 6.3 we can find $\tau_0 \in \widetilde{L}(A_{\text{inf}})$ and $\tau_1 \in \widetilde{B}(A_{\text{inf}}[[p]^{-1}]) \cap \text{End}(\Lambda) \otimes A_{\text{inf}}$ such that $\tau' \equiv \tau_1 \tau_0$ in $\text{End}(\Lambda) \otimes \mathcal{O}_{\mathbb{C}_p}$ (via the map $\vartheta_{0,p-1,\omega}^{st}$). We define δ_0 and

δ_1 similarly. Then

$$(\delta_1 \delta_0) \cdot (\tau_1 \tau_0) \equiv [\underline{p}^{\frac{1}{p-1}}] \mathbf{1}_{2g}$$

modulo $\text{Ker}(\vartheta)$. We now pass to $A_{r,p-1,\omega}^{\text{st}}$, for $1 > r > \frac{1}{p-1}$. It is a $A_{0,p-1,\omega}^{\text{st}}$ -algebra and $\vartheta_{0,p-1,\omega}$ extends to the morphism $\vartheta_{r,p-1,\omega}$ by Lemma 6.1. Corollary 6.2 implies that

$$(\delta_1 \delta_0) \cdot (\tau_1 \tau_0) = [\underline{p}^{\frac{1}{p-1}}] D \tag{8}$$

with $D \in G(A_{r,p-1,\omega}^{\text{st}})$.

Let us now work with $A_{r,s,[\underline{p}]^r \omega}^{\text{st}}$ with $s = p - 1 - r$. Using Lemma 6.1 again, we know that it is a $A_{r,p-1,\omega}^{\text{st}}$ -algebra and $\vartheta_{r,p-1,\omega}$ extends to the morphism $\vartheta_{r,s,[\underline{p}]^r \omega}$. Moreover, $\text{Ker}(\vartheta_{0,p-1,\omega}) \subset [\underline{p}]^r A_{r,s,\omega}^{\text{st}}$ by Lemma 6.1. Hence, $\tau' \equiv \tau_1 \tau_0$ modulo $[\underline{p}]^r A_{r,s,[\underline{p}]^r \omega}^{\text{st}}$. Consider the matrix

$$\tau'' := (\tau_1 \tau_0)^{-1} \tau' = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} \in G_{\mathbb{Z}_p}(A_{r,s,[\underline{p}]^r \omega}^{\text{st}}[\underline{p}]^{-1}) \cap \text{End}(\Lambda) \otimes A_{r,p-1,\omega}^{\text{st}}[[\underline{p}]^{-1}].$$

Due to (8) we have that $(\tau_1 \tau_0)^{-1}$ is in $\text{End}(\Lambda) \otimes \left([\underline{p}^{\frac{-1}{p-1}}] \cdot A_{r,s,[\underline{p}]^r \omega}^{\text{st}}\right)$ so that $[\underline{p}^{\frac{1}{p-1}}] \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$ is in $\text{End}(\Lambda) \otimes A_{r,s,[\underline{p}]^r \omega}^{\text{st}}$ and is congruent to $[\underline{p}^{\frac{1}{p-1}}] \mathbf{1}_{2g}$ modulo $[\underline{p}]^r A_{r,s,[\underline{p}]^r \omega}^{\text{st}}$. Hence τ'' is a matrix congruent to $\mathbf{1}_{2g}$ modulo $[\underline{p}]^{r-\frac{1}{p-1}}$ and, hence, it is invertible as $A_{r,s,[\underline{p}]^r \omega}^{\text{st}}$ is $[\underline{p}]$ -adically complete and separated by Lemma 6.1, that is $\tau'' \in G(A_{r,s,[\underline{p}]^r \omega}^{\text{st}})$.

In conclusion, we have $\tau = \beta \ell$ with $\beta := \tilde{\mu}(\omega) \tau_1 \in \tilde{\gamma} \tilde{B}_-(A_{r,s,[\underline{p}]^r \omega}^{\text{st}}[[\underline{p}]^{-1} \cdot \omega^{-1}])$ and $\ell := \tau_0 \cdot \tau'' \in G_{\mathbb{Z}_p}(A_{r,s,[\underline{p}]^r \omega}^{\text{st}})$ such that $\bar{\ell} \equiv \bar{\tau}_0 \in L(k)$; here $\bar{\ell}$, resp. $\bar{\tau}_0$, is the image of ℓ , resp. τ_0 in $G_{\mathbb{Z}_p}(k) = H(k)$ via the map $j_{r,s} : A_{r,s,[\underline{p}]^r \omega}^{\text{st}} \rightarrow k$ defined by $\vartheta_{r,s,[\underline{p}]^r \omega}$ composed with the quotient $\mathcal{O}_{\mathbb{C}_p} \rightarrow k$.

Frobenius φ on \widehat{A}_{st} extends to a map $\varphi : A_{r,s,[\underline{p}]^r \omega}^{\text{st}} \rightarrow A_{pr,s,[\underline{p}]^{pr} \varphi(\omega)}^{\text{st}}$ (see Lemma 6.1). The latter is well defined, together with the map $j_{pr,s} : A_{pr,s,[\underline{p}]^{pr} \varphi(\omega)}^{\text{st}} \rightarrow k$, if and only if $pr < s = (p - 1) - r$, i.e., if there exists $r > \frac{1}{p-1}$ such that $(p + 1)r < p - 1$ which is equivalent to require that $p > 3$ and explains the assumption. Then

$$\tau^{(p)} = \beta' \ell', \quad \beta' = \varphi(\beta) \in \tilde{z} \tilde{B}(A_{pr,s,[\underline{p}]^{pr} \varphi(\omega)}^{\text{st}}[[\underline{p}]^{-1} \cdot \varphi(\omega)^{-1}]), \quad \ell' = \varphi(\ell) \in G_{\mathbb{Z}_p}(A_{r,s,[\underline{p}]^r \omega}^{\text{st}}).$$

By construction $\bar{\ell}' = \bar{\ell}^{(p)} \in L^{(p)}(k)$. Here $\varphi(\beta)$ and $\varphi(\ell)$ are defined by applying φ to the entries of the matrices β and ℓ . They lie in $\tilde{z} \tilde{B}$ as $\varphi(y) = z$ and in $G_{\mathbb{Z}_p}$ respectively as they are both subgroups of $\text{GSp}(\Lambda)$ defined over \mathbb{Z}_p .

Finally $\omega = [\underline{p}]([\underline{p}]^{pr-1} \frac{p}{[\underline{p}]^{pr}} - 1)$ is equal to $[\underline{p}]$ times a unit in $A_{pr,s,[\underline{p}]^{pr}\varphi(\omega)}^{\text{st}}$. Analogously $\varphi(\omega) = p(1 - \frac{[\underline{p}]^s}{p} [\underline{p}]^{p-s})$ is p times a unit in $A_{r,s,[\underline{p}]^r\omega}^{\text{st}}$. Hence both $A_{r,s,[\underline{p}]^r\omega}^{\text{st}}$ and $A_{pr,s,[\underline{p}]^{pr}\varphi(\omega)}^{\text{st}}$ map to $A_{pr,s,[\underline{p}]^{pr}p}^{\text{st}}$. We let R^+ be the localization of $A_{pr,s,[\underline{p}]^{pr}p}^{\text{st}}$ with respect to the maximal ideal defined by the kernel of the map $j_{pr,s}: A_{pr,s,[\underline{p}]^{pr}p}^{\text{st}} \rightarrow k$. As $t = ([\varepsilon^{\frac{1}{p}}] - 1)\omega u$ with u a unit in R^+ and $\omega = [\underline{p}]$ times a unit in R^+ , inverting t or inverting $[\underline{p}]$ in R^+ gives the same ring R . As $p = [\underline{p}]^{p-1} = p \cdot \frac{[\underline{p}]^{p-1}}{p}$ then p is invertible in R as well. As R^+ is p -torsion free and $p \in [\underline{p}]^{pr} A_{pr,s,[\underline{p}]^{pr}p}^{\text{st}}$, then R^+ is also $[\underline{p}]$ -torsion free.

For (iii) the same argument for $s \cdot \tau$ in place of τ holds. We only need to prove that one obtains a decomposition with the same $\bar{\ell}$. Define

$$\tau''' := \tilde{\mu}(\omega)^{-1} s \cdot \tau = (\tilde{\mu}(\omega)^{-1} s \tilde{\mu}(\omega)) (\tilde{\mu}(\omega)^{-1} \tau).$$

Then $s' = \tilde{\mu}(\omega)^{-1} s \tilde{\mu}(\omega)$ is congruent to $\mathbf{1}_{2g}$ modulo ω as $s \in R_u \tilde{P}(\widehat{A}_{\text{st}})$. Hence, $\tau''' \equiv \tau_1 \tau_0$ in $\text{End}(\Lambda) \otimes \mathcal{O}_{\mathbb{C}_p}$. Then the argument continues as before and we get a decomposition $s \cdot \tau = \beta'' \ell''$ with $\beta'' \in \tilde{y} \tilde{B}(R)$ and $\ell'' \in G(R^+)$ and $\ell'' \equiv \bar{\tau}_0 \equiv \ell_0$ modulo \mathfrak{m} as wanted. \square

Remark 5.3 If one is interested in a decomposition of τ as in Proposition 5.2, one can work with B_{dR} and the proof is much easier. It is the need to decompose $\tau^{(p)}$ as well that forced us to look into the finer structure of \widehat{A}_{st} . As we need the existence of a positive rational number r such that $r > \frac{1}{p-1}$ and $(p+1)r < p-1$ the hypothesis $p \geq 5$ is forced on us.

5.4 Conclusion

Recall that $\tau = {}_{\mathcal{A}}M_{\mathcal{D}}(\alpha_{\text{st}})$. It follows from (6) and (7) that we have the following equality in $G(R)$:

$$\tau \tilde{h}(\tilde{h}^{(p)})^{-1} ((u_+ \cdot \tau)^{(p)})^{-1} = w \tilde{z}^{-1} r_-.$$

Decompose $\tau = \beta \ell$ and $u_+ \cdot \tau = \rho q$ as in Proposition 5.2 with ℓ and $q \in G_{\mathbb{Z}_p}(R^+)$ such that $\bar{\ell} = \bar{q} =: \ell_0 \in L(k)$ modulo \mathfrak{m} and β and $\rho \in {}^y \tilde{B}(R)$. Consider the class

$$\gamma := \ell \tilde{h}(\tilde{h}^{(p)})^{-1} (q^{(p)})^{-1} \in \tilde{y} \tilde{B}(R^+) \backslash G(R^+) / \tilde{z} \tilde{B}(R^+).$$

Its image γ_0 via $j: R^+ \rightarrow k$ is $\gamma_0 = \ell_0 h_0 (h_0^{(p)})^{-1} (\ell_0^{(p)})^{-1} \in \tilde{y} B(k) \backslash G(k) / \tilde{z} B(k)$. We also have

$$\gamma \equiv \beta \gamma (\rho^{(p)})^{-1} = \tau \tilde{h}(\tilde{h}^{(p)})^{-1} ((u_+ \cdot \tau)^{(p)})^{-1} = w \tilde{z}^{-1} r_- = w \tilde{z}^{-1}$$

in $\tilde{y}\tilde{B}(R)\backslash G(R)/\tilde{z}\tilde{B}(R)$ as $r_- \in \tilde{z}\tilde{B}(R)$ by (6). In conclusion,

$$\gamma_0 = \ell_0 h_0 (h_0^{(p)})^{-1} (\ell_0^{(p)})^{-1} \leq_- w \tilde{z}^{-1}$$

for the Bruhat order \leq_- on $G/\tilde{z}\tilde{B}$ (with respect to the left action of $\tilde{y}\tilde{B}$).

Recall that we have chosen h_0 so that $\tilde{y}^{-1}h_0(h_0^{(p)})^{-1}\tilde{z} = v \in {}^L W$. It is the same class as $\tilde{y}^{-1}\gamma_0\tilde{z}$ for the order relation \preceq on ${}^L W$. Consider the isomorphism

$$\delta: G/\tilde{z}\tilde{B} \rightarrow G/\tilde{B}, [g] \mapsto [\tilde{y}^{-1}g\tilde{z}];$$

it is equivariant with respect to the left action of $\tilde{y}\tilde{B}$ on the left hand side, resp. \tilde{B} on the right hand side. In particular, it is order preserving considering on G/\tilde{B} the standard Bruhat order. Hence,

$$\delta(\ell_0 h_0 (h_0^{(p)})^{-1} (\ell_0^{(p)})^{-1}) \leq \tilde{y}^{-1}w$$

for the standard Bruhat order. Recall that the class of \tilde{y} in the Weyl group W is ${}^L w_0$ so that the class of \tilde{y}^{-1} is w_0^L . Thanks to Lemma 7.2, the map

$$\sigma_0: {}^L W \longrightarrow {}^L W, \quad w \mapsto y^{-1}w$$

is an order reversing bijection and we have

$$v = \delta(\ell_0 h_0 (h_0^{(p)})^{-1} (\ell_0^{(p)})^{-1}) \preceq \sigma_0(w)$$

in ${}^L W$ as wanted.

6 Appendix I: Fontaine’s rings

Given the tilt \mathbb{C}_p^b of \mathbb{C}_p and the tilt $\mathcal{O}_{\mathbb{C}_p}^b$ of $\mathcal{O}_{\mathbb{C}_p}$, we have Fontaine’s classical ring A_{cris} . It is the p -adic completion of the DP envelope of $A_{\text{inf}} = \mathbb{W}(\mathcal{O}_{\mathbb{C}_p}^b)$ with respect to the kernel of the surjective map $\vartheta: A_{\text{inf}} \rightarrow \mathcal{O}_{\mathbb{C}_p}$. The ideal $\text{Ker}(\vartheta)$ is principal. Generators are, for example, $([\epsilon]-1)/([\epsilon^{1/p}]-1)$ or $p-[p]$ where $\epsilon = (1, \zeta_p, \zeta_{p^2}, \dots) \in \mathcal{O}_{\mathbb{C}_p}^b$ is a compatible sequence of primitive p^n -th roots of unity and $\underline{p} = (p, p^{1/p}, p^{1/p^2}, \dots) \in \mathcal{O}_{\mathbb{C}_p}^b$ is a compatible sequence of p^n -th roots of p . Let ω be a generator of $\text{Ker}(\vartheta)$. We also have the ring $\widehat{A}_{\text{st}} := A_{\text{cris}}\langle X \rangle$, introduced by Breuil [3], given by the p -adic completion of the DP polynomial ring $A_{\text{cris}}[X]$. The map ϑ extends to a map $\vartheta_{\text{st}}: \widehat{A}_{\text{st}} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ sending $X \mapsto 0$. It is also endowed with a Frobenius, extending the classical one on A_{cris} , requiring that $\varphi(X) = (1 + X)^p - 1$.

Given elements r and $s \in \mathbb{Q}$ with $0 \leq r \leq s$ let

$$A_{r,s} = A_{\text{inf}} \left\{ \frac{p}{[p]^r}, \frac{[p]^s}{p} \right\},$$

be the p -adic completion of the ring $A_{\text{inf}}[W, Z]/([p]^r W - p, pZ - [p]^s, WZ - [p]^{s-r})$. It coincides with the p -adic completion of the subring $A_{\text{inf}} \left[\frac{p}{[p]^r}, \frac{[p]^s}{p} \right]$ of $\text{Frac}(A_{\text{inf}})$ by [2, Cor. 2.2]. For $\gamma \in A_{r,s}$ a non-zero element we define

$$A_{r,s,\gamma}^{\text{st}} = A_{\text{inf}} \left\{ \frac{p}{[p]^r}, \frac{[p]^s}{p}, \frac{X}{p\gamma} \right\}$$

to be the p -adic completion of the ring

$$A_{\text{inf}}[W, Z, X, Y]/([p]^r W - p, pZ - [p]^s, WZ - [p]^{s-r}, p\gamma Y - X).$$

For $r < 1 < s$ the map ϑ_{st} extends to a map

$$\vartheta_{r,s,\gamma}: A_{r,s,\gamma}^{\text{st}} \longrightarrow \mathcal{O}_{\mathbb{C}_p}, \quad W \mapsto p^{1-r}, Z \mapsto p^{s-1}, Y \mapsto 0.$$

For $r < s$ we also have the map

$$j_{r,s}: A_{r,s,\gamma}^{\text{st}} \longrightarrow k,$$

that on A_{inf} is the composite of ϑ with the quotient map $\mathcal{O}_{\mathbb{C}_p} \rightarrow k$ on the residue field and that sends W, Z, X and Y to 0. It coincides with the composite of the quotient map $\mathcal{O}_{\mathbb{C}_p} \rightarrow k$ with $\vartheta_{r,s,\gamma}$ whenever the latter exists.

Lemma 6.1 *Assume that $p \geq 3$. Then*

- (1) *For every $s \leq p - 1$ the map $A_{\text{inf}} \rightarrow A_{r,s,\gamma}^{\text{st}}$ extends to a map $\widehat{A}_{\text{st}} \rightarrow A_{r,s,\gamma}^{\text{st}}$. We have $\text{Ker}(\vartheta_{\text{st}}) \subset [p]^{\min(r,1)} A_{r,s,\gamma}^{\text{st}}$ and $\text{Ker}(\vartheta_{\text{st}}) \subset \omega A_{r,s,\gamma}^{\text{st}}$ if $\gamma \in \omega A_{r,s}$. Moreover, the ring $A_{r,s,\gamma}^{\text{st}}$ is p -torsion free and p -adically, and hence $[p]$ -adically, complete and separated.*
- (2) *For $r < 1 < s$ the kernel of $\vartheta_{r,s,\gamma}$ is generated by $(\omega, W - [p]^{1-r}, Z - [p]^{s-1}, Y)$.*
- (3) *For $r < r' < 1 < s' < s$ and $\alpha \in A_{r',s'}$ a non-zero element, there are unique morphisms of A_{inf} -algebras $A_{r,s} \rightarrow A_{r',s'}$, resp. of \widehat{A}_{st} -algebras $A_{r,s,\gamma}^{\text{st}} \rightarrow A_{r',s',\alpha\gamma}^{\text{st}}$ sending $W \mapsto [p]^{r'-r} W, Z \mapsto [p]^{s-s'} Z, Y \mapsto \alpha Y$.*
- (4) *For $pr < s$ Frobenius φ on \widehat{A}_{st} extends uniquely to a ring homomorphism $\varphi: A_{r,s,\gamma}^{\text{st}} \rightarrow A_{pr,s,\varphi(\gamma)}^{\text{st}}$ such that $\varphi(W) = W, \varphi(Z) = [p]^{(p-1)s} Z, \varphi(Y) = Y \cdot \frac{(1+X)^p - 1}{X}$.*

Proof (1) The ring $A_{r,s}$ is proven to be p -torsion free and p -adically complete and separated in [2]. As $A_{r,s,\gamma}^{\text{st}}$ is the p -adic completion of $A_{r,s}[X, Y]/(p\gamma Y - X) \cong A_{r,s}[Y]$, also $A_{r,s,\gamma}^{\text{st}}$ is p -torsion free and p -adically complete and separated.

By definition of \widehat{A}_{st} to prove the first claim it suffices to show that in $A_{r,s,\gamma}^{st}$ the ideal generated by ω and X admits divided powers for $\omega = p - [p]$. As $A_{r,s,\gamma}^{st}$ is p -torsion free, it suffices to prove that $\omega^{p^n} \in p^n! \omega A_{r,s}$ and $X^{p^n} \in p^n! \gamma A_{r,s}^{st}$. The p -adic valuation of $p^n!$ is $\frac{p^n-1}{p-1}$. As $p^n - \frac{p^n-1}{p-1} \geq p^{n-1}$ for $p \geq 3$ the second claim is clear. Next we show that $\omega^{p^n} \in p^{\frac{p^n-1}{p-1}} \omega A_{r,s}$. First $\omega^{p-1} \in (p, [p])^{p-1} A_{inf} \subset (p, [p]^{p-1}) A_{inf} \subset p A_{r,s}$ as $\frac{[p]^{p-1}}{p} \in A_{r,s}$ by assumption. Hence $\omega^p \in p \omega A_{r,s}$. Assuming that $\omega^{p^n} \in p^{\frac{p^n-1}{p-1}} \omega A_{r,s}$ one deduces that $\omega^{p^{n+1}} \in p^{\frac{p^{n+1}-p}{p-1}} \omega^p A_{r,s} \subset p^{\frac{p^{n+1}-p}{p-1}} p \omega A_{r,s}$. As $p^{\frac{p^{n+1}-p}{p-1}+1} = p^{\frac{p^{n+1}-1}{p-1}}$ we conclude that $\omega^{p^{n+1}} \in p^{\frac{p^{n+1}-1}{p-1}} \omega A_{r,s}$. This then holds for every n by induction and implies the claim.

For the second statement notice that the ideal $\text{Ker}(\vartheta_{st})$ is generated by the DP powers of ω and X . We have shown that the divided powers of X lie in $p A_{r,s,\gamma}^{st}$ and that the divided powers of ω lie in $\omega A_{r,s} \subset (p, [p]) A_{r,s}$. As $(p, [p]) A_{r,s} \subset ([p]^r, [p]) A_{r,s}$ the claim follows.

The last statement follows remarking that $[p]^s = \frac{[p]^s}{p} \cdot p$ in $A_{r,s,\gamma}^{st}$ and the latter is p -adically complete and separated.

(2), (3) and (4) are direct computations left to the reader using the explicit description of the rings involved. □

Corollary 6.2 *Take $s \geq p - 1$. Let M be an $n \times n$ matrix with coefficients in \widehat{A}_{st} .*

- (1) *Let \overline{M} be the reduction of M modulo $\text{Ker}(\vartheta_{st})$. Assume that there exist r and $q \in \mathbb{Q}$ with $0 \leq q < \min(r, 1)$ and $r \leq s$ and an $n \times n$ matrix \overline{N} with coefficients in $\mathcal{O}_{\mathbb{C}_p}$ such that $\overline{NM} = p^q \mathbf{1}_n$. Then there exists an $n \times n$ matrix N with coefficients in $\widehat{A}_{r,s,\gamma}^{st}$ such that $MN = NM = [p]^q \mathbf{1}_n$.*
- (2) *Let \overline{M} be the reduction of M modulo p . Assume that there exist q and $r \in \mathbb{N}[1/p]$ with $0 \leq q < r \leq s$ and an $n \times n$ matrix \overline{N} with coefficients in $\widehat{A}_{st}/p\widehat{A}_{st}$ such that $\overline{NM} = [p]^q \mathbf{1}_n$. Then there exists an $n \times n$ matrix N with coefficients in $\widehat{A}_{r,s,\gamma}^{st}$ such that $MN = NM = [p]^q \mathbf{1}_n$.*

Proof (1) Lift \overline{N} to a matrix N' with coefficients in \widehat{A}_{st} . Then $N'M = [p]^q \mathbf{1}_n + A$ with A a matrix with coefficients in $\text{Ker}(\vartheta_{st})$. Then A is a matrix with coefficients in $[p]^{\min(r,1)} A_{r,s,\gamma}^{st}$ by Lemma 6.1. Hence, $[p]^q \mathbf{1}_n + A = [p]^q B$ with B a matrix with coefficients in $A_{r,s,\gamma}^{st}$ congruent to the identity modulo $[p]^{\min(r,1)-q}$. Thus B is invertible as its determinant is a unit by Lemma 6.1. Take $N := B^{-1}N'$.

- (2) As before lift \overline{N} to a matrix N' with coefficients in \widehat{A}_{st} . Then $N'M = [p]^q \mathbf{1}_n + A$ with A with coefficients in $p\widehat{A}_{st}$ and, hence, with coefficients in $[p]^r A_{r,s,\gamma}^{st}$. Hence, $[p]^q \mathbf{1}_n + A = [p]^q B$ with B a matrix with coefficients in $A_{r,s,\gamma}^{st}$ congruent to the identity modulo $[p]^{r-q}$. Thus B is invertible and $N := B^{-1}N'$ is the sought for matrix. □

6.1 An approximation result

The map $\vartheta : A_{\text{inf}} \rightarrow \mathcal{O}_{\mathbb{C}_p}$ extends to a map $\vartheta : A_{\text{inf}}[[\underline{p}]^{-1}] \rightarrow \mathbb{C}_p$, sending $[\underline{p}] \mapsto p$, which is surjective. Following the conventions in the paper we fix $G_{\mathbb{Z}_p} \subset \text{GSp}_{2g, \mathbb{Z}_p}$ a reductive subgroup and a parabolic subgroup $\tilde{P} \subset G_{\mathcal{O}_{\mathbb{C}_p}}$ containing a Borel subgroup $\tilde{B} \subset \tilde{P}$ defined over \mathbb{Z}_p . We let $\tilde{L} \subset \tilde{P}$ be the induced standard Levi subgroup. We write B, L, P for the induced subgroups of $G_{\mathbb{C}_p}$.

Lemma 6.3 *Let $h \in G_{\mathbb{Z}_p}(A_{\text{inf}})$ and let $\bar{A} \in {}^h\tilde{P}(\mathbb{C}_p)$ be an element such that, viewed as a matrix in $\text{GSp}_{2g}(\mathbb{C}_p)$, it has coefficients in $M_{2g \times 2g}(\mathcal{O}_{\mathbb{C}_p})$. Then, there exist matrices $a \in {}^h\tilde{L}(A_{\text{inf}})$ and $A' \in {}^h\tilde{B}(A_{\text{inf}}[[\underline{p}]^{-1}])$ such that A' , viewed as matrix in $\text{GSp}_{2g}(A_{\text{inf}}[[\underline{p}]^{-1}])$, has coefficients in $M_{2g \times 2g}(A_{\text{inf}})$ and $(A' \cdot a) \equiv \bar{A}$ modulo $\text{Ker}(\vartheta)$.*

Proof We prove the statement for $h = \mathbf{1}$ and $\bar{A} \in \tilde{P}(\mathbb{C}_p)$. The other statement is proven from this by conjugating by h .

Decompose $\bar{A} = u \cdot \ell$ with $u \in R_u P(\mathbb{C}_p)$ (the unipotent radical of P) and $\ell \in L(\mathbb{C}_p)$. Let $\tilde{B}_L := \tilde{B} \cap \tilde{L}$ be the Borel subgroup of \tilde{L} defined by \tilde{B} . By the properness of the Grassmanian variety $\tilde{B}_L \backslash \tilde{L}$ we have that

$$\tilde{B}_L(\mathbb{C}_p) \backslash \tilde{L}(\mathbb{C}_p) = (\tilde{B}_L \backslash \tilde{L})(\mathbb{C}_p) = (\tilde{B}_L \backslash \tilde{L})(\mathcal{O}_{\mathbb{C}_p}) = \tilde{B}_L(\mathcal{O}_{\mathbb{C}_p}) \backslash \tilde{L}(\mathcal{O}_{\mathbb{C}_p}).$$

Hence we can write $\ell \in L(\mathbb{C}_p) = \tilde{L}(\mathbb{C}_p)$ as $u' \cdot \bar{a}$ with $u' \in \tilde{B}_L(\mathbb{C}_p)$ and $\bar{a} \in \tilde{L}(\mathcal{O}_{\mathbb{C}_p})$. As the map $\tilde{L}(A_{\text{inf}}) \rightarrow \tilde{L}(\mathcal{O}_{\mathbb{C}_p})$ is surjective by the smoothness of \tilde{L} we can lift \bar{a} to an element $a \in \tilde{L}(A_{\text{inf}})$.

Define $\bar{A}' := uu' \in B(\mathbb{C}_p)$. We are left to show that we can lift \bar{A}' to an element $A' \in \tilde{B}(A_{\text{inf}}[[\underline{p}]^{-1}])$ which has coefficients in A_{inf} when viewed in $M_{2g \times 2g}(A_{\text{inf}}[[\underline{p}]^{-1}])$. As \tilde{B} over $\mathcal{O}_{\mathbb{C}_p}$ is the semidirect product of a split torus \tilde{T} and a unipotent group \tilde{U} , and the latter is a split extension of \mathbb{G}_a 's, we are left to prove the statement for $\mathbb{G} := \mathbb{G}_m$ and $\mathbb{G} := \mathbb{G}_a$. Given a non-zero element $s \in \mathbb{G}(\mathbb{C}_p) \subset \mathbb{C}_p$, let α be its p -adic valuation. Then $s = p^\alpha s_0$ with $s_0 \in \mathbb{G}(\mathcal{O}_{\mathbb{C}_p})$. As $\mathbb{G}(A_{\text{inf}}) \rightarrow \mathbb{G}(\mathcal{O}_{\mathbb{C}_p})$ is surjective, we can lift s_0 to an element $\tilde{s}_0 \in \mathbb{G}(A_{\text{inf}})$ and then $\tilde{s} := [\underline{p}^\alpha]\tilde{s}_0 \in \mathbb{G}(A_{\text{inf}}[[\underline{p}]^{-1}])$ lifts s as wanted. We need to show that the lift can be taken in $M_{2g \times 2g}(A_{\text{inf}})$. Notice that this can be proven after conjugating by an element of $\text{GL}_{2g}(\mathcal{O}_{\mathbb{C}_p})$.

We claim that, after possibly conjugating by an element of $\text{GL}_{2g}(\mathcal{O}_{\mathbb{C}_p})$, we may assume that $\tilde{B} \subset \tilde{B}^{\text{std}}$, with $\tilde{B}^{\text{std}} \subset \text{GL}_{2g, \mathbb{Z}_p}$ the standard Borel subgroup of upper triangular matrices, and that \tilde{T} is contained in the standard torus of \tilde{B}^{std} . Indeed, extend \tilde{B} to a Borel subgroup \tilde{B}' of $\text{GL}_{2g, \mathcal{O}_{\mathbb{C}_p}}$: first extend it over $\overline{\mathbb{Q}}_p$ by extending the connected solvable subgroup $\tilde{B}_{\mathbb{Q}_p}$ to a maximal connected solvable subgroup $\tilde{B}'_{\overline{\mathbb{Q}}_p}$ of $\text{GL}_{2g, \overline{\mathbb{Q}}_p}$. Since the scheme of Borel subgroups $\tilde{B}^{\text{std}}_{\mathbb{Z}_p} \backslash \text{GL}_{2g, \mathbb{Z}_p}$ is proper, the $\overline{\mathbb{Q}}_p$ -valued point defined by $\tilde{B}'_{\overline{\mathbb{Q}}_p}$ extends to a $\overline{\mathbb{Z}}_p$ -valued point so that, after possibly conjugation by an element of $\text{GL}_{2g}(\overline{\mathbb{Z}}_p)$, we may assume that $\tilde{B}_{\mathbb{Q}_p} \subset \tilde{B}^{\text{std}}_{\mathbb{Q}_p}$ and, hence, $\tilde{B} \subset \tilde{B}^{\text{std}}$.

As maximal tori in \tilde{B}^{std} are conjugate over $\mathcal{O}_{\mathbb{C}_p}$ we may also assume that \tilde{T} is contained in the standard torus of \tilde{B}^{std} of diagonal matrices.

By our assumption we now have that \overline{A}' lies in the upper triangular matrices in $M_{2g \times 2g}(\mathcal{O}_{\mathbb{C}_p})$. Consider the components \overline{A}'_1 on $\tilde{T}(\mathbb{C}_p)$ and \overline{A}'_2 in $\tilde{U}(\mathbb{C}_p)$. By construction they lie in the diagonal matrices and the upper triangular matrices in $M_{2g \times 2g, \mathcal{O}_{\mathbb{C}_p}}$ respectively. In particular, the various projections of \overline{A}'_2 via the decomposition (as schemes) $\tilde{U} \cong \mathbb{G}_a^r$ all lie in $\mathbb{G}_a(\mathcal{O}_{\mathbb{C}_p})$ and hence \overline{A}'_2 can be lifted to a matrix A'_2 in $\tilde{U}(A_{\text{inf}}) \subset M_{2g \times 2g}(A_{\text{inf}})$.

Fix an isomorphism $\tilde{T} \cong \mathbb{G}_m^d$ and lift $\overline{A}'_1 = (s'_1, \dots, s'_d)$ to an element A'_1 of $\tilde{T}[A_{\text{inf}}[\underline{p}]^{-1}]$ as explained above. Its image in GL_{2g} is a diagonal matrix of the form $([p^{\alpha_1}] \tilde{s}_1, \dots, [p^{\alpha_{2g}}] \tilde{s}_{2g})$ with $\tilde{s}_1, \dots, \tilde{s}_{2g} \in \mathbb{G}_m(A_{\text{inf}})$ and $\alpha_1, \dots, \alpha_{2g} \in \mathbb{Q}$. As the image of \overline{A}'_1 in the diagonal matrices of GL_{2g} lies in $M_{2g \times 2g}(\mathcal{O}_{\mathbb{C}_p})$ then each α_i must be non-negative. Then $A' := A'_1 \cdot A'_2$ is a lift of \overline{A}' in $\tilde{B}(A_{\text{inf}}[[\underline{p}]^{-1}])$ with the required property. \square

7 Appendix II: some facts about Weyl groups

We recall some facts from [7, Sect. 1.4] and [21, Sect. 3.5]. Let H be a connected reductive group over an algebraically closed field k . Fix a maximal torus T and a Borel B . Let P be a parabolic subgroup of H of type I . Every element w in W can be written uniquely as $w = w^J w_J$ for $w_J \in W_J$ and $w^J \in W^J$ and then $\ell(w) = \ell(w_J) + \ell(w^J)$. Similarly we have a unique decomposition $w = w_I {}^I w$ for $w_I \in W_I$ and ${}^I w \in {}^I W$ and then $\ell(w) = \ell(w_I) + \ell({}^I w)$. The groups W, W_J and W_I contain unique elements of maximal length, denoted by $w_0, w_{J,0}, w_{I,0}$ respectively. By maximality they are elements of order 2 and w_0 has the property that it sends positive roots to negative roots and that for every $w \in W$

$$\ell(w w_0) = \ell(w_0 w) = \ell(w_0) - \ell(w).$$

In particular, $J := {}^{w_0} I$ is the type opposite to I . We also deduce from the uniqueness of the decomposition above that in W^J and in ${}^I W$ we have unique elements of maximal length $w_0^J \in W^J$ and ${}^I w_0 \in {}^I W$, characterized by the property that $w_{I,0} {}^I w_0 = w_0 = w_0^J w_{J,0}$. Conjugation by w_0 defines an involution $\iota_0: W \rightarrow W$ preserving the length of elements. In particular, it preserves the set of simple reflections Δ . Then,

Lemma 7.1 *The map $\iota_0: W \rightarrow W, w \mapsto w_0 w w_0$, has the following properties*

- (a) ι_0 identifies $W_J = \iota_0(W_I)$ and $J = \iota_0(I)$. In particular, it induces mutually inverse and length preserving isomorphisms $W_J \rightarrow W_I$ and $W_I \rightarrow W_J$;
- (b) ι_0 identifies $W^J = \iota_0(W^I)$ (resp. ${}^J W = \iota_0({}^I W)$) and induces mutually inverse and length preserving bijections $W^J \rightarrow W^I$ and $W^I \rightarrow W^J$ (resp. ${}^J W \cong {}^I W$);
- (c) ι_0 identifies $w_{I,0} = \iota_0(w_{J,0}) = w_0 w_{J,0} w_0$ and, in particular, we have $w_0^J = {}^I w_0$ and $w_0^I = {}^J w_0$.

Proof All assertions are trivial except for the identity $w_0^J = {}^I w_0$ that follows from

$${}^I w_0 = w_{I,0} w_0 = w_0 w_{J,0} w_0 w_0 = w_0 w_{J,0} = w_0^J.$$

□

Lemma 7.2 (1) *The map $W \rightarrow W$, sending $w \rightarrow w^{-1}$, is an order and length preserving bijection. If $w = w^I w_I$ with $w^I \in W^I$ and $w_I \in W_I$ then $w^{-1} = w_I^{-1} (w^I)^{-1}$ and $w_I^{-1} \in W_I$ and $(w^I)^{-1} = {}^I w^{-1} \in {}^I W$.*

(2) *The map $W \rightarrow W$, sending $w \mapsto w_0 w w_{I,0}$, induces an order reversing bijection $W^I \rightarrow W^I$ such that $\ell(w_0 w w_{I,0}) = \ell(w_0^I) - \ell(w)$.*

Similarly the map $W \rightarrow W$, sending $w \mapsto w_{I,0} w w_0$, induces an order reversing bijection ${}^I W \rightarrow {}^I W$ such that $\ell(w_{I,0} w w_0) = \ell(w_0^I) - \ell(w)$.

(3) *The map $\sigma_0: W \rightarrow W$, sending $w \mapsto w_0^I w$, satisfies the following properties:*

- (i) *it is a bijection and it sends bijectively ${}^I W \rightarrow {}^J W$;*
- (ii) *it satisfies $\ell(w_0^I w) = \ell(w_0^I) - \ell(w)$ for every $w \in {}^I W$;*
- (iii) *it reverses the orders \preceq on ${}^I W$ and ${}^J W$, defined as follows. Let \leq be the Bruhat order on W and let $z = \varphi(w_0^I)$. For w and $w' \in {}^I W$ (resp. in ${}^J W$) we say that $w' \preceq w$ iff there exists $u \in W_I$ (resp. $u \in W_J$) such that $uw'z\varphi(u)^{-1}z^{-1} \leq w$ (resp. $uw'\varphi(u)^{-1} \leq w$).*

Proof (1) is clear.

(2) We prove the first statement. The second is obtained from the first using the order and length preserving bijection $W^I \rightarrow {}^I W$ given by $w \mapsto w^{-1}$ obtained from (1). First we compute lengths. For every $w \in W^I$ and every $y \in W_I$ we have $\ell(wy) = \ell(w) + \ell(y)$. Then $\ell(w_0 wy) = \ell(w_0) - \ell(y) - \ell(w)$. So for $y = w_{I,0}$ we get that $\ell(w_0 w w_{I,0}) = \ell(w_0) - \ell(w_{I,0}) - \ell(w) = \ell(w_0^I) - \ell(w)$ as wanted. On the other hand the function $y \mapsto \ell(w_0 wy)$ is minimized for $\ell(y)$ maximal, i.e., for $y = w_{I,0}$. Hence, $w_0 w w_{I,0}$ is the element in the coset $w_0 w W_I$ of minimal length, i.e., $w_0 w w_{I,0} \in W^I$.

(3) As $w_0^I = w_0 w_{I,0}$, the given map is the composite of the map $W \rightarrow W$, $w \mapsto w_{I,0} w w_0$, and of the map $\iota_0: W \rightarrow W$, $w \mapsto w_0 w w_0$. These are both involutions and hence σ_0 is a bijection on W . It sends bijectively ${}^I W$ to ${}^J W$, reversing the Bruhat orders \leq , thanks to Claim (2) and Lemma 7.1.

Write $y := w_0^I$ for simplicity. Take w and $w' \in {}^I W$ such that $w' \preceq w$ and let $u \in W_I$ be such that $uw'z\varphi(u)^{-1}z^{-1} \leq w$. This implies that $uw'z\varphi(u)^{-1}z^{-1} \in {}^I W$. We deduce that $yuw'z\varphi(u)^{-1}z^{-1} = (yuy^{-1})(y w')\varphi(yuy^{-1})^{-1} \geq yw$. Write $t := yu^{-1}y^{-1}$. Then $t(yw)\varphi(t)^{-1} \leq yw'$. Note that $yuy^{-1} = w_0^I u^I w_0 = w_0(w_{I,0} u w_{I,0}) w_0$ as $y^{-1} = {}^I w_0$ by Claim (1). Since $w_{I,0} u w_{I,0} \in W_I$, we conclude that $t \in W_J$ thanks to Lemma 7.1 so that $yw \preceq yw'$. □

Acknowledgements The author thanks the referees for their comments and suggestions.

Declarations

Conflict of interest The author declares that no conflict of interest exists, that is, he has no financial or personal relationship with a third party whose interests could be positively or negatively influenced by the present article’s content. All data generated or analysed during this study are included in this article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Bédard, R.: On the Brauer liftings for modular representations. *J. Algebra* **93**, 332–353 (1985)
- Berger, L.: Représentations p -adiques et équations différentielles. *Invent. Math.* **148**, 219–284 (2002)
- Breuil, C.: Représentations p -adiques semi-stables et transversalité de Griffiths. *Math. Ann.* **307**, 191–224 (1997)
- Brinon, O.: Représentations p -adiques cristallines de de Rham dans le cas relatif. *Mémoire. SMF* **112** (2008)
- Caraiani, A., Scholze, P.: On the generic part of the cohomology of compact unitary Shimura varieties. *Ann. Math.* **186**, 649–766 (2017)
- Chai, C.-L., Faltings, G.: *Degeneration of Abelian Varieties*. *Ergebnisse der Mathematik*, vol. 22. Springer, Berlin (1990)
- Goldring, W., Koskivirta, J.-S.: Strata Hasse invariants, Hecke algebras and Galois representations. *Invent. Math.* **217**, 887–984 (2019)
- He, X.: The G -stable pieces of the wonderful compactification. *Trans. Am. Math. Soc.* **359**, 300–3024 (2007)
- He, X.: G -stable pieces and partial flag varieties. In: *Representation Theory*. *Contemp. Math.*, vol. 478, pp. 61–70 (2009)
- Kim, W., Madapusi Pera, K.: 2-adic integral canonical models. *Forum Math. Sigma* **4**, 34 (2016)
- Kisin, M.: Integral models for Shimura varieties of abelian type. *J. Am. Math. Soc.* **23**, 967–1012 (2010)
- Kisin, M.: Mod p points on Shimura varieties of abelian type. *J. Am. Math. Soc.* **30**, 819–914 (2017)
- Lee, D.U.: Nonemptiness of Newton strata of Shimura varieties of Hodge type. *Algebra Number Theory* **12**, 259–283 (2018)
- Lusztig, G.: On the Green polynomials of classical groups. *Proc. Lond. Math. Soc.* **33**, 443–475 (1976)
- Lusztig, G.: On the finiteness of the number of unipotent classes. *Invent. Math.* **34**, 201–213 (1976)
- Madapusi Pera, K.: Toroidal compactifications of integral models of Shimura varieties of Hodge type. *Ann. Sci. Éc. Norm. Supér.* **52**, 393–514 (2019)
- Moonen, B., Wedhorn, T.: Discrete invariants of varieties in positive characteristic. *Int. Math. Res. Not.* **72**, 3855–3903 (2004)
- Pilloni, V., Stroth, B.: Cohomologie cohérente et représentations Galoisienne. *Ann. Math. Québec* **40**, 167–202 (2016)
- Pink, R.: *Arithmetical Compactification of Mixed Shimura Varieties* (Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, 1989). Universität Bonn Mathematisches Institut, Bonner Mathematische Schriften (1990)
- Pink, R., Wedhorn, T., Ziegler, P.: Algebraic Zip data. *Doc. Math.* **16**, 253–300 (2011)
- Pink, R., Wedhorn, T., Ziegler, P.: F -zips with additional structure. *Pacific J. Math.* **274**, 183–236 (2015)
- Scholze, P.: On torsion in the cohomology of locally symmetric varieties. *Ann. Math.* **182**, 945–1066 (2015)
- Wedhorn, T., Ziegler, P.: Tautological rings of Shimura varieties and cycle classes of Ekedahl-Oort strata, Available at [arXiv:1811.04843v3](https://arxiv.org/abs/1811.04843v3)
- Wortman, D.: The μ -ordinary locus for Shimura varieties of Hodge type, Available at [arXiv:1310.6444v1](https://arxiv.org/abs/1310.6444v1)
- Yu, C.-F.: On non-emptiness of basic loci of Shimura varieties. *MFO Workshop on Reductions of Shimura Varieties*. *Math. F. Oberwolfach Report No. 39/2015* (2016)

-
26. Zhang, C.: Ekedahl-Oort strata for good reductions of Shimura varieties of Hodge type. *Can. J. Math.* **70**, 451–480 (2018)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.