



Character bounds for regular semisimple elements and asymptotic results on Thompson’s conjecture

Michael Larsen¹ · Jay Taylor² · Pham Huu Tiep³

Received: 20 April 2022 / Accepted: 8 December 2022 / Published online: 23 January 2023
© The Author(s) 2023

Abstract

For every integer k there exists a bound $B = B(k)$ such that if the characteristic polynomial of $g \in \mathrm{SL}_n(q)$ is the product of $\leq k$ pairwise distinct monic irreducible polynomials over \mathbb{F}_q , then every element x of $\mathrm{SL}_n(q)$ of support at least B is the product of two conjugates of g . We prove this and analogous results for the other classical groups over finite fields; in the orthogonal and symplectic cases, the result is slightly weaker. With finitely many exceptions (p, q) , in the special case that $n = p$ is prime, if g has order $\frac{q^p-1}{q-1}$, then every non-scalar element $x \in \mathrm{SL}_p(q)$ is the product of two conjugates of g . The proofs use the Frobenius formula together with upper bounds for values of unipotent and quadratic unipotent characters in finite classical groups.

Contents

1	Introduction	2
2	Combinatorics	4
2.1	Arrays	4
2.2	Fourier transform	5
2.3	Hooks	5

The first author was partially supported by the NSF grant DMS-2001349. The third author gratefully acknowledges the support of the NSF (Grants DMS-1840702 and DMS-2200850), the Joshua Barlaz Chair in Mathematics, and the Charles Simonyi Endowment at the Institute for Advanced Study (Princeton). The authors are grateful to the referee for careful reading and helpful comments on the paper, in particular for the suggestion to add Corollary 2 to the paper.

✉ Jay Taylor
jay.taylor@manchester.ac.uk
Michael Larsen
mjlarson@indiana.edu
Pham Huu Tiep
pht19@math.rutgers.edu

¹ Department of Mathematics, Indiana University, Bloomington, IN 47405, USA

² Department of Mathematics, The University of Manchester, Oxford Road, Manchester M13 9PL, UK

³ Department of Mathematics, Rutgers University, Piscataway, NJ 08854, USA

2.4 Symbols	6
3 More combinatorics	7
3.1 A combinatorial result of Asai	7
3.2 More symbols	11
4 Hyperoctahedral groups	13
5 Lusztig series	17
6 A character bound from the Mackey formula	18
7 Further bounds from Deligne–Lusztig characters	20
8 Quadratic unipotent characters	23
9 Character degrees	26
10 Applications to asymptotic variants of Thompson’s conjecture	33
10.1 Type A	33
10.2 Types BCD	35
10.3 Another result for SL	38
References	44

1 Introduction

A conjecture of Thompson (cf. [2, p. 3]) states that each finite non-abelian simple group G contains a conjugacy class $C \subseteq G$ such that $C^2 = G$. The main result of [9] shows that this conjecture holds, unless G is a finite simple group of Lie type over a field of at most 8 elements. Note that Thompson’s conjecture (for a given simple group G) implies Ore’s conjecture [32] (for the same group G), which states that every element of G is a commutator, and which is now a theorem thanks to [26]. However, there has not been any further progress on Thompson’s conjecture until very recently, when the conjecture has been shown, see [21, Theorem 7.7], to hold for simple symplectic, unitary, or orthogonal groups (with some condition on the center of the Schur cover in the two latter cases) of sufficiently large rank.

Inspired by these results, we would like to study an asymptotic version of Thompson’s conjecture when G is one of the finite classical groups $SL_n(q)$, $SU_n(q)$, $Sp_{2n}(q)$, $SO_{2n+1}(q)$, and $SO_{2n}^\pm(q)$, which are all closely related to simple groups. This asymptotic version treats target elements of sufficiently large support. We prove that regular semisimple conjugacy classes $C = g^G$ satisfy our asymptotic version of Thompson’s conjecture whenever the characteristic polynomial of g is close to being irreducible.

If $G = Cl(V)$ is a finite classical group, with natural module $V = \mathbb{F}_q^n$, we define the support $\text{supp}(x)$ of an element $x \in G$ to be the codimension of the largest eigenspace of x on $V \otimes_{\mathbb{F}_q} \overline{\mathbb{F}_q}$. The following is one of our main results and generalizes [21, Theorem 7.8].

Theorem 1 *For all integers $k \in \mathbb{Z}_{\geq 1}$ there exists an explicit constant $B = B(k) > 0$ such that for all $n \in \mathbb{Z}_{\geq 1}$ and all prime powers q the following statement holds. Suppose G is one of $SL_n(q)$, $SU_n(q)$, $Sp_{2n}(q)$, $SO_{2n+1}(q)$, and $SO_{2n}^\pm(q)$, and $g \in G$ is a regular semisimple element whose characteristic polynomial on the natural module is a product of k pairwise distinct irreducible polynomials, of pairwise distinct degrees if G is of type Sp or SO . Then $g^G \cdot g^G$ contains every element $x \in [G, G]$ with $\text{supp}(x) \geq B$.*

In fact, in the Sp and SO cases, we prove a slightly stronger result, see Theorem 10.2. We also note that the assumption $x \in [G, G]$ is superfluous in the SL , SU , and Sp cases (since $G = [G, G]$ in these cases, aside from known exceptions with $n \leq 3$), but necessary in the SO case (since in this case $[G, G]$ has index 2 in G and so $g^G \cdot g^G \subseteq [G, G]$).

In a special family of particularly favorable cases, Theorem 10.7 shows that all non-central elements of G lie in C^2 .

If $\text{Irr}(G)$ denotes the set of the complex irreducible characters of G , then the well-known formula of Frobenius states that $x \in G$ is contained in $g^G \cdot g^G$ if and only if

$$\sum_{\chi \in \text{Irr}(G)} \frac{\chi(g)^2 \chi(x^{-1})}{\chi(1)} \neq 0 \tag{1.1}$$

Similarly, $x \in G$ is contained in $g^G \cdot (g^{-1})^G$ if and only if

$$\sum_{\chi \in \text{Irr}(G)} \frac{|\chi(g)|^2 \chi(x^{-1})}{\chi(1)} \neq 0 \tag{1.2}$$

Our proof of Theorem 1, but using (1.2) instead of (1.1), implies the following asymptotic version of Ore’s conjecture, in which one fixes the shape of one of the two arguments of the commutator:

Corollary 2 *For all integers $k \in \mathbb{Z}_{\geq 1}$ there exists an explicit constant $B = B(k) > 0$ such that for all $n \in \mathbb{Z}_{\geq 1}$ and all prime powers q the following statement holds. Suppose G is one of $\text{SL}_n(q)$, $\text{SU}_n(q)$, $\text{Sp}_{2n}(q)$, $\text{SO}_{2n+1}(q)$, and $\text{SO}_{2n}^{\pm}(q)$, and $g \in G$ is a regular semisimple element whose characteristic polynomial on the natural module is a product of k pairwise distinct irreducible polynomials, of pairwise distinct degrees if G is of type Sp or SO . Also set $\tilde{G} := \text{GL}_n(q)$ if $G = \text{SL}_n(q)$, $\tilde{G} := \text{GU}_n(q)$ if $G = \text{SU}_n(q)$, and $\tilde{G} := G$ otherwise. Then $g^{\tilde{G}} \cdot (g^{-1})^{\tilde{G}}$ contains every element $x \in [G, G]$ with $\text{supp}(x) \geq B$. In particular, every element $x \in [G, G]$ with $\text{supp}(x) \geq B$ is a commutator $[y, z]$, where $y \in g^{\tilde{G}}$ and $z \in \tilde{G}$.*

To prove (1.1) or (1.2), we need sufficiently good upper bounds on $|\chi(g)|$. To get these we realise our group as the fixed point subgroup \mathbf{G}^F of a Frobenius endomorphism $F : \mathbf{G} \rightarrow \mathbf{G}$ on a connected reductive algebraic group \mathbf{G} and use the Deligne–Lusztig theory [7].

To illustrate our techniques suppose $\mathbf{G}^F = \text{Sp}_{2n}(q)$ or $\text{SO}_{2n+1}(q)$. To each element w of the Weyl group $W \cong C_2 \wr \mathfrak{S}_n$ of \mathbf{G} , Deligne and Lusztig have associated a virtual character R_w of \mathbf{G}^F whose irreducible constituents are called *unipotent* characters. The subspace $\text{Class}_0(\mathbf{G}^F) \subseteq \text{Class}(\mathbf{G}^F)$ of all \mathbb{C} -valued class functions spanned by $\{R_w \mid w \in W\}$ is the space of uniform unipotent class functions.

If χ is a unipotent character then the (uniform) projection of χ onto $\text{Class}_0(\mathbf{G}^F)$ is known to have the form

$$\mathcal{R}_{f_\chi} = \frac{1}{|W|} \sum_{w \in W} f_\chi(w) R_w$$

for some class function $f_\chi \in \text{Class}(W)$, which is not irreducible in general. If $g \in \mathbf{G}^F$ is semisimple then its characteristic function is *uniform*, which means $\chi(g) = \mathcal{R}_{f_\chi}(g)$ for all χ .

Our first step towards understanding $\mathcal{R}_{f_\chi}(g)$ is to show that f_χ satisfies a version of the *recursive Murnaghan–Nakayama rule* (or MN-rule), see Theorem 4.8. This is a consequence of a fundamental combinatorial result of Asai [3], [4] that relates the classical MN-rule for the irreducible characters of W and Lusztig’s Fourier transform, whose proof we give in Sect. 3. If $g \in \mathbf{G}^F$ is a regular semisimple element such that $\mathbf{C}_{\mathbf{G}^F}^g$ is a torus of type w then $|\chi(g)| = |\mathcal{R}_{f_\chi}(g)| = |f_\chi(w)|$ and we recover the MN-rule of Lübeck–Malle [27, Thm. 3.3].

By working with uniform projections we may apply these results to non-unipotent characters, and we do so to obtain bounds on $|\chi(g)|$ whenever χ is a *quadratic unipotent character* and the *cycle type* of $g \in \mathbf{G}^F$ is a product of $k \geq 1$ pairwise distinct cycles (see §8 for precise definitions). In fact, following an argument of Larsen–Shalev [22] we obtain a bound

on $|\chi(g)|$ that depends only on k , see Corollary 8.2 in the quadratic unipotent case. Bounds for arbitrary characters, involving k and n , are given in Corollaries 7.6 and 7.8. Using the Mackey formula for tori we also give a bound in Theorem 6.2 that works for any irreducible character of any finite reductive algebraic group.

Now treating all characters χ in (1.1) involves a reduction to Levi subgroups using Deligne–Lusztig induction. The characters that contribute to the sum the most have a heavily restricted form. Our character bounds allow us to obtain sufficiently good bounds on the sum. Aside from these immediate applications, we believe our character bounds for regular semisimple elements will be useful in other situations as well.

2 Combinatorics

For any set X we will denote by $\text{Pow}(X)$ the set of all subsets of X of *finite cardinality*. This is naturally an \mathbb{F}_2 -vector space under symmetric difference, which we denote by

$$A \oplus B = (A \cup B) - (A \cap B)$$

for any $A, B \in \text{Pow}(X)$. Moreover, it is equipped with a nondegenerate symmetric bilinear form

$$\langle -, - \rangle : \text{Pow}(X) \times \text{Pow}(X) \rightarrow \mathbb{Z}/2\mathbb{Z}$$

given by $\langle A, B \rangle = |A \cap B| \pmod{2}$. If $e \in \mathbb{Z}$ then we let

$$\text{Pow}_e(X) = \{A \in \text{Pow}(X) \mid |A| \equiv e \pmod{2}\}.$$

We set $X^{(2)} = X \times \mathbb{Z}/2\mathbb{Z}$. If $X \subseteq Y$ then $X^{(2)} \subseteq Y^{(2)}$. Elements of $X^{(2)}$ will be identified with their representatives in $X \times \{0, 1\}$. We denote by $\delta : X^{(2)} \rightarrow \mathbb{Z}/2\mathbb{Z}$ the projection onto the second factor.

Set $\mathbb{N} = \{1, 2, 3, \dots\}$ and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. If $A \in \text{Pow}(\mathbb{N}_0)$ then we define the *rank* of A to be $\rho(A) = \sum_{a \in A} a - \binom{|A|}{2}$. For each $k \in \mathbb{N}_0$ we define a map $(-)^{\rightarrow k} : \text{Pow}(\mathbb{N}_0) \rightarrow \text{Pow}(\mathbb{N}_0)$ by setting $A^{\rightarrow k} = \{0, \dots, k-1\} \sqcup \{a+k \mid a \in A\}$. This gives an equivalence relation \sim by setting $A \sim B$ if $A = B^{\rightarrow k}$ or $B = A^{\rightarrow k}$ for some $k \in \mathbb{N}_0$.

We denote by $[A]$ the equivalence class containing A and $\mathcal{B} = \text{Pow}(\mathbb{N}_0)/\sim$ the set of all equivalence classes. These are called β -sets. The rank $\rho([A]) = \rho(A)$ of $[A] \in \mathcal{B}$ is well defined. If $n \in \mathbb{N}_0$ then $\mathcal{B}_n \subseteq \mathcal{B}$ denotes all β -sets of rank n .

2.1 Arrays

The elements of $\text{Pow}(\mathbb{Z}^{(2)})$ will be called *arrays*. They will be identified with their images under the natural bijection $\text{Pow}(\mathbb{Z}^{(2)}) \rightarrow \text{Pow}(\mathbb{Z}) \times \text{Pow}(\mathbb{Z})$ given by $X \mapsto (X^0, X^1)$, where $X^i = \{x \in \mathbb{Z} \mid (x, i) \in X\}$. We say X^0 is the *top row* of X and X^1 the *bottom row* of X .

Following Lusztig [28], and modifying the notation of [34], we consider elements of $\tilde{\mathcal{P}} = \text{Pow}(\mathbb{N}_0^{(2)}) \subseteq \text{Pow}(\mathbb{Z}^{(2)})$. Recall the *rank* of $X \in \tilde{\mathcal{P}}$ is defined to be

$$\text{rk}(X) = \sum_{x^0 \in X^0} x^0 + \sum_{x^1 \in X^1} x^1 - \left[\left(\frac{|X| - 1}{2} \right)^2 \right] = \rho(X^0) + \rho(X^1) + \left[\left(\frac{\text{def } X}{2} \right)^2 \right] \tag{2.1}$$

where $\text{def}(\Lambda) = |X^0| - |X^1|$ is the *defect* of Λ .

For $d \in \mathbb{Z}$ we let $\tilde{\mathcal{P}}^d \subseteq \tilde{\mathcal{P}}$ be the set of arrays of defect d . We set $\tilde{\mathcal{P}}^{\text{od}} = \bigsqcup_{d \in \mathbb{Z}} \tilde{\mathcal{P}}^{2d+1}$ and $\tilde{\mathcal{P}}^{\text{ev}} = \bigsqcup_{d \in \mathbb{Z}} \tilde{\mathcal{P}}^{2d}$ so that $\tilde{\mathcal{P}} = \tilde{\mathcal{P}}^{\text{od}} \sqcup \tilde{\mathcal{P}}^{\text{ev}}$. For $n \in \mathbb{N}_0$ we let $\tilde{\mathcal{P}}_n \subseteq \tilde{\mathcal{P}}$ be the set of arrays of rank n . We then set $\tilde{\mathcal{P}}_n^d = \tilde{\mathcal{P}}_n \cap \tilde{\mathcal{P}}^d$, $\tilde{\mathcal{P}}_n^{\text{od}} = \tilde{\mathcal{P}}_n \cap \tilde{\mathcal{P}}^{\text{od}}$, and $\tilde{\mathcal{P}}_n^{\text{ev}} = \tilde{\mathcal{P}}_n \cap \tilde{\mathcal{P}}^{\text{ev}}$.

Each $X \in \tilde{\mathcal{P}}$ gives rise to the following subsets of \mathbb{N}_0 : $X^\cup := X^0 \cup X^1$, $X^\cap = X^0 \cap X^1$, $X^\ominus = X^0 \ominus X^1$. We let $\text{Sim}(X) = \{Y \in \tilde{\mathcal{P}} \mid Y^\cup = X^\cup \text{ and } Y^\cap = X^\cap\}$ be the *similarity class* of X . All elements of $\text{Sim}(X)$ have the same rank but different defects.

2.2 Fourier transform

We associate to each $X \in \tilde{\mathcal{P}}$ an associated *special array*

$$X_{\text{sp}} := \{(y, 0), (y, 1) \mid y \in X^\cap\} \cup \{(x, |X^\ominus| + \langle X^\ominus, \{0, \dots, x\} \rangle) \mid x \in X^\ominus\}. \tag{2.2}$$

Note that if $Y \in \text{Sim}(X)$ then $Y_{\text{sp}} = X_{\text{sp}}$. Moreover, if $x \in X^\ominus$ then we have

$$\langle X_{\text{sp}}, \{x\} \rangle \equiv |X^\ominus| + \langle X^\ominus, \{0, \dots, x\} \rangle. \tag{2.3}$$

The defect $\text{def}(X_{\text{sp}}) \in \{0, 1\}$ of this array satisfies $\text{def}(X_{\text{sp}}) \equiv |X^\ominus| \pmod{2}$. Thus we have an integer

$$s(X) = 2^{(|X^\ominus| - \text{def}(X_{\text{sp}}))/2} \in \mathbb{N}.$$

We have a map $\sharp : \tilde{\mathcal{P}} \rightarrow \text{Pow}(\mathbb{N}_0)$ given by $Y^\sharp = Y^1 \ominus Y_{\text{sp}}^1 \subseteq Y^\ominus$. This restricts to a bijection $\sharp : \text{Sim}(X) \rightarrow \text{Pow}(X^\ominus)$ for any $X \in \tilde{\mathcal{P}}$. With this we define a \mathbb{C} -linear map $\tilde{\mathcal{R}} : \mathbb{C}[\tilde{\mathcal{P}}] \rightarrow \mathbb{C}[\tilde{\mathcal{P}}]$ by setting

$$\tilde{\mathcal{R}}(X) = \frac{1}{s(X)} \sum_{Y \in \text{Sim}(X)} (-1)^{\langle X^\sharp, Y^\sharp \rangle} Y.$$

Here $\mathbb{C}[\tilde{\mathcal{P}}]$ denotes the free \mathbb{C} -module with basis $\tilde{\mathcal{P}}$ and $\langle -, - \rangle$ is the symmetric \mathbb{F}_2 -bilinear form defined above. Up to scaling this is the Fourier transform of the abelian group $\text{Pow}(X^\ominus)$.

Remark 2.1 We briefly make a few comments regarding the conventions and definitions in [28]. If $M = X^1 \cap X^\ominus$ and $M_0 = X_{\text{sp}}^1 \cap X^\ominus$ then we have $X^\sharp = M \ominus M_0$, which is denoted by M^\sharp in [28, §4.5, §4.6]. We have another set $M'_0 = X_{\text{sp}}^0 \cap X^\ominus = X^\ominus - M_0$. If $|X^\ominus|$ is even then M_0 and M'_0 are distinguished by the condition that $\sum_{x \in M_0} x < \sum_{x \in M'_0} x$.

This definite choice of M_0 over M'_0 , using X_{sp} , is used in [28, §4.18]. Moreover, our definition of X_{sp} agrees with the convention in [29, 17.2], that the smallest entry of X^\ominus occurs in the lower row of X_{sp} . This is different to the definition of a distinguished symbol given in [13, 4.4.3].

2.3 Hooks

Each $(d, i) \in \mathbb{Z}^{(2)}$ determines an injective function $\mathcal{D}_{d,i} : \mathbb{Z}^{(2)} \rightarrow \mathbb{Z}^{(2)}$ given by $\mathcal{D}_{d,i}((x, j)) = (x - d, i + j)$. This induces a map $\mathcal{D}_{d,i} : \text{Pow}(\mathbb{Z}^{(2)}) \rightarrow \text{Pow}(\mathbb{Z}^{(2)})$ that is \mathbb{F}_2 -linear. For any $X, H \in \text{Pow}(\mathbb{Z}^{(2)})$ we define

$$X \setminus_{d,i} H = X \ominus H \ominus \mathcal{D}_{d,i}(H) \in \text{Pow}(\mathbb{Z}^{(2)}).$$

We write this as $X \setminus H$ when (d, i) is clear from the context or by $X \setminus_{d,i} \lambda = X \setminus \lambda$ when $H = \{\lambda\}$ is a singleton. For brevity we write the map $\mathcal{D}_{0,1}$ simply as $(-)^{\text{op}}$.

If $X \in \tilde{\mathcal{P}}$ then the elements of the set

$$\mathcal{H}_{d,i}(X) = \{\lambda \in X \mid \mathcal{D}_{d,i}(\lambda) \in \mathbb{N}_0^{(2)} - X\}$$

are called the (d, i) -hooks of X . Elsewhere in the literature $(d, 0)$ -hooks and $(d, 1)$ -hooks, with $d > 0$, are called d -hooks and d -cohooks respectively. Following [34] we define the *leg parity* of $\lambda = (x, j) \in \mathcal{H}_{d,i}(X)$ to be

$$l_{d,i}(\lambda, X) = \langle \{0, \dots, x\}, X^j \rangle + \langle \{0, \dots, x - d\}, (X \setminus_{d,i} \lambda)^{i+j} \rangle.$$

If $d > 0$ and $i = 0$ then this has the same parity as the usual notion of the leg length of a hook. This is also easily seen to agree with the definitions in [13, 4.4.10].

Remark 2.2 The similarity relation can be rephrased in terms of $(0, 1)$ -hooks. Specifically, we have bijections $\mathcal{H}_{0,1}(X) \rightarrow X^\ominus$ and $\mathcal{H}_{0,1}(X) \rightarrow \text{Sim}(X)$ given by $H \mapsto H^\cup = H^\ominus$ and $H \mapsto X \setminus_{0,1} H$ respectively. Moreover, if $Y = X \setminus_{0,1} H$ then we have $Y^j = X^j \ominus H^\ominus$ for any $j \in \{0, 1\}$.

Recall that $\delta : X^{(2)} \rightarrow \mathbb{Z}/2\mathbb{Z}$ is the second projection map. For a pair $(d, i) \in \mathbb{Z}^{(2)}$, with $d \neq 0$, and $j \in \{0, 1\}$ we define a \mathbb{C} -linear map $\tilde{\mathcal{H}}_{d,i}^j : \mathbb{C}[\tilde{\mathcal{P}}] \rightarrow \mathbb{C}[\tilde{\mathcal{P}}]$ by setting

$$\tilde{\mathcal{H}}_{d,i}^j(X) = \sum_{\lambda \in \mathcal{H}_{d,i}(X)} (-1)^{j\delta(\lambda) + l_{d,i}(\lambda, X)} X \setminus_{d,i} \lambda$$

for any $X \in \tilde{\mathcal{P}}$. Note that if $Y = X \setminus_{d,i} \lambda$, with $\lambda \in \mathcal{H}_{d,i}(X)$, then $\text{def}(Y) = \text{def}(X) - 2i(-1)^{\delta(\lambda)}$ so $\text{def}(Y) \equiv \text{def}(X) \equiv |X^\ominus| \pmod{2}$.

2.4 Symbols

The map $(-)^{\rightarrow k} : \text{Pow}(\mathbb{N}_0) \rightarrow \text{Pow}(\mathbb{N}_0)$ defined above, for $k \in \mathbb{N}_0$, extends to map $\tilde{\mathcal{P}} \rightarrow \tilde{\mathcal{P}}$ given by $(A, B)^{\rightarrow k} = (A^{\rightarrow k}, B^{\rightarrow k})$. As before this yields an equivalence relation on $\tilde{\mathcal{P}}$. We denote by $[X]$ the equivalence class containing $X \in \tilde{\mathcal{P}}$ and $\tilde{\mathcal{S}}$ the set of equivalence classes. The equivalence class $[X]$ is called an (ordered) symbol.

Given $\lambda = (x, j) \in \mathbb{Z}^{(2)}$ and $k \in \mathbb{Z}$ let $\lambda + k = (x + k, j)$. If $X \in \tilde{\mathcal{P}}$ and $k \in \mathbb{N}_0$ then it is readily checked that:

- $\mathcal{H}_{d,i}(X^{\rightarrow k}) = \{\lambda + k \mid \lambda \in \mathcal{H}_{d,i}(X)\}$,
- $X^{\rightarrow k} \setminus_{d,i} (\lambda + k) = (X \setminus_{d,i} \lambda)^{\rightarrow k}$
- $l_{d,i}(\lambda + k, X^{\rightarrow k}) = l_{d,i}(\lambda, X)$.

Thus the maps $\tilde{\mathcal{R}}$ and $\tilde{\mathcal{H}}_{d,i}^j$ preserve the kernel of the natural quotient map $\mathbb{C}[\tilde{\mathcal{P}}] \rightarrow \mathbb{C}[\tilde{\mathcal{S}}]$ and so factor through endomorphisms of $\mathbb{C}[\tilde{\mathcal{S}}]$ which we denote in the same way.

Recall that $X \in \tilde{\mathcal{P}}$ is degenerate if $X^0 = X^1$. We let

$$\mathcal{S} = \{[X] \mid X \in \tilde{\mathcal{P}} \text{ and } X^0 \neq X^1\} \cup \{[X]_\pm \mid X \in \tilde{\mathcal{P}} \text{ and } X^0 = X^1\}$$

where $[X] = \{[X], [X^{\text{op}}]\}$. We take the rank and defect of $[X] \in \mathcal{S}$ to be $\text{rk}([X]) = \text{rk}(X)$ and $\text{def}([X]) = |\text{def}(X)|$. We can then partition \mathcal{S} with respect to the rank and defect as in Sect. 2.1.

3 More combinatorics

3.1 A combinatorial result of Asai

We will now prove the following fundamental combinatorial observation of Asai that relates the maps $\tilde{\mathcal{R}}$ and $\tilde{\mathcal{H}}_{d,i}^j$. This was first stated by Asai in [3, Lem. 2.8.3] and [4, Lem. 1.5.3] where it is left as a “direct calculation”. However, as pointed out by Lübeck–Malle [27, §3.4] a sign is missing in the statement in [3].

Waldspurger also states a version of this result [34, §2], where it is left as “un calcul fastidieux mais élémentaire”, but the conventions of [34] are different leading to a different statement. Specifically the analogue of our map $\tilde{\mathcal{R}}$, denoted by \mathcal{F} in [34], is not the same, as can be seen by evaluating it on the similarity class $\{(\emptyset, \{1, 2\}), (\{1\}, \{2\}), (\{2\}, \{1\}), (\{1, 2\}, \emptyset)\}$.

Aside from being an important ingredient in our work here, Asai’s combinatorics form a core basis for the block theory of finite reductive groups and solutions of Lusztig’s conjecture on almost characters for classical groups. In this second application the correctness of signs is crucial. In light of the importance of Asai’s statements, we provide some details regarding the proof.

We note that some of the main ideas of the proof have recently appeared in [30, Prop. 6], where a weaker statement, leading essentially to Theorem 4.8, is proved. Unfortunately there are several errors in the proofs of [30] that are corrected by our arguments here.

Theorem 3.1 (Asai) *For any $0 \neq d \in \mathbb{Z}$ we have the following equalities of linear endomorphisms of $\mathbb{C}[\tilde{\mathcal{P}}]$:*

- (i) $\tilde{\mathcal{H}}_{d,0}^0 \circ \tilde{\mathcal{R}} = \tilde{\mathcal{R}} \circ \tilde{\mathcal{H}}_{d,0}^0$,
- (ii) $\tilde{\mathcal{H}}_{d,0}^1 \circ \tilde{\mathcal{R}} = -\Theta \circ \tilde{\mathcal{R}} \circ \tilde{\mathcal{H}}_{d,1}^0$,

where $\Theta : \mathbb{C}[\tilde{\mathcal{P}}] \rightarrow \mathbb{C}[\tilde{\mathcal{P}}]$ is the \mathbb{C} -linear map defined by $\Theta(X) = (-1)^{\text{def}(X)} X$.

From now on $0 \neq d \in \mathbb{Z}$ and $i \in \{0, 1\}$ are fixed. Given $x \in \mathbb{N}$ we denote by $\mathcal{X}(X, x)$ the set of pairs (H, λ) with $H \subseteq \mathcal{H}_{0,1}(X)$, and $\lambda \in \mathcal{H}_{d,0}(X \setminus H)$ is such that $\lambda = (x, j)$ for some $j \in \{0, 1\}$. Correspondingly we denote by $\mathcal{Y}(X, x)$ the set of pairs (μ, G) where $G \subseteq \mathcal{H}_{0,1}(X)$, and $\mu \in \mathcal{H}_{d,i}(X \setminus G)$ satisfies $\mu = (x, j)$ for some $j \in \{0, 1\}$.

Given $X \in \tilde{\mathcal{P}}$ we then have

$$\tilde{\mathcal{H}}_{d,0}^i(\tilde{\mathcal{R}}(\Lambda)) = \sum_{x \in \mathbb{N}} \sum_{(H, \lambda) \in \mathcal{X}(X, x)} \frac{1}{s(X)} (-1)^{i\delta(\lambda) + \langle X^\sharp, (X \setminus H)^\sharp \rangle + I_{d,0}(\lambda, X \setminus H)} X \setminus H \setminus \lambda \quad (3.1)$$

and

$$\tilde{\mathcal{R}}(\tilde{\mathcal{H}}_{d,i}^0(\Lambda)) = \sum_{x \in \mathbb{N}} \sum_{(\mu, G) \in \mathcal{Y}(\Lambda, x)} \frac{1}{s(X \setminus \mu)} (-1)^{\langle (X \setminus \mu)^\sharp, (X \setminus \mu \setminus G)^\sharp \rangle + I_{d,i}(\mu, X)} X \setminus \mu \setminus G. \quad (3.2)$$

Before proving (i) of Theorem 3.1 we start with a lemma.

Lemma 3.2 *Assume $i = 0$ and $(H, \lambda) \in \mathcal{X}(X, x)$ and $(\mu, G) \in \mathcal{Y}(X, x)$ are two terms satisfying one of the following:*

- (i) $(\mu, G^\ominus) = (\lambda, H^\ominus)$ and $\langle \{x, x - d\}, H^\ominus \rangle = 0$,
- (ii) $(\mu, G^\ominus) = (\lambda^{op}, H^\ominus \ominus \{x, x - d\})$ and $\langle \{x, x - d\}, H^\ominus \rangle = \langle \{x, x - d\}, X^\ominus \rangle = 1$.

Then we have

$$(-1)^{\langle X^\sharp, (X \setminus H)^\sharp \rangle + I_{d,0}(\lambda, X \setminus H)} = (-1)^{\langle (X \setminus \mu)^\sharp, (X \setminus \mu \setminus G)^\sharp \rangle + I_{d,0}(\mu, X)}.$$

Proof Let $Y = X \setminus_{0,1} H$, $U = X \setminus_{d,0} \mu$, and $V = U \setminus_{0,1} G$. We have

$$\langle X^\sharp, Y^\sharp \rangle + \langle U^\sharp, V^\sharp \rangle = |X^1| + |U^1| + |X_{sp}^1| + |U_{sp}^1| + \langle X^\sharp, H^\ominus \rangle + \langle U^\sharp, G^\ominus \rangle$$

because $V^\sharp = U^\sharp \ominus G^\ominus$ and $Y^\sharp = X^\sharp \ominus H^\ominus$. The sum of the first four terms is 0 because $|X^1| = |U^1|$, as μ is a $(d, 0)$ -hook, and a straightforward check shows that $|U_{sp}^1| = |X_{sp}^1|$. We thus have

$$\langle X^\sharp, Y^\sharp \rangle + \langle U^\sharp, V^\sharp \rangle = \langle U^1, H^\ominus \ominus G^\ominus \rangle + \langle U_{sp}^1, H^\ominus \ominus G^\ominus \rangle + \langle X^1 \ominus U^1, H^\ominus \rangle + \langle X_{sp}^1 \ominus U_{sp}^1, H^\ominus \rangle.$$

As $U^{\delta(\mu)} = X^{\delta(\mu)} \ominus \{x, x - d\}$ and $Y^{\delta(\lambda)} = X^{\delta(\lambda)} \ominus H^\ominus$, it is straightforward to see that

$$l_{d,0}(\lambda, Y) + l_{d,0}(\mu, X) = \langle \{0, \dots, x\} \ominus \{0, \dots, x - d\}, X^{\delta(\lambda)} \ominus X^{\delta(\mu)} \ominus H^\ominus \rangle.$$

We have to show the sum of these two expressions is 0.

For any $z \in U^\ominus \cap X^\ominus$ it follows from Eq. 2.3 that

$$\langle U_{sp}^1 \ominus X_{sp}^1, \{z\} \rangle = \langle U^\ominus \ominus X^\ominus, \{0, \dots, z\} \rangle = \langle \{0, \dots, x\} \ominus \{0, \dots, x - d\}, \{z\} \rangle$$

because $U^\ominus \ominus X^\ominus = \{x - d, x\}$ so $|X^\ominus| \equiv |U^\ominus| \pmod{2}$. Thus for any subset $Z \subseteq U^\ominus \cap X^\ominus$ we get

$$\langle U_{sp}^1 \ominus X_{sp}^1, Z \rangle = \langle \{0, \dots, x\} \ominus \{0, \dots, x - d\}, Z \rangle.$$

Note that $X^1 \ominus U^1$ is either \emptyset or $\{x, x - d\}$ depending on whether $\delta(\mu) = 0$ or 1. Hence the statement clearly follows if (i) holds. So assume (ii) holds. As $H^\ominus \subseteq X^\ominus$ we must have $H^\ominus \cap \{x, x - d\} = X^\ominus \cap \{x, x - d\}$. We will assume this is $\{x\}$ as the case where it is $\{x - d\}$ is identical. This means $x - d \notin X^\cup$ because $x - d \notin X^\cap$.

Clearly $x \notin U^\cup$ and $x - d \in U^\ominus$ so

$$\langle U^1, H^\ominus \ominus G^\ominus \rangle + \langle X^1 \ominus U^1, H^\ominus \rangle = \langle U^1, \{x - d\} \rangle + \langle X^1, \{x\} \rangle = 0.$$

Now $\langle U_{sp}^1, H^\ominus \ominus G^\ominus \rangle + \langle X_{sp}^1 \ominus U_{sp}^1, \{x\} \rangle = \langle U_{sp}^1, \{x - d\} \rangle + \langle X_{sp}^1, \{x\} \rangle$ is the same as

$$\langle \{x, x - d\}, \{0, \dots, x - d\} \rangle + \langle X^\ominus, \{0, \dots, x\} \ominus \{0, \dots, x - d\} \rangle,$$

and the first term is equivalent to $\langle \{x\}, \{0, \dots, x\} \ominus \{0, \dots, x - d\} \rangle$. Adding this to

$$\langle X_{sp}^1 \ominus U_{sp}^1, H^\ominus \ominus \{x\} \rangle$$

gives the statement because $H^\ominus \ominus \{x\} \subseteq U^\ominus \cap X^\ominus$. □

Proof of Theorem 3.1(i) We can assume $x \in X^\cup$ since otherwise $\mathcal{X}(X, x)$ and $\mathcal{Y}(X, x)$ are empty, and there is nothing to show. Similarly, this is the case if $x - d \in X^\cap$. The proof divides into several cases distinguished by the distribution of $x - d$ and x amongst the rows of X .

Case 1: $x \in X^\ominus$ and $x - d \notin X^\cup$. We have a bijection $\mathcal{X}(X, x) \rightarrow \mathcal{Y}(X, x)$ given by

$$(H, \lambda) \mapsto (\mu, G) := \begin{cases} (\lambda, H) & \text{if } H^\ominus \cap \{x, x - d\} = \emptyset \\ (\lambda^{\text{op}}, H \ominus \{\lambda^{\text{op}}, \mathcal{D}_{d,0}(\lambda^{\text{op}})\}) & \text{if } H^\ominus \cap \{x, x - d\} = \{x\} \end{cases}$$

such that $X \setminus H \setminus \lambda = X \setminus \mu \setminus G$. Now $(X \setminus \mu)^\ominus = (X^\ominus - \{x\}) \sqcup \{x - d\}$, so we have $s(X \setminus \mu) = s(X)$. Hence, by Lemma 3.2 the coefficients of $X \setminus H \setminus \lambda$ and $X \setminus \mu \setminus G$ in Eqs.(3.1) and (3.2) are the same.

Case 2: $x \in X^\cap$ and $x - d \notin X^\cup$. We have a fixed-point free involution $' : \mathscr{D}(\Lambda, x) \rightarrow \mathscr{D}(\Lambda, x)$ given by $(\mu, G)' = (\mu^{\text{op}}, G')$, where G' is defined by

$$G \ominus G' = \begin{cases} \{\mu, \mathcal{D}_{d,0}(\mu^{\text{op}})\} & \text{if } G^\ominus \cap \{x, x - d\} = \emptyset \\ \{\mu^{\text{op}}, \mathcal{D}_{d,0}(\mu^{\text{op}})\} & \text{if } G^\ominus \cap \{x, x - d\} = \{x\} \\ \{\mu, \mathcal{D}_{d,0}(\mu)\} & \text{if } G^\ominus \cap \{x, x - d\} = \{x - d\} \\ \{\mu^{\text{op}}, \mathcal{D}_{d,0}(\mu)\} & \text{if } G^\ominus \cap \{x, x - d\} = \{x, x - d\}. \end{cases}$$

This bijection satisfies

$$X \setminus \mu \setminus G = X \setminus \mu^{\text{op}} \setminus G' \text{ and } \langle \{x, x - d\}, G^\ominus \rangle = \langle \{x, x - d\}, G'^\ominus \rangle$$

because $G^\ominus \ominus G'^\ominus = \{x, x - d\}$. Given $e \in \{0, 1\}$ we let $\mathscr{D}^e(\Lambda, x)$ be the set of pairs $\{(\mu, G), (\mu, G)'\}$ with $\langle \{x, x - d\}, G^\ominus \rangle = \bar{e}$, where bar indicates reduction (mod 2).

We claim the coefficients of $V = X \setminus \mu \setminus G$ and $V' = X \setminus \mu^{\text{op}} \setminus G'$ in Eq. (3.2) differ by $(-1)^{\langle \{x, x - d\}, G^\ominus \rangle}$. If $U = X \setminus \mu$ and $U' = X \setminus \mu^{\text{op}}$ then as in Lemma 3.2 we get that

$$\langle U^\sharp, V^\sharp \rangle + \langle U'^\sharp, V'^\sharp \rangle = \langle U^\sharp, G^\ominus \rangle + \langle U'^\sharp, G'^\ominus \rangle.$$

As $G^\ominus \ominus G'^\ominus = \{x, x - d\}$, this term can be written as

$$\begin{aligned} & \langle U^1, \{x, x - d\} \rangle + \langle U^1_{\text{sp}}, \{x, x - d\} \rangle + \langle U^1 \ominus U'^1, G^\ominus \rangle + \langle U^1_{\text{sp}} \ominus U'^1_{\text{sp}}, G^\ominus \rangle \\ & = 1 + \langle U^\ominus, \{0, \dots, x\} \ominus \{0, \dots, x - d\} \rangle + \langle \{x, x - d\}, G^\ominus \rangle + 0 \\ & = \langle X^\ominus, \{0, \dots, x\} \ominus \{0, \dots, x - d\} \rangle + \langle \{x, x - d\}, G^\ominus \rangle. \end{aligned}$$

Finally, as in Lemma 3.2,

$$l_{d,0}(\mu, X) + l_{d,0}(\mu^{\text{op}}, X) = \langle \{0, \dots, x\} \ominus \{0, \dots, x - d\}, X^\ominus \rangle.$$

Therefore, given a pair $\{(\mu, G), (\mu^{\text{op}}, G')\} \in \mathscr{D}^1(X, x)$, the corresponding terms $X \setminus \mu \setminus G$ and $X \setminus \mu^{\text{op}} \setminus G'$ in Eq. (3.2) cancel. We have a bijection $\mathscr{X}(X, x) \rightarrow \mathscr{D}^0(\Lambda, x)$ given by $(H, \lambda) \mapsto \{(\lambda, H), (\lambda, H)'\}$ such that $X \setminus H \setminus \lambda = X \setminus \lambda \setminus H$. Now $s(X \setminus \lambda) = 2s(X)$ because $(X \setminus \lambda)^\ominus = X^\ominus \sqcup \{x, x - d\}$, but the coefficients of $X \setminus \lambda \setminus H$ and $X \setminus \lambda^{\text{op}} \setminus H'$ combine to yield the coefficient of $X \setminus H \setminus \lambda$ by Lemma 3.2.

Case 3: $x \in X^\ominus$ and $x - d \in X^\ominus$. Identically to Case 2 we have a fixed-point free involution $' : \mathscr{X}(X, x) \rightarrow \mathscr{X}(X, x)$ denoted by $(H, \lambda)' = (H', \lambda^{\text{op}})$. This bijection satisfies

$$X \setminus H' \setminus \lambda^{\text{op}} = X \setminus H \setminus \lambda \text{ and } \langle \{x, x - d\}, H^\ominus \rangle = \langle \{x, x - d\}, H'^\ominus \rangle.$$

One can check that when the terms exist, the coefficients of $X \setminus H \setminus \lambda$ and $X \setminus H' \setminus \lambda^{\text{op}}$ in Eq. (3.1) differ by $(-1)^{1 + \langle \{x, x - d\}, X^1 \rangle}$. Given $e \in \{0, 1\}$ we let $\mathscr{X}^e(X, x)$ be the set of pairs $\{(H, \lambda), (H, \lambda)'\}$ with $\langle \{x, x - d\}, H^\ominus \rangle = \bar{e}$.

If $x - d$ and x occur in the same row of X then we must have $\mathscr{X}^0(X, x) = \{\emptyset\}$ and $\mathscr{X}(X, x) = \{\emptyset\}$. Moreover, each pair in $\mathscr{X}^1(X, x)$ gives rise to terms in Eq. (3.1) that cancel. If $x - d$ and x occur in opposite rows of X then we must have $\mathscr{X}^1(X, x) = \{\emptyset\}$. Moreover, we have a bijection $\mathscr{D}(X, x) \rightarrow \mathscr{D}^0(\Lambda, x)$ given by $(\mu, G) \mapsto \{(G, \mu), (G, \mu)'\}$ such that $X \setminus \mu \setminus G = X \setminus G \setminus \mu = X \setminus G' \setminus \mu^{\text{op}}$. In this case $(X \setminus \mu)^\ominus = X^\ominus - \{x, x - d\}$ so $2s(X \setminus \mu) = s(X)$ but $X \setminus H \setminus \lambda$ and $X \setminus H' \setminus \lambda^{\text{op}}$ have the same coefficient in Eq. (3.1). Thus, the coefficients of $X \setminus \mu \setminus G = X \setminus G \setminus \mu$ agree by Lemma 3.2

Case 4: $x \in X^\cap$ and $x - d \in X^\ominus$. As in Case 1, we have a bijection $\mathcal{X}(X, x) \rightarrow \mathcal{Y}(X, x)$ given by

$$(H, \lambda) \mapsto (\mu, G) := \begin{cases} (\lambda, H) & \text{if } H^\ominus \cap \{x, x - d\} = \emptyset \\ (\lambda^{\text{op}}, H \ominus \{\lambda, \mathcal{D}_{d,0}(\lambda)\}) & \text{if } H^\ominus \cap \{x, x - d\} = \{x - d\} \end{cases}$$

such that $X \setminus H \setminus \lambda = X \setminus \mu \setminus G$. Clearly $(X \setminus \mu)^\ominus = (X^\ominus - \{x - d\}) \sqcup \{x\}$ so $s(X \setminus \mu) = s(X)$. Again, by Lemma 3.2 the coefficients of $X \setminus H \setminus \lambda$ and $X \setminus \mu \setminus G$ \square in Eqs. (3.1) and (3.2) are the same.

We now consider the proof of (ii) of Theorem 3.1. The argument is exactly the same as (i), proceeding through the same cases. The bijection in Case (ii) is defined identically simply replacing $\mathcal{D}_{d,0}$ with $\mathcal{D}_{d,1}$. Instead of providing a direct analogue of Lemma 3.2 we instead check directly in each case that the signs of the corresponding coefficients agree. As an example, we treat the analogue of Case 1, leaving the remaining cases to the reader.

Proof of Theorem 3.1(ii) Assume $x \in X^\ominus$ and $x - d \notin X^\cup$. We have a bijection $\mathcal{X}(X, x) \rightarrow \mathcal{Y}(X, x)$ given by

$$(H, \lambda) \mapsto (\mu, G) := \begin{cases} (\lambda, H \ominus \{\mathcal{D}_{d,1}(\lambda)\}) & \text{if } H^\ominus \cap \{x, x - d\} = \emptyset \\ (\lambda^{\text{op}}, H \ominus \{\lambda^{\text{op}}\}) & \text{if } H^\ominus \cap \{x, x - d\} = \{x\} \end{cases}$$

such that $X \setminus H \setminus \lambda = X \setminus \mu \setminus G$. As in the proof of (i) of Theorem 3.1, we need only check that the sign of the coefficient of $Y \setminus \lambda$ in Eq. (3.1) and $(-1)^{1+\text{def}(V)}$ times the sign of the coefficient of $V = U \setminus G$ in Eq. (3.2) agree, where $Y = X \setminus H$ and $U = X \setminus \mu$. We check this directly.

As μ is a $(d, 1)$ -hook, we have $|U^1| = |X^1| \pm 1$, so arguing as in the proof of Lemma 3.2, we see that $1 + \text{def}(V) + \langle X^\sharp, Y^\sharp \rangle + \langle U^\sharp, V^\sharp \rangle$ is

$$\text{def}(V) + \langle U^1, H^\ominus \ominus G^\ominus \rangle + \langle U_{\text{sp}}^1, H^\ominus \ominus G^\ominus \rangle + \langle X^1 \ominus U^1, H^\ominus \rangle + \langle X_{\text{sp}}^1 \ominus U_{\text{sp}}^1, H^\ominus \rangle.$$

Moreover, this time $l_{d,0}(\lambda, Y) + l_{d,1}(\mu, X)$ is

$$\langle \{0, \dots, x\} \ominus \{0, \dots, x - d\}, X^{\delta(\lambda)} \ominus X^{\delta(\mu)} \ominus H^\ominus \rangle + \langle \{0, \dots, x - d\}, X^\ominus \ominus \{x\} \rangle.$$

We consider the two cases of the bijection above separately.

Suppose first that $H^\ominus \cap \{x, x - d\} = \emptyset$. Clearly $\langle X^1 \ominus U^1, H^\ominus \rangle = 0$, and as $H^\ominus \subseteq U^\ominus$, we have $\langle U_{\text{sp}}^1 \ominus X_{\text{sp}}^1, H^\ominus \rangle = \langle \{0, \dots, x - d\} \ominus \{0, \dots, x\}, H^\ominus \rangle$ as in Lemma 3.2. Now

$$\begin{aligned} \langle U_{\text{sp}}^1, H^\ominus \ominus G^\ominus \rangle &= \langle U_{\text{sp}}^1, \{x - d\} \rangle = |U^\ominus| + \langle U^\ominus, \{0, \dots, x - d\} \rangle \\ &= \text{def}(V) + \langle X^\ominus \ominus \{x, x - d\}, \{0, \dots, x - d\} \rangle \end{aligned}$$

because $x \notin U^\cup$. Summing the above terms it suffices to show that

$$(-1)^{\delta(\lambda)} = (-1)^{1+(U^1, H^\ominus \ominus G^\ominus)} = (-1)^{1+(U^1, \{x-d\})},$$

and this is straightforward.

Finally we assume that $H^\ominus \cap \{x, x - d\} = \{x\}$. This time $H^\ominus \ominus \{x\} \subseteq X^\ominus \cap U^\ominus$ so

$$\langle U_{\text{sp}}^1 \ominus X_{\text{sp}}^1, H^\ominus \ominus \{x\} \rangle = \langle \{0, \dots, x - d\} \ominus \{0, \dots, x\}, H^\ominus \ominus \{x\} \rangle.$$

Moreover, $\langle X_{\text{sp}}^1 \ominus U_{\text{sp}}^1, \{x\} \rangle = \langle X_{\text{sp}}^1, \{x\} \rangle = \overline{\text{def}(V)} + \langle X^\ominus, \{0, \dots, x\} \rangle$ and $\langle U^1, H^\ominus \ominus G^\ominus \rangle = \langle U_{\text{sp}}^1, H^\ominus \ominus G^\ominus \rangle = 0$ because $H^\ominus \ominus G^\ominus = \{x\}$ and $x \notin U^\cup$. As before it suffices

to show that

$$(-1)^{\delta(\lambda)} = (-1)^{1+(X^1 \ominus U^1, H^\ominus)} = (-1)^{1+(X^1 \ominus U^1, \{x\})},$$

and again this is straightforward. □

3.2 More symbols

We have a linear map $[-] : \mathbb{C}[\tilde{\mathcal{P}}] \rightarrow \mathbb{C}[\mathcal{S}]$ defined such that if $X \in \tilde{\mathcal{P}}$ is nondegenerate then $\llbracket X \rrbracket \in \mathcal{S}$ and if X is degenerate then $\llbracket X \rrbracket = \llbracket X \rrbracket_+ + \llbracket X \rrbracket_-$. This clearly factors through a map $\mathbb{C}[\tilde{\mathcal{S}}] \rightarrow \mathbb{C}[\mathcal{S}]$. The image of $[-]$ has a natural complement in $\mathbb{C}[\mathcal{S}]$, namely $\langle \llbracket X \rrbracket_+ - \llbracket X \rrbracket_- \mid X \in \tilde{\mathcal{P}} \text{ is degenerate} \rangle_{\mathbb{C}}$.

We wish to understand to what extent the endomorphisms $\tilde{\mathcal{R}}$ and $\tilde{\mathcal{H}}_{d,i}^j$ of $\mathbb{C}[\tilde{\mathcal{S}}]$ factor through $[-]$. As the term arises frequently we let

$$d(X) = (\text{def}(X) - \text{def}(X_{\text{sp}}))/2 \in \mathbb{N}_0$$

for any $X \in \tilde{\mathcal{P}}$. We then define a \mathbb{C} -linear map $\varepsilon : \mathbb{C}[\tilde{\mathcal{S}}] \rightarrow \mathbb{C}[\tilde{\mathcal{S}}]$ by setting $\varepsilon(\llbracket X \rrbracket) = (-1)^{d(X)} \llbracket X \rrbracket$. The following easy observations are stated in [34, §2].

Lemma 3.3 *For any $(d, i) \in \mathbb{Z}^{(2)}$ and $j \in \{0, 1\}$ we have the following equalities of linear endomorphisms of $\mathbb{C}[\tilde{\mathcal{S}}]$:*

- (i) $\tilde{\mathcal{R}} \circ (-)^{\text{op}} = \varepsilon \circ \tilde{\mathcal{R}}$,
- (ii) $(-)^{\text{op}} \circ \tilde{\mathcal{R}} = \tilde{\mathcal{R}} \circ \varepsilon$,
- (iii) $(-)^{\text{op}} \circ \tilde{\mathcal{H}}_{d,i}^j = (-1)^j \tilde{\mathcal{H}}_{d,i}^j \circ (-)^{\text{op}}$,
- (iv) $\varepsilon \circ (-)^{\text{op}} = \Theta \circ (-)^{\text{op}} \circ \varepsilon$.

Proof (i). If $X \in \tilde{\mathcal{P}}$ then $\text{Sim}(X) = \text{Sim}(X^{\text{op}}) = \text{Sim}(X)^{\text{op}}$. As $(X^{\text{op}})^\sharp = X^\ominus \ominus X^\sharp$, we see that the coefficient of $Y \in \text{Sim}(\Lambda)$ in $\tilde{\mathcal{R}}(X^{\text{op}})$ is $(-1)^{\langle X^\ominus, Y^\sharp \rangle} = (-1)^{\langle X^\ominus, Y^1 \rangle + \langle X^\ominus, X_{\text{sp}}^1 \rangle}$ times the corresponding coefficient in $\tilde{\mathcal{R}}(X)$. Now,

$$\begin{aligned} 2d(X) &= (|X^\ominus| - 2|Y^1 \cap X^\ominus|) - (|X^\ominus| - 2|X_{\text{sp}}^1 \cap X^\ominus|) \\ &= 2(|X_{\text{sp}}^1 \cap X^\ominus| - |Y^1 \cap X^\ominus|). \end{aligned}$$

(ii). This is similar to (i) using that $\langle X^\sharp, Y^\sharp \ominus (Y^{\text{op}})^\sharp \rangle = \langle X^\sharp, X^\ominus \rangle$.

(iii). It is straightforward to check that for any $X \in \tilde{\mathcal{P}}$, we have $\mathcal{H}_{d,i}(X^{\text{op}}) = \mathcal{H}_{d,i}(X)^{\text{op}}$ and $l_{d,i}(\lambda, X) = l_{d,i}(\lambda^{\text{op}}, X^{\text{op}})$, which gives the statement.

(iv). As $\text{def}(X^{\text{op}}) = -\text{def}(X)$, we have $d(X^{\text{op}}) = d(X) - \text{def}(X)$. □

We define for each integer $e \in \mathbb{Z}$ the set

$$\tilde{\mathcal{P}}^{\equiv e} = \{X \in \tilde{\mathcal{P}} \mid d(X) \equiv e \pmod{2}\}.$$

This gives a partition $\tilde{\mathcal{P}} = \tilde{\mathcal{P}}^{\equiv 0} \sqcup \tilde{\mathcal{P}}^{\equiv 1}$ and partitions $\tilde{\mathcal{P}}^{\text{od}} = \tilde{\mathcal{P}}^{\text{od},0} \sqcup \tilde{\mathcal{P}}^{\text{od},1}$ and $\tilde{\mathcal{P}}^{\text{ev}} = \tilde{\mathcal{P}}^{\text{ev},0} \sqcup \tilde{\mathcal{P}}^{\text{ev},1}$. Note that $(-)^{\text{op}}$ swaps $\tilde{\mathcal{P}}^{\text{od},0}$ and $\tilde{\mathcal{P}}^{\text{od},1}$ but stabilises $\tilde{\mathcal{P}}^{\text{ev},0}$ and $\tilde{\mathcal{P}}^{\text{ev},1}$. Thus we get a partition $\mathcal{S}^{\text{ev}} = \mathcal{S}^{\text{ev},0} \sqcup \mathcal{S}^{\text{ev},1}$, where $\mathcal{S}^{\text{ev},0}$ contains all $\llbracket X \rrbracket_\pm$ with $X \in \tilde{\mathcal{P}}^{\text{ev},0}$ degenerate.

If $X \in \tilde{\mathcal{P}}$ then we let $\text{Sim}_e(X) = \text{Sim}(X) \cap \tilde{\mathcal{P}}^{\equiv e}$. Under the map $^\sharp : \tilde{\mathcal{P}} \rightarrow \text{Pow}(\mathbb{N}_0)$, we have $X \in \tilde{\mathcal{P}}^{\equiv e}$ if and only if $|X^\sharp| \equiv e \pmod{2}$. Now we define $\tilde{\mathcal{R}}_e : \mathbb{C}[\tilde{\mathcal{P}}] \rightarrow \mathbb{C}[\tilde{\mathcal{P}}]$ by setting

$$\tilde{\mathcal{R}}_e(X) = \frac{1}{s(X)} \sum_{Y \in \text{Sim}_e(X)} (-1)^{\langle X^\sharp, Y^\sharp \rangle} Y$$

so that $\tilde{\mathcal{R}} = \tilde{\mathcal{R}}_0 + \tilde{\mathcal{R}}_1$.

Lemma 3.4 *For any $X \in \tilde{\mathcal{P}}$ and $e \in \mathbb{Z}$ we have:*

- (i) $\tilde{\mathcal{R}}_e(X^{op}) = (-1)^e \tilde{\mathcal{R}}_e(X)$,
- (ii) $\tilde{\mathcal{R}}_e(X)^{op} = (-1)^{d(X)} \tilde{\mathcal{R}}_{e+\text{def}(X)}(X)$

In particular, for any $X \in \tilde{\mathcal{P}}^{\equiv 1}$ we have $\llbracket \tilde{\mathcal{R}}_e(X) \rrbracket = 0$.

Proof This follows by projecting (i) and (ii) of Lemma 3.3 onto each summand of the decomposition $\mathbb{C}[\tilde{\mathcal{P}}] = \mathbb{C}[\tilde{\mathcal{P}}^{\equiv 0}] \oplus \mathbb{C}[\tilde{\mathcal{P}}^{\equiv 1}]$. Here we use that $\text{Sim}_e(X)^{op} = \text{Sim}_{e+\text{def}(X)}(X)$, which follows from the above remarks. For the final statement note that (ii) shows that if $X \in \tilde{\mathcal{P}}^{\equiv 1}$ then $\llbracket \tilde{\mathcal{R}}_e(X) \rrbracket = -\llbracket \tilde{\mathcal{R}}_{e+\text{def}(X)}(X) \rrbracket = -\llbracket \tilde{\mathcal{R}}_e(X) \rrbracket$. \square

By (i) of Lemma 3.4 we have \mathcal{R}_0 factors through $\llbracket - \rrbracket$ to give an endomorphism of its image. We extend this to an endomorphism of $\mathbb{C}[\mathcal{S}]$ by letting it fix pointwise the complement defined above (in other words, $\mathcal{R}_0(\llbracket X \rrbracket_{\pm}) = \llbracket X \rrbracket_{\pm}$ for all degenerate $X \in \tilde{\mathcal{P}}$). We also denote this by \mathcal{R}_0 . We consider the subspaces $\mathbb{C}[\mathcal{S}^{od}]$ and $\mathbb{C}[\mathcal{S}^{ev}]$ separately. First let $\mathcal{A}^{od} = \mathcal{U}^{od} = \mathbb{C}[\mathcal{S}^{od}]$.

Note that $\mathcal{R}_0(\llbracket X \rrbracket)$ is simply the Fourier transform on the abelian group $\text{Pow}_0(X^{\ominus})$. As such \mathcal{R}_0^2 is the identity on $\mathbb{C}[\mathcal{S}^{od}]$. To have a compatible notation we consider \mathcal{R}_0 as a map $\mathcal{U}^{od} \rightarrow \mathcal{A}^{od}$ and denote by $\mathcal{Q}_0 : \mathcal{A}^{od} \rightarrow \mathcal{U}^{od}$ its inverse.

Assume $(d, i) \in \mathbb{Z}^{(2)}$ with $d \neq 0$. It follows from Lemma 3.3 that the restriction of the map $\tilde{\mathcal{H}}_{d,i}^0$ to $\mathbb{C}[\tilde{\mathcal{S}}^{od}]$ factors through a well-defined endomorphism of \mathcal{U}^{od} and \mathcal{A}^{od} which we denote by $\mathcal{H}_{d,i}^0$. Similarly $\tilde{\mathcal{H}}_{d,i}^1 \circ \varepsilon$ factors through an endomorphism which we denote by $\mathcal{H}_{d,i}^1$. The following is now simply a consequence of Theorem 3.1.

Proposition 3.5 *For any $(d, i) \in \mathbb{Z}^{(2)}$, with $d \neq 0$, we have commutative diagrams*

$$\begin{array}{ccc}
 \mathcal{A}^{od} & \xrightarrow{\mathcal{Q}_0} & \mathcal{U}^{od} & & \mathcal{U}^{od} & \xrightarrow{\mathcal{R}_0} & \mathcal{A}^{od} \\
 \mathcal{H}_{d,i}^0 \downarrow & & \downarrow \mathcal{H}_{d,i}^0 & & \mathcal{H}_{d,i}^0 \downarrow & & \downarrow \mathcal{H}_{d,i}^0 \\
 \mathcal{A}^{od} & \xrightarrow{\mathcal{Q}_0} & \mathcal{U}^{od} & & \mathcal{U}^{od} & \xrightarrow{\mathcal{R}_0} & \mathcal{A}^{od}
 \end{array}$$

If $X \in \tilde{\mathcal{P}}$ has even defect then $\text{Sim}_0(X) = \text{Sim}_0(X)^{op}$. As remarked in the proof of Lemma 3.3 we have $(Y^{op})^{\sharp} = Y^{\sharp} \ominus X^{\ominus}$ so $\langle X^{\sharp}, (Y^{op})^{\sharp} \rangle = \langle X^{\sharp}, Y^{\sharp} \rangle$ for any $Y \in \text{Sim}_0(X)$. Thus if $X \in \tilde{\mathcal{P}}^{ev}$ is nondegenerate then

$$\mathcal{R}_0(\llbracket X \rrbracket) = \frac{2}{s(X)} \sum_{Y \in \text{Sim}_0(X)} (-1)^{\langle X^{\sharp}, Y^{\sharp} \rangle} \llbracket Y \rrbracket$$

where $\overline{\text{Sim}}_e(X) = \{Y, Y^{op} \mid Y \in \text{Sim}_e(X)\}$. This is the Fourier transform on the abelian group $\overline{\text{Pow}}_0(X^{\ominus}) = \text{Pow}_0(X^{\ominus}) / \{\emptyset, X^{\ominus}\}$.

If $e \in 2\mathbb{Z}$ is an even integer then we let $\mathcal{A}^{ev,e} = \mathcal{U}^{ev,e} = \mathbb{C}[\mathcal{S}^{ev,0}] \subseteq \mathbb{C}[\mathcal{S}^{ev}]$. The map \mathcal{R}_0 restricts to an involution on the subspace $\mathbb{C}[\mathcal{S}^{ev,0}]$. As above we consider \mathcal{R}_0 as a map $\mathcal{A}^{ev,e} \rightarrow \mathcal{U}^{ev,e}$ with inverse $\mathcal{Q}_0 : \mathcal{U}^{ev,e} \rightarrow \mathcal{A}^{ev,e}$.

Let $\tilde{\mathcal{P}}^{ev,nd} \subseteq \tilde{\mathcal{P}}^{ev,0}$ be the subset of nondegenerate arrays. For any odd integer $e \in 2\mathbb{Z} + 1$ we let $\mathcal{U}^{ev,e} = \mathbb{C}[\mathcal{S}^{ev,1}]$ and define the quotient space

$$\mathcal{A}^{ev,e} = \mathbb{C}[\tilde{\mathcal{P}}^{ev,nd}] / \langle \llbracket X \rrbracket + \llbracket X^{op} \rrbracket \mid X \in \tilde{\mathcal{P}}^{ev,nd} \rangle_{\mathbb{C}}.$$

By Lemma 3.4 the map $\tilde{\mathcal{R}}_e$ factors through a map $\mathcal{A}^{ev,e} \rightarrow \mathcal{Q}^{ev,e}$ which we denote by \mathcal{R}_e . We define a right inverse $\mathcal{Q}_e : \mathcal{Q}^{ev,e} \rightarrow \mathcal{A}^{ev,e}$ of this map by setting

$$\mathcal{Q}_e(\llbracket X \rrbracket) = \frac{1}{s(X)} \sum_{Y \in \text{Sim}_0(X)} (-1)^{\langle X^\sharp, Y^\sharp \rangle} \langle\langle Y \rangle\rangle = \frac{2}{s(X)} \sum_{\bar{Y} \in \overline{\text{Sim}}_0(X)} (-1)^{\langle X^\sharp, Y^\sharp \rangle} \langle\langle Y \rangle\rangle$$

where $\langle\langle - \rangle\rangle : \mathbb{C}[\tilde{\mathcal{P}}^{ev,nd}] \rightarrow \mathcal{A}^{ev,e}$ is the natural quotient map.

It follows easily from Lemma 3.3 that the endomorphism $\tilde{\mathcal{H}}_{d,0}^i$ of $\mathbb{C}[\tilde{\mathcal{P}}^{ev,nd}]$ factors through a well defined map $\mathcal{H}_{d,0}^i : \mathcal{A}^{ev,e} \rightarrow \mathcal{A}^{ev,e+i}$ for each $e \in \mathbb{Z}$. Similarly we have $\mathcal{H}_{d,i}^0$ factors through a map $\mathcal{Q}^{ev,e} \rightarrow \mathcal{Q}^{ev,e+i}$.

Proposition 3.6 *For any $(d, i) \in \mathbb{Z}^{(2)}$, with $d \neq 0$, and any $e \in \mathbb{Z}$ we have commutative diagrams*

$$\begin{array}{ccc} \mathcal{Q}^{ev,e} & \xrightarrow{\mathcal{Q}_e} & \mathcal{A}^{ev,e} & & \mathcal{A}^{ev,e} & \xrightarrow{\mathcal{R}_e} & \mathcal{Q}^{ev,e} \\ (-1)^i \mathcal{H}_{d,i}^0 \downarrow & & \downarrow \mathcal{H}_{d,0}^i & & \mathcal{H}_{d,0}^i \downarrow & & \downarrow (-1)^i \mathcal{H}_{d,i}^0 \\ \mathcal{Q}^{ev,e+i} & \xrightarrow{\mathcal{Q}_{e+i}} & \mathcal{A}^{ev,e+i} & & \mathcal{A}^{ev,e+i} & \xrightarrow{\mathcal{R}_{e+i}} & \mathcal{Q}^{ev,e+i} \end{array}$$

4 Hyperoctahedral groups

Assume $(\mathcal{I}, <)$ is a finite totally ordered set of cardinality $|\mathcal{I}| = 2n$. Denote by $\dagger : \mathcal{I} \rightarrow \mathcal{I}$ the unique order reversing bijection on \mathcal{I} . We say $a \in \mathcal{I}$ is positive or negative if $a > a^\dagger$ or $a < a^\dagger$ respectively. This gives a decomposition $\mathcal{I} = \mathcal{I}^+ \sqcup \mathcal{I}^-$ into subsets of cardinality n . If $\mathcal{O} \subseteq \mathcal{I}$ is a subset then $(\mathcal{O}, <)$ is also a totally ordered set.

Example 4.1 We could take $\mathcal{I} = \{-n < \dots < -1 < 1 < \dots < n\}$ then for any $a \in \mathcal{I}$ we have $a^\dagger = -a$ so $\mathcal{I}^+ = \{1, \dots, n\}$ and $\mathcal{I}^- = \{-1, \dots, -n\}$.

If $\mathfrak{S}_{\mathcal{I}}$ is the symmetric group on \mathcal{I} then we define $W_{\mathcal{I}} = \mathbf{C}_{\mathfrak{S}_{\mathcal{I}}}(\sigma)$ to be the centraliser of the involution $\sigma = \prod_{a \in \mathcal{I}^+} (a, a^\dagger)$. Let $\delta_{\mathcal{I}} : \mathfrak{S}_{\mathcal{I}} \rightarrow \mathbb{Z}/2\mathbb{Z}$ be the unique non-trivial homomorphism. Given $e \in \{0, 1\}$ we let $W_{\mathcal{I}}^e = \{w \in W_{\mathcal{I}} \mid \delta_{\mathcal{I}}(w) = e\}$ so that we have a decomposition $W_{\mathcal{I}} = W_{\mathcal{I}}^0 \sqcup W_{\mathcal{I}}^1$ into the cosets of $W_{\mathcal{I}}^0 \triangleleft W_{\mathcal{I}}$. Note we have a semidirect product decomposition $W_{\mathcal{I}} = N_{\mathcal{I}} \rtimes H_{\mathcal{I}}$ where $N_{\mathcal{I}} = \langle (a, a^\dagger) \mid a \in \mathcal{I}^+ \rangle$ and $H_{\mathcal{I}} = \{w \in W_{\mathcal{I}} \mid {}^w \mathcal{I}^+ = \mathcal{I}^+ \} \cong \mathfrak{S}_{\mathcal{I}^+}$.

For any σ -stable subset $\mathcal{O} \subseteq \mathcal{I}$, equivalently $\mathcal{O} = \mathcal{O}^\dagger$, we have a natural injective homomorphism $W_{\mathcal{O}} \rightarrow W_{\mathcal{I}}$ whose image is the pointwise stabiliser of $\mathcal{I} \setminus \mathcal{O}$. We identify $W_{\mathcal{O}}$ with its image in $W_{\mathcal{I}}$.

We say $w \in W_{\mathcal{I}}$ is an \mathcal{I} -cycle if the subgroup $\langle w, \sigma \rangle \leq W_{\mathcal{I}}$ acts transitively on \mathcal{I} . Thus $w = nh$, with $n \in N_{\mathcal{I}}$ and $h \in H_{\mathcal{I}}$ acting on \mathcal{I}^+ as cycle of length n . The following is an elementary calculation.

Lemma 4.2 *If $w \in W_{\mathcal{I}}$ is an \mathcal{I} -cycle then $\mathbf{C}_{W_{\mathcal{I}}}(w) = \langle w \rangle$ if $\delta_{\mathcal{I}}(w) = 1$ and $\mathbf{C}_{W_{\mathcal{I}}}(w) = \langle w \rangle \rtimes \langle \sigma \rangle$ if $\delta_{\mathcal{I}}(w) = 0$. In either case $|\mathbf{C}_{W_{\mathcal{I}}}(w)| = |\mathcal{I}| = 2n$.*

Now suppose $w \in W_{\mathcal{I}}$ and $\mathcal{I}/\langle w, \sigma \rangle = \{\mathcal{O}_1, \dots, \mathcal{O}_k\}$. Then we can write $w = w_1 \cdots w_k$ as a pairwise commuting product with $w_i \in W_{\mathcal{O}_i}$ an \mathcal{O}_i -cycle. Such a decomposition, which we call a *cycle decomposition*, is unique up to reordering. We call $k \geq 1$ the *cycle length*.

There is a unique ordering of the orbits such that $(-1)^{\delta_{\mathcal{O}_1}(w_1)}|\mathcal{O}_1^+| \geq \dots \geq (-1)^{\delta_{\mathcal{O}_k}(w_k)}|\mathcal{O}_k^+|$. With this ordering we call $((-1)^{\delta_{\mathcal{O}_1}(w_1)}|\mathcal{O}_1^+|, \dots, (-1)^{\delta_{\mathcal{O}_k}(w_k)}|\mathcal{O}_k^+|)$ the signed cycle type of the element. It determines the conjugacy class $\text{Cl}_{W_{\mathcal{I}}}(w)$ uniquely. If these inequalities are all strict then we say w has *pairwise distinct cycles*.

Lemma 4.3 *Let $w \in W_{\mathcal{I}}$ be as above and let $\mathcal{I} = \mathcal{I}_1 \sqcup \dots \sqcup \mathcal{I}_m$ be a decomposition into σ -stable sets such that $P = W_{\mathcal{I}_1} \cdots W_{\mathcal{I}_m} \leq W_{\mathcal{I}}$ contains w . If w has pairwise distinct cycles then the following hold:*

- (i) $C_{W_{\mathcal{I}}}(w) = C_{W_{\mathcal{O}_1}}(w_1) \cdots C_{W_{\mathcal{O}_k}}(w_k)$,
- (ii) $|C_{W_{\mathcal{I}}}(w)| \leq 2^k \cdot n^k$,
- (iii) $C_P(w) = C_W(w)$.

Proof Part (i) is given by the uniqueness of the cycle decomposition. Part (ii) follows from (i) and Lemma 4.2. Part (iii) follows from (i) because $\langle w, \sigma \rangle$ stabilises each \mathcal{I}_j , so they must be a union of the \mathcal{O}_i . □

Recall from Sect. 2 that we have define the β -sets \mathcal{B}_n . After [14, §6.4.1] we have a bijection $\mathcal{B}_n \rightarrow \text{Irr}(\mathfrak{S}_{\mathcal{I}^+})$ which we denote by $[A] \mapsto \chi_{[A]}$. Under the natural isomorphism $\mathfrak{S}_{\mathcal{I}^+} \cong H_{\mathcal{I}^+}$ we get a bijection $\mathcal{B}_n \rightarrow \text{Irr}(H_{\mathcal{I}})$.

If $\delta \in \{0, 1\}$ then this yields a bijection $\tilde{\mathcal{S}}_n^\delta \rightarrow \text{Irr}(W_{\mathcal{I}})$, denoted by $[X] \mapsto \rho_{[X]}$, defined as follows. First note that for any $[X] \in \tilde{\mathcal{S}}_n^\delta$ we have $\text{rk}([X^0]) + \text{rk}([X^1]) = n$ by Eq. (2.1). Now choose a σ -stable partition $\mathcal{I} = \mathcal{I}_0 \sqcup \mathcal{I}_1$ such that $|\mathcal{I}_j| = 2 \cdot \text{rk}(X^j)$ with $j \in \{0, 1\}$ (note these subsets may be empty). We then have

$$\rho_{[X]} = \text{Ind}_{W_{\mathcal{I}_0} W_{\mathcal{I}_1}}^{W_{\mathcal{I}}} (\tilde{\chi}_{[X^0]} \boxtimes (\varepsilon_{\mathcal{I}_1} \tilde{\chi}_{[X^1]}))$$

where $\tilde{\chi}_{[X^j]}$ is the inflation of $\chi_{[X^j]} \in \text{Irr}(H_{\mathcal{I}_j})$ under the map $W_{\mathcal{I}_j} \rightarrow H_{\mathcal{I}_j}$. These characters satisfy the following MN-rule (or Murnaghan–Nakayama rule).

Proposition 4.4 *Let $\mathcal{O} \in \mathcal{I}/\langle w, N_{\mathcal{I}} \rangle$ be an orbit for some element $w \in W_{\mathcal{I}}$. Then we have a unique decomposition $w = w_1 w_2 = w_2 w_1$ with $w_1 \in W_{\mathcal{O}}$ and $w_2 \in W_{\mathcal{I} \setminus \mathcal{O}}$. If $(d, j) = (|\mathcal{O}^+|, \delta_{\mathcal{O}}(w_1))$ then for any $[X] \in \tilde{\mathcal{S}}_n^\delta$, with $\delta \in \{0, 1\}$, we have*

$$\rho_{[X]}(w) = \sum_{\lambda \in \mathcal{H}_{d,0}(X)} (-1)^{j\delta(\lambda) + I_{d,0}(\lambda, X)} \rho_{[X \setminus \lambda]}(w_2),$$

Proof We refer the reader to [14, Thm. 10.3.1]. For the correspondence between hooks of partitions and hooks of β -sets see [31, §I.1]. □

In [22, Theorem 7.2], a bound is given for the character values of the symmetric group at a given element in terms of its cycle length. The argument in [22] is a consequence of the MN-rule together with analogues of the following easy observations.

Lemma 4.5 *Let $X \in \tilde{\mathcal{P}}$ be an array with an (e, j) -hook $\lambda \in \mathcal{H}_{e,j}(X)$ for some $(e, j) \in \mathbb{N}_0^{(2)}$. Then for any $(d, i) \in \mathbb{N}_0^{(2)}$ we have*

$$\mathcal{H}_{d,i}(X) \subseteq \mathcal{H}_{d,i}(X \setminus_{e,j} \lambda) \cup \{\lambda, \mathcal{D}_{e-d,i+j}(\lambda)\}.$$

Proof Let $\mu \in \mathcal{H}_{d,i}(X)$ so $\mathcal{D}_{d,i}(\mu) \in \mathbb{N}_0^{(2)}$. If $\mu \neq \lambda$ then $\mu \in Y := X \setminus_{e,j} \lambda$, and if μ is not a (d, i) -hook of Y then clearly $\mathcal{D}_{d,i}(\mu) \in Y - X = \{\mathcal{D}_{e,j}(\lambda)\}$. □

Lemma 4.6 *If $(n, i) \in \mathbb{N}_0^{(2)}$ then $X \in \tilde{\mathcal{P}}_n$ has at most one (n, i) -hook.*

Proof Suppose $\lambda \in \mathcal{H}_{n,i}(X)$ is such a hook. Then $X \setminus_{n,i} \lambda$ is of rank 0 and so has no hooks. Hence, by the previous lemma the only possible (n, i) -hooks are λ and λ^{op} . But it is easily seen that if λ^{op} were a hook of X then it would also be one of $X \setminus_{n,i} \lambda$, which is impossible. \square

Theorem 4.7 *Fix an integer $1 \leq k \leq n$. Then for each element $w \in W_{\mathcal{I}}$ of cycle length k and each irreducible character $\chi \in \text{Irr}(W_{\mathcal{I}})$ we have*

$$|\chi(w)| \leq 2^{k-1} \cdot k!$$

Moreover, if $w \in W_{\mathcal{I}}^0$ then for all $\chi \in \text{Irr}(W_{\mathcal{I}}^0)$ we have $|\chi(w)| \leq (2^k + 1) \cdot 2^{k-1} \cdot k! \leq 2^{2k} \cdot k!$.

Proof Let $\chi = \rho_{[X]}$ with $[X] \in \tilde{\mathcal{S}}_n^1$. We argue by induction on k . Suppose $k = 1$. If $\chi(w) \neq 0$ then by Proposition 4.4, $[X]$ has an $(n, 0)$ -hook, but by Lemma 4.6, there is at most one such hook, so $|\chi(w)| \leq 1$. So assume $k > 1$. We have $\mathcal{I}/\langle w, \sigma \rangle = \{\mathcal{O}_1, \dots, \mathcal{O}_k\}$ and we set $d_i = |\mathcal{O}_i|$ for any $1 \leq i \leq k$. We also let $w = w_1 w_2$ with $w_1 \in W_{\mathcal{O}_1}$ and $w_2 \in W_{\mathcal{I} \setminus \mathcal{O}_1}$.

Clearly we may assume that $\chi(w) \neq 0$. By Proposition 4.4 and the induction hypothesis we see that

$$|\chi(w)| \leq \sum_{\lambda \in \mathcal{H}_{d_1,0}(X)} |\rho_{[X \setminus \lambda]}(w_2)| \leq |\mathcal{H}_{d_1,0}(X)| \cdot 2^{k-2} \cdot (k-1)!$$

So it suffices to show that $|\mathcal{H}_{d_1,0}(X)| \leq 2k$.

Repeatedly applying Proposition 4.4, we see that there exist arrays $X = X_0, X_1, \dots, X_k \in \tilde{\mathcal{P}}$ such that for $1 \leq i \leq k$ we have $X_i = X_{i-1} \setminus_{d_i,0} \lambda_i$ for some hook $\lambda_i \in \mathcal{H}_{d_i,0}(X_{i-1})$. As

$$\text{rk}(X_i) = \text{rk}(X_0) - (d_1 + \dots + d_i),$$

we have $\text{rk}(X_k) = 0$, so $\mathcal{H}_{d_1,0}(X_k) = \emptyset$. By Lemma 4.5 we have $|\mathcal{H}_{d_1,0}(X_i)| \leq |\mathcal{H}_{d_1,0}(X_{i+1})| + 2$, which yields the desired bound.

Now assume $\chi \in \text{Irr}(W_{\mathcal{I}}^0)$. If χ extends to $W_{\mathcal{I}}$ then we are done so assume this is not the case. Then $\chi = \text{Res}_{W_{\mathcal{I}}^0}^{W_{\mathcal{I}}}(\rho_{[X]})$ for some degenerate symbol $[X] \in \tilde{\mathcal{S}}_n^0$ and is the sum $\chi_+ + \chi_-$ of two distinct irreducible characters. Clearly $\chi_{\pm}(w) = \frac{1}{2}(\chi(w) + \Delta(w))$ where $\Delta = \chi_+ - \chi_-$ is the difference character.

If $\Delta(w) = 0$ then we are done, so we may assume that $\Delta(w) \neq 0$. A result of Stembridge [33, Theorem 7.5] shows that, in this case, $|\Delta(w)| = 2^k |\chi_{[A]}(x)|$ for some character $\chi_{[A]} \in \text{Irr}(\mathfrak{S}_{\mathcal{I}^+})$ and some element $x \in \mathfrak{S}_{\mathcal{I}^+}$ which is a product of k disjoint cycles. Hence, by [22, Theorem 7.2], we have $|\Delta(w)| \leq 2^{2k-1} \cdot k!$, which easily gives the bound. \square

For an ordered symbol $[X] \in \tilde{\mathcal{S}}_n$ of rank n we define a corresponding class function

$$\phi_{[X]} = \frac{1}{s(X)} \sum_{\substack{Y \in \text{Sim}(X) \\ \text{def}(Y) = \text{def}(Y_{\text{sp}})}} (-1)^{\langle X^{\sharp}, Y^{\sharp} \rangle} \rho_{[Y]} \in \text{Class}(W_{\mathcal{I}}).$$

Note that $\{Y \in \text{Sim}(X) \mid \text{def}(Y) = \text{def}(Y_{\text{sp}})\} \subseteq \text{Sim}_0(X)$ so this is essentially the projection of the Fourier transform $\tilde{\mathcal{R}}_0(X)$ onto the subspace $\mathbb{Q}[\tilde{\mathcal{S}}_n^{\delta}]$, where $\delta = \text{def}(X_{\text{sp}})$. Somewhat remarkably these functions also satisfy a version of the MN-rule, which is the main point of Theorem 3.1.

Theorem 4.8 Let $\mathcal{O} \in \mathcal{I}/\langle w, N_{\mathcal{I}} \rangle$ be an orbit for some element $w \in W_{\mathcal{I}}$. Then we have a unique decomposition $w = w_1 w_2 = w_2 w_1$ with $w_1 \in W_{\mathcal{O}}$ and $w_2 \in W_{\mathcal{I} \setminus \mathcal{O}}$. If $(d, j) = (|\mathcal{O}^+|, \delta_{\mathcal{O}}(w_1))$ then for any $[X] \in \tilde{\mathcal{S}}_n$, we have

$$\phi_{[X]}(w) = (-1)^{j(1+\text{def}(X))} \sum_{\lambda \in \mathcal{H}_{d,j}(X)} (-1)^{l_{d,j}(\lambda, X)} \phi_{[X \setminus \lambda]}(w_2).$$

Proof By Proposition 4.4 we have

$$\phi_{[X]}(w) = \frac{1}{s(X)} \sum_{\substack{Y \in \text{Sim}(X) \\ \text{def}(Y) = \text{def}(X_{\text{sp}})}} \sum_{\lambda \in \mathcal{H}_{d,0}(Y)} (-1)^{j\delta(\lambda) + (X^{\sharp}, Y^{\sharp}) + l_{d,0}(\lambda, X)} \rho_{[Y \setminus \lambda]}(w_2).$$

Let $\delta = \text{def}(X_{\text{sp}}) \in \{0, 1\}$. Under the isomorphism $\mathbb{Q}[\text{Irr}(W_{\mathcal{I} \setminus \mathcal{O}})] \xrightarrow{\sim} \mathbb{Q}[\tilde{\mathcal{S}}_{n-d}^{\delta}]$ defined above the right hand side is identified with an expression in $\mathbb{Q}[\tilde{\mathcal{S}}]$.

Under the decomposition $\mathbb{Q}[\tilde{\mathcal{S}}] = \bigoplus_{e \in \mathbb{Z}} \mathbb{Q}[\tilde{\mathcal{S}}^e]$ this is the projection of $\tilde{\mathcal{H}}_{d,0}^j(\tilde{\mathcal{R}}([X]))$ onto the subspace $\mathbb{Q}[\tilde{\mathcal{S}}^{\delta}]$. If $\text{def}(X)$ is odd then by Theorem 3.1 we have $\tilde{\mathcal{H}}_{d,0}^j(\tilde{\mathcal{R}}([X])) = \tilde{\mathcal{R}}(\tilde{\mathcal{H}}_{d,j}^0([X]))$ and projecting the right hand side of this onto $\mathbb{Q}[\tilde{\mathcal{S}}^{\delta}]$ gives us that

$$\begin{aligned} \phi_{[X]}(w) &= \sum_{\lambda \in \mathcal{H}_{d,\delta}(X)} \frac{1}{s(X \setminus \lambda)} \sum_{\substack{Y \in \text{Sim}(X \setminus \lambda) \\ \text{def}(Y) = \text{def}(Y_{\text{sp}})}} (-1)^{((X \setminus \lambda)^{\sharp}, Y^{\sharp}) + l_{d,j}(\lambda, X)} \rho_{[Y]}(w_2) \\ &= \sum_{\lambda \in \mathcal{H}_{d,\delta}(X)} (-1)^{l_{d,j}(\lambda, X)} \phi_{[X \setminus \lambda]}(w_2). \end{aligned}$$

If $\text{def}(X)$ is even then the same holds, but we must multiply through by $(-1)^j$. □

Theorem 4.9 Fix an integer $1 \leq k \leq n$. Then for each element $w \in W_{\mathcal{I}}$ of cycle length k and each symbol $[X] \in \mathcal{S}_n$ of rank n , we have

$$|\phi_{[X]}(w)| \leq 2^{k-1} \cdot k!.$$

Proof The proof is identical to that of Theorem 4.7. □

We now associate functions to unordered symbols as follows. If $\llbracket X \rrbracket \in \mathcal{S}_n^{\text{od}}$ has odd defect then we simply let $\phi_{\llbracket X \rrbracket} = \phi_{[X]} \in \text{Class}(W_{\mathcal{I}})$. Now fix $e \in \{0, 1\}$. Then for any $\llbracket X \rrbracket \in \mathcal{S}_n^{\text{ev},e}$ we let

$$\phi_{\llbracket X \rrbracket} = \text{Res}_{W_{\mathcal{I}}^e}^{W_{\mathcal{I}}}(\phi_{[X]}) = \frac{1}{s(X)} \sum_{\substack{Y \in \text{Sim}(X) \\ \text{def}(Y) = 0}} (-1)^{\langle X^{\sharp}, Y^{\sharp} \rangle} \text{Res}_{W_{\mathcal{I}}^e}^{W_{\mathcal{I}}}(\rho_{[Y]}) \in \text{Class}(W_{\mathcal{I}}^e).$$

Note that $\text{Res}_{W_{\mathcal{I}}^e}^{W_{\mathcal{I}}}(\rho_{[Y^{\text{op}}]}) = (-1)^e \text{Res}_{W_{\mathcal{I}}^e}^{W_{\mathcal{I}}}(\rho_{[Y]})$ so these functions are nonzero.

Remark 4.10 With some additional justifications, the statement in Theorem 4.8 may now equally be seen to hold for the class functions $\phi_{\llbracket X \rrbracket}$. If $\llbracket X \rrbracket$ has odd defect then the same statement holds verbatim.

Assume now in the statement of Theorem 4.8 that $\delta(w) = e$, so that $w \in W_{\mathcal{I}}^e$. Then it makes sense to consider $\phi_{\llbracket X \rrbracket}(w)$ for any $\llbracket X \rrbracket \in \mathcal{S}_n^{\text{ev},e}$. Clearly $w_2 \in W_{\mathcal{I}}^{e+j}$, and if $\lambda \in \mathcal{H}_{d,j}(X)$ then $\llbracket X \setminus \lambda \rrbracket \in \mathcal{S}_n^{\text{ev},e+j}$, so the term $\phi_{\llbracket X \setminus \lambda \rrbracket}(w_2)$ makes sense. Hence, restricting the symbols to $\mathcal{S}_n^{\text{ev},2e}$, the statement in Theorem 4.8 continues to hold.

5 Lusztig series

In the next few sections we consider a general connected reductive group \mathbf{G} defined over $\mathbb{F} = \overline{\mathbb{F}}_p$ with Frobenius endomorphism $F : \mathbf{G} \rightarrow \mathbf{G}$. We will follow the setup in [28], see also the exposition in [13, Chap. 2]. This setting, whilst a little less frequently used, is more convenient as we wish to discuss character values of Deligne–Lusztig characters. In this section we just outline some notation.

Let $\mathbf{T} \leq \mathbf{B} \leq \mathbf{G}$ be a fixed F -stable maximal torus and Borel subgroup of \mathbf{G} . Let $X = X(\mathbf{T}) = \text{Hom}(\mathbf{T}, \mathbb{G}_m)$, which we view as a \mathbb{Z} -module. For any $\phi : \mathbf{T} \rightarrow \mathbf{T}$ a morphism of algebraic groups we denote by $\phi^* : X \rightarrow X$ the map given by $\phi^*(\chi) = \chi \circ \phi$ for all $\chi \in X$.

For each $w \in W := \mathbf{N}_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ we fix an element $n_w \in \mathbf{N}_{\mathbf{G}}(\mathbf{T})$ such that $w = n_w \mathbf{T}$. Let $\iota_g : \mathbf{G} \rightarrow \mathbf{G}$, with $g \in \mathbf{G}$, be the inner automorphism given by $\iota_g(x) = gxg^{-1}$. Then $Fw := F\iota_{n_w}$ and $wF := \iota_{n_w}F$ are also Frobenius endomorphisms of \mathbf{G} stabilising \mathbf{T} . We write w^* instead of $(\iota_{n_w}|_{\mathbf{T}})^*$ and for brevity we let ${}^w\lambda = w^{*-1}(\lambda)$ for all $w \in W$ and $\lambda \in X$.

If $\mathbb{Z}_{(p)}$ is the localisation of \mathbb{Z} at the prime ideal $(p) \subseteq \mathbb{Z}$ containing p then $V = \mathbb{Z}_{(p)} \otimes_{\mathbb{Z}} X$ is a free $\mathbb{Z}_{(p)}$ -module of finite rank. We will identify X with its image in V under the canonical map $x \mapsto 1 \otimes x$ and similarly any homomorphism $\gamma : X \rightarrow X$ is identified with $1 \otimes \gamma : V \rightarrow V$. Note the quotient V/X is a torsion \mathbb{Z} -module.

Abusing terminology we call $W_{\text{aff}} = X \rtimes W$ the affine Weyl group. We have a natural action of W_{aff} on V given by $(\chi, z) \cdot \lambda = \chi + {}^z\lambda$ for any $(\chi, z) \in W_{\text{aff}}$ and $\lambda \in V$. Let $W_{\text{aff}}(\lambda)$ be the stabiliser of λ and let $W(\lambda) = \{w \in W \mid \lambda - {}^w\lambda \in X\}$ be the projection of $W_{\text{aff}}(\lambda)$ onto W .

If $\mathbb{Z}\Phi \subseteq X$ is the \mathbb{Z} -submodule generated by the roots then $W_{\text{aff}}^\circ = \mathbb{Z}\Phi \rtimes W \triangleleft W_{\text{aff}}$ is the usual affine Weyl group. Again we let $W_{\text{aff}}^\circ(\lambda)$ be the stabiliser of λ and let $W^\circ(\lambda)$ be the projection of $W_{\text{aff}}^\circ(\lambda) \triangleleft W_{\text{aff}}(\lambda)$ on to W . We have $W^\circ(\lambda)$ is the kernel of the homomorphism $W(\lambda) \rightarrow X/\mathbb{Z}\Phi$ given by $w \mapsto (\lambda - {}^w\lambda) + \mathbb{Z}\Phi$.

Given $\lambda \in V$ and $w \in W$ we let $\lambda_w = F^*\lambda - {}^w\lambda \in V$. Note that $(\lambda + \lambda')_w = \lambda_w + \lambda'_w$ for any $\lambda, \lambda' \in V$. Moreover, for any $z \in W$ a straightforward calculation shows that

$$F^{-1}(z)\lambda_w = ({}^z\lambda)_{F^{-1}(z)wz^{-1}}. \tag{5.1}$$

The set $\mathbf{Z}_W(\lambda, F) = \{w \in W \mid \lambda_w \in X\}$ is either empty or a coset of the group $W(\lambda)$.

Denote by $\mathcal{C}_W(X, F)$ (resp. $\mathcal{D}_W(X, F)$) the set of all pairs (λ, w) (resp. (λ, a)) with $\lambda \in V$ and $w \in \mathbf{Z}_W(\lambda, F)$ (resp. $a = wW^\circ(\lambda) \subseteq \mathbf{Z}_W(\lambda, F)$). By (5.1), we have a natural action of W_{aff} on $\mathcal{C}_W(X, F)$ via

$$(\chi, z) \cdot (\lambda, w) = (\chi + {}^z\lambda, F^{-1}(z)wz^{-1})$$

and a similar action on $\mathcal{D}_W(X, F)$ as $W({}^z\lambda) = {}^zW(\lambda)$ for any $z \in W$ and $\lambda \in V$.

We now fix an injective homomorphism $\kappa : \mathbb{F}^\times \rightarrow \mathbb{C}^\times$. As \mathbf{T}^{wF} is a p' -group, we have a bijection $X(\mathbf{T}^{wF}) \rightarrow \text{Irr}(\mathbf{T}^{wF})$ given by $\chi \mapsto \kappa \circ \chi$. Given a pair $(\lambda, w) \in \mathcal{C}_W(X, F)$ we set

$$\lambda_{wF} = \kappa \circ (\lambda_w|_{\mathbf{T}^{wF}}) \in \text{Irr}(\mathbf{T}^{wF}).$$

The following is straightforward; see [28, Lem. 6.2] and [13, Lem. 2.4.8].

Lemma 5.1 *Fix $w \in W$ and let $V(w) = \{\lambda \in V \mid \lambda_w \in X\}$. Then the map $V(w) \rightarrow \text{Irr}(\mathbf{T}^{wF})$ defined by $\lambda \mapsto \lambda_{wF}$ is a surjective \mathbb{Z} -module homomorphism with kernel $X \subseteq V(w)$.*

Given $w \in W$ and $\theta \in \text{Irr}(\mathbf{T}^{wF})$ we denote by $R_w^{\mathbf{G}}(\theta)$ the virtual character of \mathbf{G}^F defined in [7, Def. 1.9]. As usual we extend this by linearity to a map on all class functions.

Moreover, for any $(\lambda, w) \in \mathcal{C}_W(X, F)$ we set $R_w^{\mathbf{G}}(\lambda) := R_w^{\mathbf{G}}(\lambda_{wF})$. We then define for any pair $(\lambda, a) \in \mathcal{D}_W(X, F)$ the set

$$\mathcal{E}(\mathbf{G}^F, \lambda, a) = \{\rho \in \text{Irr}(\mathbf{G}^F) \mid \langle R_w^{\mathbf{G}}(\lambda), \rho \rangle \neq 0 \text{ for some } w \in a\}.$$

This is a rational Lusztig series of \mathbf{G}^F contained in the geometric series

$$\mathcal{E}(\mathbf{G}^F, \lambda) = \{\rho \in \text{Irr}(\mathbf{G}^F) \mid \langle R_w^{\mathbf{G}}(\lambda), \rho \rangle \neq 0 \text{ for some } w \in \mathbf{Z}_W(\lambda, F)\}$$

indexed by $\lambda \in V$. We have $\mathcal{E}(\mathbf{G}^F, \lambda, a) = \mathcal{E}(\mathbf{G}^F, \mu, b)$ if and only if (λ, a) and (μ, b) are in the same W_{aff} -orbit.

Suppose now that $\iota : \mathbf{G} \rightarrow \tilde{\mathbf{G}}$ is a regular embedding. Then $\tilde{\mathbf{T}} = \mathbf{T} \cdot \mathbf{Z}(\mathbf{G})$ is an F -stable maximal torus of $\tilde{\mathbf{G}}$ and if we let $\tilde{X} = X(\tilde{\mathbf{T}})$ then we have a surjective \mathbb{Z} -module homomorphism $\iota^* : \tilde{X} \rightarrow X$ given by $\iota^*(\chi) = \chi \circ \iota$. Note this maps the roots of $\tilde{\mathbf{G}}$ in \tilde{X} bijectively onto the roots of \mathbf{G} in X . Through ι we identify W with the Weyl group $\mathbf{N}_{\tilde{\mathbf{G}}}(\tilde{\mathbf{T}})/\tilde{\mathbf{T}}$ of $\tilde{\mathbf{G}}$.

Lemma 5.2 *For any $(\tilde{\lambda}, a) \in \mathcal{D}_W(\tilde{X}, F)$ we have $W(\tilde{\lambda}) = W^\circ(\tilde{\lambda}) = W^\circ(\lambda)$ and*

$$\mathcal{E}(\mathbf{G}^F, \lambda, a) = \{\rho \in \text{Irr}(\mathbf{G}^F) \mid \langle \rho, \text{Res}_{\mathbf{G}^F}^{\tilde{\mathbf{G}}^F}(\tilde{\rho}) \rangle \neq 0 \text{ for some } \tilde{\rho} \in \mathcal{E}(\tilde{\mathbf{G}}^F, \tilde{\lambda}, a)\}$$

where $\lambda = \iota^*(\tilde{\lambda})$.

Proof By the above remark we have $\tilde{\lambda} - w\tilde{\lambda} \in \mathbb{Z}\Phi$ if and only if $\lambda - w\lambda = \iota^*(\tilde{\lambda} - w\tilde{\lambda}) \in \mathbb{Z}\Phi$ which shows that $W^\circ(\tilde{\lambda}) = W^\circ(\lambda)$. From the proof of [8, Lem. 11.2.1] we see that the image of the map $W(\tilde{\lambda}) \rightarrow \tilde{X}/\mathbb{Z}\Phi$ has p' -order but as $\mathbf{Z}(\tilde{\mathbf{G}})$ is connected the quotient $\tilde{X}/\mathbb{Z}\Phi$ has trivial p' -torsion so $W(\tilde{\lambda}) = W^\circ(\tilde{\lambda})$. The second statement is [5, Prop. 11.7]. \square

6 A character bound from the Mackey formula

We denote by $W:F$ the semidirect product of W with the group $\langle F \rangle \leq \text{Aut}(W)$ such that $FwF^{-1} = F(w)$ for all $w \in W$. The unique coset $WF \subseteq W:F$ of W containing F is a W -set under conjugation and for $w \in W$ we write $\mathbf{C}_W(wF)$ for the stabiliser of wF under this action.

For each $w \in W$ we choose an element $g_w \in \mathbf{G}$ such that $g_w^{-1}F(g_w) = n_w \in \mathbf{N}_{\mathbf{G}}(\mathbf{T})$. As usual the map $wF \mapsto \mathbf{T}_w := {}^{g_w}\mathbf{T}$ yields a bijection between the orbits of W acting on WF and the \mathbf{G}^F -classes of F -stable maximal tori. Note that $t \mapsto {}^{g_w}t$ gives an isomorphism $\mathbf{T}^{wF} \rightarrow \mathbf{T}_w^F$. In particular, we have a bijection $\text{Irr}(\mathbf{T}_w^F) \rightarrow \text{Irr}(\mathbf{T}^{wF})$ given by $\theta \mapsto {}^{g_w^{-1}}\theta = \theta \circ \iota_{g_w}$.

Remark 6.1 For $\theta \in \text{Irr}(\mathbf{T}_w^F)$ we have a Deligne–Lusztig character $R_{\mathbf{T}_w}^{\mathbf{G}}(\theta)$ defined as in [7, 1.20] which satisfies $R_{\mathbf{T}_w}^{\mathbf{G}}(\theta) = R_w^{\mathbf{G}}({}^{g_w^{-1}}\theta)$. We will implicitly use this equality in what follows.

We have an action of $\mathbf{C}_W(wF)$ on $\text{Irr}(\mathbf{T}^{wF})$ by setting ${}^z\theta = \theta \circ \iota_{n_z}^{-1}$ for any $z \in \mathbf{C}_W(wF)$ and $\theta \in \text{Irr}(\mathbf{T}^{wF})$. We denote by $\mathbf{C}_W(wF, \theta) \leq \mathbf{C}_W(wF)$ the stabiliser of $\theta \in \text{Irr}(\mathbf{T}^{wF})$. Now for any $(\lambda, w) \in \mathcal{C}_W(X, F)$ and $z \in W$ it follows from (5.1) that ${}^z(\lambda_{wF}) = (F(z))^{*-1}\lambda|_{z w F z^{-1}}$. Therefore,

$$F(\mathbf{C}_W(wF, \lambda_{wF})) = \mathbf{C}_W(\lambda)(Fw), \tag{6.1}$$

where the latter stabiliser is calculated with respect to the action of $W(\lambda)$ on $FZ_W(\lambda, F)$.

A regular semisimple element $g \in G^F$ is said to be of type $wF \in WF$ if $C_G^\circ(g)$ is G^F -conjugate to T_w . Of course, the type is only determined up to W -conjugacy. Moreover, an element g of type wF is then G^F -conjugate to an element of the form ${}^{g_w}t$ with $t \in T^{wF}$.

As in [8], if H is a finite group then we denote by $\pi_h^H \in \text{Class}(H)$ the function taking the value $|C_H(h)|$ on $\text{Cl}_H(h)$ and the value 0 on $H - \text{Cl}_H(h)$. For any $f \in \text{Class}(H)$ we then have $\langle f, \pi_h^H \rangle = f(h)$. We also write $[g, h] = g^{-1}h^{-1}gh$ for the commutator of $g, h \in H$.

Theorem 6.2 *Assume $g \in G^F$ is a regular semisimple element of type wF . Fix a pair $(\lambda, a) \in \mathcal{D}_W(X, F)$ and let*

$$\mathcal{X}_w(\lambda, a) := \{z \in W \mid [z, Fw]w^{-1}a = W^\circ(\lambda)\}.$$

If $\chi \in \mathcal{E}(G^F, \lambda, a)$ and $x \in a$ then

$$|\chi(g)| \leq \sum_{z \in C_W(Fw) \setminus \mathcal{X}_w(\lambda, a) / C_W(Fx, \lambda)} (|C_W(Fw)| / |C_{W(z\lambda)}(Fw)|^{\frac{1}{2}}) \leq |W| \cdot |C_W(Fw)|$$

Proof Assume $t \in T^{wF}$ is such that ${}^{g_w}t \in \text{Cl}_{G^F}(g)$. By [7, Prop. 9.18], see also [8, Prop. 10.3.6], we have $\pi_g^{G^F} = R_w^G(\pi_t^{T^{wF}}) = \sum_{\theta \in \text{Irr}(T^{wF})} \theta(t^{-1})R_w^G(\theta)$. For any $\theta \in \text{Irr}(T^{wF})$ we have by the Mackey formula for tori, see [7, Thm. 6.8] or [8, Cor. 9.3.1], that

$$\sum_{\chi \in \text{Irr}(G^F)} \langle \chi, R_w^G(\theta) \rangle^2 = \langle R_w^G(\theta), R_w^G(\theta) \rangle = |C_W(wF, \theta)|.$$

Applying this to $\langle \chi, \pi_g^{G^F} \rangle$ we get

$$|\chi(g)| \leq \sum_{\theta \in \text{Irr}(T^{wF})} |\langle \chi, R_w^G(\theta) \rangle| \leq \sum_{\theta \in \text{Irr}(T^{wF})} |C_W(wF, \theta)|^{\frac{1}{2}}. \tag{6.2}$$

Let $V(w)$ be as in Lemma 5.1 so that $V(w)/X \cong \text{Irr}(T^{wF})$. If $\mu \in V(w)$ is such that $\langle \chi, R_w^G(\mu) \rangle \neq 0$ then we must have $(\mu, wW^\circ(\mu))$ and (λ, a) are in the same W_{aff} -orbit by the disjointness of Lusztig series. This happens if and only if there exists a $z \in W$ such that $\mu - {}^z\lambda \in X$ and

$$F^{-1}(z)az^{-1} = wW^\circ(\mu) = wzW^\circ(\lambda)z^{-1}.$$

This last condition is equivalent to $z \in \mathcal{X}_w(\lambda, a)$.

If $z \in \mathcal{X}_w(\lambda, a)$ and $v \in C_W(Fx, \lambda)$ then one checks easily that $zv \in \mathcal{X}_w(\lambda, a)$, and clearly ${}^{zv}\lambda = {}^z\lambda$. Hence, the sum in (6.2) can be taken over $\mathcal{X}_w(\lambda, a) / C_W(Fx, \lambda)$ with θ replaced by $({}^z\lambda)_{wF}$. The group $C_W(Fw) = F(C_W(wF))$ acts on $\mathcal{X}_w(\lambda, a)$ by left multiplication, and the term in the sum is constant on orbits. Hence, we can sum as in the statement once we multiply through by the size of the orbit $|C_W(Fw) / C_{W(z\lambda)}(Fw)|$. We now cancel terms using (6.1). □

Remark 6.3 The above follows the same argument as [17, Thm. 5.4] where one finds the bound $|C_W(Fw)|$. The above yields this bound when the sum contains only one term. However, this does not hold in general so this should be corrected as above. In the extreme cases where $W(\lambda) = W$ or $W^\circ(\lambda) = \{1\}$ then this does give the bound $|C_W(Fw)|$.

7 Further bounds from Deligne–Lusztig characters

The following result giving the value of a Deligne–Lusztig character at a regular semisimple element is well known. We simply translate this into the setup we utilise here, see [12, Prop. 4.5.8] for an equivalent formulation.

Proposition 7.1 *Assume $g \in \mathbf{G}^F$ is a regular semisimple element of type wF and let $t \in \mathbf{T}^{wF}$ be such that ${}^s w t \in \mathbf{C}_{\mathbf{G}^F}(g)$. Then for any $(\lambda, x) \in \mathcal{C}_W(X, F)$ we have*

$$R_x^{\mathbf{G}}(\lambda)(g) = \sum_{\substack{\mu \in V(w)/X \\ (\mu, x) \in \mathbf{C}_W(\lambda, w)}} |\mathbf{C}_{W(\mu)}(Fw)| \mu_{wF}(t) = |\mathbf{C}_{W(\lambda)}(Fx)| \sum_{\substack{z \in W/\mathbf{C}_{W(\lambda)}(Fx) \\ Fw = zFxz^{-1}}} ({}^z \lambda)_{wF}(t).$$

In particular, we have $|R_x^{\mathbf{G}}(\lambda)(g)| \leq |\mathbf{C}_W(Fw)|$.

Proof As noted in the proof of Theorem 6.2 we simply have to calculate $\langle R_x^{\mathbf{G}}(\lambda), R_w^{\mathbf{G}}(\pi_t^{\mathbf{T}^{wF}}) \rangle$. The statement now follows immediately from the inner product formula for Deligne–Lusztig characters and the identification $V(w)/X \cong \text{Irr}(\mathbf{T}^{wF})$. \square

To make invoking Lusztig’s classification results for the irreducible characters of \mathbf{G}^F simpler it will be beneficial to assume that $\mathbf{Z}(\mathbf{G})$ is connected. As usual we invoke a regular embedding to achieve this. We note that for our purposes we do not need the significantly more difficult multiplicity freeness results obtained by Lusztig.

Lemma 7.2 *Assume $\mathbf{G} \rightarrow \tilde{\mathbf{G}}$ is a regular embedding and let $\chi \in \text{Irr}(\mathbf{G}^F)$ be an irreducible constituent of $\text{Res}_{\mathbf{G}^F}^{\tilde{\mathbf{G}}^F}(\tilde{\chi})$ for some $\tilde{\chi} \in \text{Irr}(\tilde{\mathbf{G}}^F)$. Then for any semisimple element $g \in \mathbf{G}^F$ we have*

$$|\chi(g)| \leq |\tilde{\mathbf{G}}^F / \tilde{\mathbf{G}}_{\chi}^F|^{-1} |\tilde{\chi}(g)| \leq |\tilde{\chi}(g)|,$$

where $\tilde{\mathbf{G}}_{\chi}^F$ is the stabiliser of χ in $\tilde{\mathbf{G}}^F$.

Proof Let $\mathbf{T} \leq \mathbf{G}$ be an F -stable maximal torus and $\theta \in \text{Irr}(\mathbf{T}^F)$. We then have $\tilde{\mathbf{T}} = \mathbf{T} \cdot \mathbf{Z}(\tilde{\mathbf{G}})$ is an F -stable maximal torus of $\tilde{\mathbf{G}}$. Let $\tilde{\theta} \in \text{Irr}(\tilde{\mathbf{T}}^F)$ be an irreducible character such that $\text{Res}_{\tilde{\mathbf{T}}^F}^{\tilde{\mathbf{G}}^F}(\tilde{\theta}) = \theta$. By [5, Prop. 10.10] we have $\text{Res}_{\mathbf{G}^F}^{\tilde{\mathbf{G}}^F}(R_{\tilde{\mathbf{T}}}^{\tilde{\mathbf{G}}}(\tilde{\theta})) = R_{\mathbf{T}}^{\mathbf{G}}(\theta)$. Therefore, by Frobenius reciprocity,

$$\langle \chi, R_{\mathbf{T}}^{\mathbf{G}}(\theta) \rangle = \langle \text{Ind}_{\mathbf{G}^F}^{\tilde{\mathbf{G}}^F}(\chi), R_{\tilde{\mathbf{T}}}^{\tilde{\mathbf{G}}}(\tilde{\theta}) \rangle.$$

This shows that for any $c \in \tilde{\mathbf{G}}^F$ we have ${}^c \chi \in \text{Irr}(\mathbf{G}^F)$ and χ have the same uniform projection. As $\pi_g^{\mathbf{G}^F}$ is a uniform function this means that $\chi(g) = \langle \chi, \pi_g^{\mathbf{G}^F} \rangle = \langle {}^c \chi, \pi_g^{\mathbf{G}^F} \rangle = {}^c \chi(g)$. Now by Clifford’s Theorem we have

$$\text{Res}_{\mathbf{G}^F}^{\tilde{\mathbf{G}}^F}(\tilde{\chi}) = e \sum_{c \in \tilde{\mathbf{G}}^F / \tilde{\mathbf{G}}_{\chi}^F} {}^c \chi$$

for some integer $e \geq 1$. Hence $\tilde{\chi}(g) = e |\tilde{\mathbf{G}}^F / \tilde{\mathbf{G}}_{\chi}^F| \chi(g)$ giving the bound. \square

Fix a pair $(\lambda, a) \in \mathcal{D}_W(X, F)$. For a class function $f \in \text{Class}(Fa)$ on the coset of $W^\circ(\lambda)$ we define a corresponding class function

$$\mathcal{R}_{\lambda, a}^{\mathbf{G}}(f) = \frac{1}{|W^\circ(\lambda)|} \sum_{x \in a} f(Fx) R_x^{\mathbf{G}}(\lambda). \tag{7.1}$$

of \mathbf{G}^F . Clearly this is contained in the subspace of all \mathbb{C} -class functions $\text{Class}(\mathbf{G}^F, \lambda, a)$ spanned by the Lusztig series $\mathcal{E}(\mathbf{G}^F, \lambda, a)$. In fact, $\mathcal{R}_{\lambda,a}^{\mathbf{G}}$ gives an isomorphism $\text{Class}(Fa) \rightarrow \text{Class}_0(\mathbf{G}^F, \lambda, a)$ onto the subspace spanned by $\{R_x^{\mathbf{G}}(\lambda) \mid x \in a\}$; see the arguments in [8, §11.6].

Suppose we choose a representative $a = wW^\circ(\lambda)$ of the coset. We may then form the semidirect product $W^\circ(\lambda):Fw$ by the group $\langle Fw \rangle \leq \text{Aut}(W^\circ(\lambda))$ as above. The coset of $W^\circ(\lambda)$ in $W^\circ(\lambda):Fw$ containing Fw can be identified, $W^\circ(\lambda)$ -equivariantly, with the same coset in $W:F$.

We denote by $\text{Irr}(Fw.W^\circ(\lambda))$ the set of restrictions $\text{Res}_{Fa}^{W^\circ(\lambda):Fw}(\tilde{\phi})$, where $\tilde{\phi} \in \text{Irr}(W^\circ(\lambda):Fw)$ restricts irreducibly to $W^\circ(\lambda)$. These functions on the coset Fa depend on our choice of representative w , so we include a period to indicate this choice. There is, however, a natural choice $w_a \in a$ which is the element of minimal length (determined by our choice of Borel subgroup \mathbf{B}).

Lemma 7.3 *Assume $g \in \mathbf{G}^F$ is a regular semisimple element of type wF . For any irreducible character $\phi \in \text{Irr}(Fx.W^\circ(\lambda))$, with $x \in a$, we have*

$$|\mathcal{R}_{\lambda,a}^{\mathbf{G}}(\phi)(g)| \leq |\mathbf{C}_W(Fw)|$$

Proof By Proposition 7.1 we have

$$|\mathcal{R}_{\lambda,a}^{\mathbf{G}}(\phi)(g)| \leq |\mathbf{C}_W(Fw)| \left(\frac{1}{|W^\circ(\lambda)|} \sum_{y \in W^\circ(\lambda)} |\phi(Fxy)| \right),$$

but by [19, Lem. 8.14(c)], the sum on the right hand side is equal to 1. □

Lemma 7.4 *Assume $g \in \mathbf{G}^F$ is a regular semisimple element of type wF . Then for any class function $f \in \text{Class}(Fa)$ we have*

$$\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f)(g) = \sum_{\substack{z \in \mathbf{C}_W(Fw) \backslash W/W^\circ(\lambda) \\ Fx = z^{-1}Fwz \in Fa}} \frac{f(Fx) \cdot R_x^{\mathbf{G}}(\lambda)(g)}{|\mathbf{C}_{W^\circ(\lambda)}(Fx)|}.$$

Proof By Proposition 7.1 we may restrict the sum over $x \in a$, found in the definition of $\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f)$, to those elements satisfying $Fx \in \text{Cl}_W(Fw)$. Alternatively, via the bijection $\mathbf{C}_W(Fw) \backslash W \rightarrow \text{Cl}_W(Fw)$, given by $\mathbf{C}_W(Fw)z \mapsto z^{-1}Fwz$, we can sum over all cosets $\mathbf{C}_W(Fw)z \in \mathbf{C}_W(Fw) \backslash W$ such that $z^{-1}Fwz \in Fa$. Grouping together elements in the same $W^\circ(\lambda)$ -orbit and bringing $|W^\circ(\lambda)|$ into the sum gives the statement. □

Observe that, if $W^\circ(\lambda) = W$ and $w \in a$ then $\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f)(g) = f(Fw)\lambda_{wF}(t)$, see [24, Prop. 3.3]. Together with Theorems 4.7 and 4.9, this implies

Corollary 7.5 *Suppose \mathbf{G} is a quasisimple group of classical type A to D. If $g \in \mathbf{G}^F$ is a regular semisimple element of cycle length k and $\chi \in \text{Irr}(\mathbf{G}^F)$ is a unipotent character, then $|\chi(g)| \leq 2^{2k} \cdot k!$.*

Now suppose $[\mathbf{G}, \mathbf{G}]$ is quasisimple of type A_{n-1} . In this case we have W is isomorphic to \mathfrak{S}_n and F induces an inner automorphism on W , which is either trivial or conjugation by the longest element. Hence, the coset WF can be identified with W so it makes sense to speak of the cycle type of an element of WF .

Corollary 7.6 *Assume all the quasi-simple components of G are of type A. If $g \in G^F$ is a regular semisimple element of type wF then*

$$|\chi(g)| \leq |C_W(Fw)|$$

for any $\chi \in \text{Irr}(G^F)$. Moreover, suppose $[G, G]$ is quasisimple of type A_{n-1} with $n \geq 2$. If wF has cycle length $k \geq 1$ then $|\chi(g)| \leq k! \cdot n^k$ for any $\chi \in \text{Irr}(G^F)$.

Proof By Lemma 7.2, we can assume that $Z(G)$ is connected. In that case every irreducible character is, up to sign, of the form $\mathcal{R}_{\lambda,a}^G(\phi)$ with $\phi \in \text{Irr}(Fw.W(\lambda))$ so this is just Lemma 7.3.

For the final statement we need only show that $|C_{\mathfrak{S}_n}(w)| \leq k! \cdot n^k$. If all cycles have the same length, say $m \geq 1$, then $C_{\mathfrak{S}_n}(w) \cong C_m \wr \mathfrak{S}_k$ so $|C_{\mathfrak{S}_n}(w)| = k! \cdot m^k$. Now an arbitrary w may be written as a pairwise commuting product $w = w_1 \cdots w_r$ such that for each $1 \leq i \leq r$ we have w_i is a product of $k_i \geq 1$ disjoint cycles of length $m_i \geq 1$ and the lengths m_1, \dots, m_r are pairwise distinct. We then have

$$C_{\mathfrak{S}_n}(w) \cong C_{\mathfrak{S}_{k_1 m_1}}(w_1) \times \cdots \times C_{\mathfrak{S}_{k_r m_r}}(w_r),$$

and by the previous calculation

$$|C_{\mathfrak{S}_n}(w)| = (k_1! \cdot m_1^{k_1}) \cdots (k_r! \cdot m_r^{k_r}) \leq k! \cdot n^{k_1 + \cdots + k_r} = k! \cdot n^k.$$

□

For the next statement we wish to define an integer $r(W, F) \geq 0$ as follows. Let $\mathbb{S} \subseteq W$ be the set of Coxeter generators determined by our choice of Borel B . Write $W = W_1 \cdots W_m$ as a direct product of its irreducible components, all of which are assumed to be of type A through D. We then have a corresponding decomposition $\mathbb{S} = \mathbb{S}_1 \sqcup \cdots \sqcup \mathbb{S}_m$. The Frobenius F permutes the W_i . Suppose first that it does so transitively. Then we define

$$r(W, F) = \begin{cases} 0 & \text{if } W_1 \text{ is of type } A_n \text{ with } n \geq 0, \\ |\mathbb{S}_1| & \text{otherwise.} \end{cases}$$

Here we consider the trivial group as being of type A_0 .

Now grouping together the W_i we can write $W = W^{(1)} \cdots W^{(n)}$ where each $W^{(i)}$ is an F -stable subgroup such that F permutes transitively its irreducible components. Hence, we are in the previous situation and we define $r(W, F) = r(W^{(1)}, F) + \cdots + r(W^{(n)}, F)$.

Theorem 7.7 *Assume $g \in G^F$ is a regular semisimple element of type wF and every quasi-simple component of G is of classical type A to D. If F is a Frobenius endomorphism then for any irreducible character $\chi \in \mathcal{E}(G^F, \lambda, a)$ we have*

$$|\chi(g)| \leq 2^r \cdot |C_W(Fw)|$$

where $w_a \in a$ is the unique element of minimal length and $r = r(W^\circ(\lambda), Fw_a)$ is defined as above.

Proof Again, by Lemma 7.2 we can assume $Z(G)$ is connected. By Lemma 5.2 this implies that $W(\lambda) = W^\circ(\lambda)$ and $a = Z_W(\lambda, F)$ so $\mathcal{E}(G^F, \lambda, a) = \mathcal{E}(G^F, \lambda)$. Recall that in [28, Chp. 4] Lusztig has defined a partition of $\text{Irr}(W(\lambda))$ into families.

Denote by $w_a \in a = Z_W(\lambda, F)$ the unique element of minimal length. The automorphism $\gamma := Fw_a$ of $W(\lambda)$ permutes the families. Suppose $\mathcal{F} \subseteq \text{Irr}(W(\lambda))$ is a γ -stable family. For each γ -fixed character $\phi \in \mathcal{F}'$ we fix an extension $\tilde{\phi} \in \text{Irr}(W(\lambda) : \gamma)$ that is realisable over \mathbb{Q} .

Suppose $\mathcal{F} \subseteq \text{Irr}(W(\lambda))$ is γ -stable and let

$$\mathcal{E}(\mathbf{G}^F, \lambda, \mathcal{F}) = \{\chi \in \text{Irr}(\mathbf{G}^F) \mid \langle \chi, \mathcal{R}_{\lambda,a}^{\mathbf{G}}(\tilde{\phi}) \rangle \neq 0 \text{ for some } \phi \in \mathcal{F}'\}.$$

These sets partition $\mathcal{E}(\mathbf{G}^F, \lambda)$; see [28, Thm. 6.17]. Associated to \mathcal{F} we have a corresponding finite group $\mathcal{G}_{\mathcal{F}}$.

As all factors are of classical type, we have $\mathcal{G}_{\mathcal{F}}$ is a (possibly trivial) elementary abelian 2-group. We also have two sets $\overline{\mathcal{M}}(\mathcal{G}_{\mathcal{F}}, \gamma)$ and $\mathcal{M}(\mathcal{G}_{\mathcal{F}}, \gamma)$ and a pairing

$$\{-, -\} : \overline{\mathcal{M}}(\mathcal{G}_{\mathcal{F}}, \gamma) \times \mathcal{M}(\mathcal{G}_{\mathcal{F}}, \gamma) \rightarrow \mathbb{C}.$$

From the formula for this pairing, we see that

$$|\{\bar{x}, x\}| = |\mathcal{G}_{\mathcal{F}}|^{-1}$$

for any $\bar{x} \in \overline{\mathcal{M}}(\mathcal{G}_{\mathcal{F}}, \gamma)$ and $x \in \mathcal{M}(\mathcal{G}_{\mathcal{F}}, \gamma)$

By [28, Thm. 4.23], we have a bijection $\mathcal{E}(\mathbf{G}^F, \lambda, \mathcal{F}) \rightarrow \overline{\mathcal{M}}(\mathcal{G}_{\mathcal{F}}, \gamma)$, which we denote by $\chi \mapsto x_{\chi}$, and an injection $\mathcal{F}' \rightarrow \mathcal{M}(\mathcal{G}_{\mathcal{F}}, \gamma)$, denoted by $\phi \mapsto x_{\tilde{\phi}}$. This latter map depends on our choice of extension. Now, by [28, 4.26.1], if $\chi \in \mathcal{E}(\mathbf{G}^F, \lambda, \mathcal{F})$ then

$$\chi(g) = \pm \sum_{\phi \in \mathcal{F}'} \{\bar{x}_{\chi}, x_{\tilde{\phi}}\} \mathcal{R}_{\lambda,a}^{\mathbf{G}}(\tilde{\phi})(g).$$

The group of roots of unity acts on $\mathcal{M}(\mathcal{G}_{\mathcal{F}}, \gamma)$, and the number of orbits is the same as $|\overline{\mathcal{M}}(\mathcal{G}_{\mathcal{F}}, \gamma)| = |\mathcal{G}_{\mathcal{F}}|$. It follows from [28, 4.21.6] that $|\mathcal{F}'| \leq |\mathcal{G}_{\mathcal{F}}|^2$. We now use Lemma 7.3. □

We assume F is a Frobenius endomorphism. If $[\mathbf{G}, \mathbf{G}]$ is quasisimple of type B_n or C_n then $W \cong W_{\mathcal{I}}$ is a hyperoctahedral group, and F induces the identity on W . If $[\mathbf{G}, \mathbf{G}]$ is quasisimple of type D_n then $W \cong W_{\mathcal{I}}^0$ and either F, F^2 , or F^3 induces the identity on W . When F^2 induces the identity on W , we have an embedding $W : F \rightarrow W_{\mathcal{I}}$. Thus, it makes sense to speak of the *cycles* of an element of the coset WF . The following is now just a simple application of Lemma 4.3.

Corollary 7.8 *Assume $[\mathbf{G}, \mathbf{G}]$ is quasisimple of type B_n ($n \geq 2$), C_n ($n \geq 2$), or D_n ($n \geq 4$), and F is a Frobenius endomorphism with F^2 inducing the identity on W . If $wF \in WF$ has cycle length $k \geq 1$ and pairwise distinct cycles, then for any regular semisimple element $g \in \mathbf{G}^F$ of type wF we have*

$$|\chi(g)| \leq 2^{n+k} \cdot n^k$$

for all $\chi \in \text{Irr}(\mathbf{G}^F)$.

8 Quadratic unipotent characters

Recall that a bound for the values of unipotent characters at regular semisimple elements was obtained in Corollary 7.5. In this section, we establish a bound for the more general class of quadratic unipotent characters. Consider a connected reductive group \mathbf{G} whose center $\mathbf{Z}(\mathbf{G})$ is connected, of dimension 1, and whose derived subgroup $\mathbf{G}_{\text{der}} = [\mathbf{G}, \mathbf{G}] \leq \mathbf{G}$ is a symplectic or special orthogonal group. We specify \mathbf{G} by its root datum $(X, \Phi, \check{X}, \check{\Phi})$. Firstly we have $X = \bigoplus_{i=0}^n \mathbb{Z}e_i$ and $\check{X} = \bigoplus_{i=0}^n \mathbb{Z}\check{e}_i$ with perfect pairing $\langle -, - \rangle : X \times \check{X} \rightarrow \mathbb{Z}$ given by $\langle e_i, \check{e}_j \rangle = \delta_{ij}$ (the Kronecker delta). We assume $n \geq 2$.

A set of simple roots $\alpha_1, \dots, \alpha_n$ and corresponding coroots $\check{\alpha}_1, \dots, \check{\alpha}_n$ are as follows. We have $(\alpha_1, \check{\alpha}_1)$ is one of the pairs $(-e_1, -2\check{e}_1)$, $(e_0 - 2e_1, -\check{e}_1)$, or $(e_0 - e_1 - e_2, -\check{e}_1 - \check{e}_2)$. Then for $2 \leq i \leq n$ we let $\alpha_i = e_{i-1} - e_i$ and $\check{\alpha}_i = \check{e}_{i-1} - \check{e}_i$. The choices correspond to whether \mathbf{G} is of type $\mathbf{B}_n, \mathbf{C}_n$, or \mathbf{D}_n , respectively.

One easily calculates that $(\langle \alpha_i, \check{\alpha}_j \rangle)$ is a Cartan matrix and $X/\mathbb{Z}\Phi \cong \mathbb{Z}$ is generated by $e_0 + \mathbb{Z}\Phi$. Let $X_{\text{der}} = X/(\mathbb{Z}e_0)$ and $\check{X}_{\text{der}} = \bigoplus_{i=1}^n \check{e}_i$. Denote by $\pi : X \rightarrow X_{\text{der}}$ the natural quotient map. We then have $(X_{\text{der}}, \bar{\Phi}, \check{X}_{\text{der}}, \check{\Phi})$ is the root datum of \mathbf{G}_{der} .

Fix a prime power $q = p^a$. We describe F by defining F^* as an endomorphism of X . For any $0 \leq i \leq n$ with $i \neq 1$ we have $F^*e_i = qe_i$. We then have F^*e_1 is either qe_1 or $q(e_0 - e_1)$ with this latter case occurring only when \mathbf{G}^F is of type ${}^2\mathbf{D}_n(q)$.

If $V_{\text{der}} = \mathbb{Z}_{(p)} \otimes_{\mathbb{Z}} X_{\text{der}}$ then we have $V_{\text{der}} \cong \bigoplus_{i=1}^n \mathbb{Z}_{(p)}\bar{e}_i$ and $V \cong \bigoplus_{i=1}^{n+1} \mathbb{Z}_{(p)}e_i$. The natural quotient map $X \rightarrow X_{\text{der}}$ extends to a surjective $\mathbb{Z}_{(p)}$ -module homomorphism $\pi : V \rightarrow V_{\text{der}}$. Given $\lambda \in V$ we have $W(\lambda) = W^\circ(\lambda) = W^\circ(\bar{\lambda})$ and if $\chi \in \mathcal{E}(\mathbf{G}^F, \lambda, a)$ then all irreducible constituents of χ are contained in $\mathcal{E}(\mathbf{G}_{\text{der}}^F, \bar{\lambda}, a)$, see Lemma 5.2.

Consider the totally ordered set $\mathcal{I} = \{-\bar{e}_n < \dots < -\bar{e}_1 < \bar{e}_1 < \dots < \bar{e}_n\}$. Letting F act on X via $q^{-1}F^*$ we get an action of $W : F$ on X_{der} that gives an injective homomorphism $W : F \rightarrow \mathfrak{S}_{\mathcal{I}}$. We implicitly identify $W : F$, hence also W , with its image which is contained in $W_{\mathcal{I}}$. Via this identification we can speak of the *signed cycle type* of any element of $W : F$.

Theorem 8.1 *Let $(\lambda, a) \in \mathcal{D}_W(X, F)$ be such that $2\bar{\lambda} \in X_{\text{der}}$ and let $g \in \mathbf{G}^F$ be a regular semisimple element of type wF . If wF has cycle length $k \geq 1$ and pairwise distinct cycles then*

$$|\chi(g)| \leq 2^{3k+4} \cdot k!$$

for any irreducible character $\chi \in \mathcal{E}(\mathbf{G}^F, \lambda, a)$.

Proof Recall that we have an isomorphism $\mathcal{R}_{\lambda,a}^{\mathbf{G}} : \text{Class}(Fa) \rightarrow \text{Class}_0(\mathbf{G}^F, \lambda, a)$. Thus, if $\text{pr} : \text{Class}(\mathbf{G}^F) \rightarrow \text{Class}(\mathbf{G}^F)$ is the projection onto the subspace of uniform functions then there exists a unique class function $f_\chi \in \text{Class}(Fa)$ such that $\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f_\chi) = \text{pr}(\chi)$.

Now $\text{pr}(\chi)$ and χ have the same value at g , so it suffices to bound the value of $\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f_\chi)$ at g . By Proposition 7.1 and Lemma 7.4, we have

$$|\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f_\chi)(g)| \leq \sum_{\substack{z \in \mathbf{C}_W(Fw) \backslash W/W(\lambda) \\ Fx = z^{-1}Fwz \in Fa}} \frac{|\mathbf{C}_W(Fx)|}{|\mathbf{C}_{W(\lambda)}(Fx)|} \cdot |f_\chi(Fx)|.$$

We can assume $\mathcal{R}_{\lambda,a}^{\mathbf{G}}(f_\chi)(g) \neq 0$ and hence assume that $Fx = z^{-1}Fwz \in Fa$ is a conjugate of Fw .

We now bound: $|\mathbf{C}_W(Fx)|/|\mathbf{C}_{W(\lambda)}(Fx)|$, the number of terms in the sum, and finally $|f_\chi(Fx)|$. Let us note that the number of terms in the sum is precisely the number of $W(\lambda)$ -orbits on Fa that meet the centraliser $\mathbf{C}_W(Fw)$.

We will take this case by case. First let us note that as wF has pairwise distinct cycles, so does its conjugate $Fx = zF(wF)F^{-1}z^{-1}$. Now, by replacing λ with an element in the same W_{aff} -orbit, we can assume that $\bar{\lambda} = \frac{1}{2}(\bar{e}_1 + \dots + \bar{e}_m)$ for some $0 \leq m \leq n$ where $\bar{\lambda} = 0$ when $m = 0$.

We set $\mathcal{I}_1 = \{\pm\bar{e}_i \mid 1 \leq i \leq m\}$ and $\mathcal{I}_0 = \mathcal{I} \setminus \mathcal{I}_1$ and let $H = W_{\mathcal{I}_0}W_{\mathcal{I}_1} \leq W_{\mathcal{I}}$. For convenience, we let $\pi_i = (\bar{e}_i, -\bar{e}_i) \in W_{\mathcal{I}}$ for any $1 \leq i \leq n$.

Type \mathbf{B}_n . We have $W(\lambda) = Fa = H$. Lusztig has shown that there is an isomorphism

$$\mathcal{U}_m^{\text{od}} \otimes_{\mathbb{C}} \mathcal{U}_{n-m}^{\text{od}} \rightarrow \text{Class}(\mathbf{G}^F, \lambda, a)$$

such that the natural basis $\{[X] \otimes [Y] \mid [X] \in \mathcal{S}_m^{\text{od}}$ and $[Y] \in \mathcal{S}_{n-m}^{\text{od}}\}$ maps onto the series $\mathcal{E}(\mathbf{G}^F, \lambda)$.

The images of the Fourier transforms $\mathcal{R}_0([X]) \otimes \mathcal{R}_0([Y])$ are Lusztig’s almost characters. By [28, 4.23], this bijection may be chosen such that if $[X] \in \tilde{\mathcal{S}}_m^1$ and $[Y] \in \tilde{\mathcal{S}}_{n-m}^1$ have defect 1 then

$$\mathcal{R}_0([X]) \otimes \mathcal{R}_0([Y]) \mapsto \mathcal{R}_{\lambda,a}^{\mathbf{G}}(\rho_{[X]} \boxtimes \rho_{[Y]}).$$

As $[X] \otimes [Y] = \mathcal{R}_0 \mathcal{Q}_0([X]) \otimes \mathcal{R}_0 \mathcal{Q}_0([Y])$, we see that if χ is the image of $[X] \otimes [Y]$ then $f_\chi = \phi_{[X]} \boxtimes \phi_{[Y]}$.

By (iii) of Lemma 4.3, we have $\mathbf{C}_W(Fx) = \mathbf{C}_H(Fx)$. Write $Fx = x_0x_1$ with $x_i \in W_{\mathcal{I}_i}$. If $k_i \geq 1$ is the cycle length of x_i then $k = k_0 + k_1$. Using Theorem 4.7, we thus get the following bound on the character value

$$|f_\chi(Fx)| = |\phi_{[X]}(x_0)| \cdot |\phi_{[Y]}(x_1)| \leq (2^{k_0-1} \cdot k_0!) \cdot (2^{k_1-1} \cdot k_1!) \leq 2^{k-2} \cdot k!.$$

If $\alpha = (\alpha_1, \dots, \alpha_{k_0})$ and $\beta = (\beta_1, \dots, \beta_{k_1})$ are the signed cycle types of x_0 and x_1 respectively then, up to reordering the entries, $\alpha \cup \beta = (\alpha_1, \dots, \alpha_{k_0}, \beta_1, \dots, \beta_{k_1})$ is the signed cycle type of Fx . Thus, there can certainly be at most $\sum_{c=0}^k \binom{k}{c} = 2^k$ terms in the above sum. Putting things together we get the bound $2^{2k-2} \cdot k!$ in this case.

Type \mathbf{C}_n . We have $W(\lambda) = W_{\mathcal{I}_0}^0 W_{\mathcal{I}_1}$ and $Fa = W_{\mathcal{I}_0}^e W_{\mathcal{I}_1}$ for some $e \in \{0, 1\}$. In this case, we have an isomorphism

$$\mathcal{U}_m^{\text{ev},e} \otimes_{\mathbb{C}} \mathcal{U}_{n-m}^{\text{od}} \rightarrow \text{Class}(\mathbf{G}^F, \lambda, a).$$

The natural basis $\{[X] \otimes [Y] \mid [X] \in \mathcal{S}_m^{\text{ev},e}$ and $[Y] \in \mathcal{S}_{n-m}^{\text{od}}\}$ maps onto the series $\mathcal{E}(\mathbf{G}^F, \lambda)$.

By [28, 4.23], this bijection may be chosen such that for any $[X] \in \tilde{\mathcal{S}}_m^0$ and $[Y] \in \tilde{\mathcal{S}}_{n-m}^1$, we have

$$\mathcal{R}_e([X]) \otimes \mathcal{R}_0([Y]) \mapsto \mathcal{R}_{\lambda,a}^{\mathbf{G}}(\text{Res}_{W_{\mathcal{I}_0}^e}^{W_{\mathcal{I}_0}}(\rho_{[X]}) \boxtimes \rho_{[Y]}).$$

As $[X] \otimes [Y] = \mathcal{Q}_e \mathcal{R}_e([X]) \otimes \mathcal{Q}_0 \mathcal{R}_0([Y])$ we see that if χ is the image of $[X] \otimes [Y]$ then $f_\chi = \phi_{[X]} \boxtimes \phi_{[Y]}$.

By (iii) of Lemma 4.3, we have $\mathbf{C}_H(Fx) = \mathbf{C}_W(Fx)$ and $|\mathbf{C}_H(Fx)|/|\mathbf{C}_W(\lambda)(Fx)| \leq |H/W(\lambda)| \leq 2$. Appealing to Theorem 4.7, when X is degenerate, we find, as above, that $|f_\chi(Fx)| \leq 2^{2k-1} \cdot k!$. As we have $|\mathbf{C}_W(Fw) \setminus W/W(\lambda)| \leq 2|\mathbf{C}_W(Fw) \setminus W/H|$, there are at most 2^{k+1} terms in the above sum. Putting things together gives the bound $2^{3k+1} \cdot k!$ in this case.

Type \mathbf{D}_n . We have $W(\lambda) = W_{\mathcal{I}_0}^0 W_{\mathcal{I}_1}^0$. If $\pi_0 = (\bar{e}_1, -\bar{e}_1)$ and $\pi_1 = (\bar{e}_{m+1}, -\bar{e}_{m+1})$ then the coset $Fa \subseteq H$ is either: $W(\lambda)$, $\pi_0 W(\lambda) = W_{\mathcal{I}_0}^1 W_{\mathcal{I}_1}^0$, $\pi_1 W(\lambda) = W_{\mathcal{I}_0}^0 W_{\mathcal{I}_1}^1$, or $\pi_0 \pi_1 W(\lambda)$. These cases are similar to the above. We have $\mathbf{C}_H(Fx) = \mathbf{C}_W(Fx)$ and $|\mathbf{C}_H(Fx)|/|\mathbf{C}_W(\lambda)(Fx)| \leq |H/W(\lambda)| \leq 4$. Arguing similarly, we get that the above sum has at most 2^{k+2} terms, and $|f_\chi(Fx)| \leq 2^{2k} \cdot k!$. Putting things together gives the bound $2^{3k+4} \cdot k!$.

We end with a comment about the final coset $\pi_0 \pi_1 W(\lambda)$. In this case we have an isomorphism $\mathcal{U}_m^{\text{ev},1} \otimes_{\mathbb{C}} \mathcal{U}_{n-m}^{\text{ev},1} \rightarrow \text{Class}(\mathbf{G}^F, \lambda, a)$. Lusztig’s Theorem in this case says that this isomorphism can be chosen such that for any $[X] \in \tilde{\mathcal{S}}_m^0$ and $[Y] \in \tilde{\mathcal{S}}_{n-m}^0$ we have

$$\mathcal{R}_1([X]) \otimes \mathcal{R}_1([Y]) \mapsto \mathcal{R}_{\lambda,a}^{\mathbf{G}}(\text{Res}_{\pi_1 \pi_2 W(\lambda)}^{W_{\mathcal{I}_0} W_{\mathcal{I}_1}}(\rho_{[X]} \boxtimes \rho_{[Y]}));$$

see the discussion in [28, §4.21]. One readily checks that if χ is the image of $[X] \otimes [Y]$ then $f_\chi = \text{Res}_{\pi_1 \pi_2 W(\lambda)}^{W_{\mathcal{I}_0} W_{\mathcal{I}_1}}(\phi_{[X]} \boxtimes \phi_{[Y]})$. □

Corollary 8.2 *Assume G is a symplectic or special orthogonal group and $\chi \in \text{Irr}(G^F)$ is a quadratic unipotent character. Furthermore, let $wF \in WF$ have cycle length $k \geq 1$ and pairwise distinct cycles. Then $|\chi(g)| \leq 2^{3k+4} \cdot k!$ for any regular semisimple element $g \in G^F$ of type wF .*

Proof Recall that being quadratic unipotent means that χ lies in a series $\mathcal{E}(G^F, \lambda, a)$ with $2\lambda \in X$. The statement is thus an immediate consequence of Theorem 8.1 and Lemmas 5.2 and 7.2. □

9 Character degrees

In this section, we prove the following theorem.

Theorem 9.1 *There exists an absolute constant $C > 0$ such that for every finite quasisimple group G of Lie type of rank r and every positive integer D , the number of irreducible characters of G of degree $\leq D$ is at most $D^{C/r}$.*

Taking C large enough, we can ignore any finite number of quasisimple groups G , and thus we may assume that $G = G^F$ for a simple, simply connected algebraic group G of rank r and a Frobenius endomorphism $F : G \rightarrow G$. The Landazuri-Seitz bound [20] implies that the minimal non-trivial character of G has degree at least $|G|^\epsilon$, where ϵ depends only on the rank of G . Therefore, we are justified in assuming that r is as large as we wish, so, in particular, G is of classical type and F is an endomorphism of Steinberg type.

Our proof closely follows the character degree estimates of Liebeck and Shalev [25]. Liebeck and Shalev prove a more precise result [25, Theorem 1.1] than Theorem 9.1 when q is sufficiently large in terms of r and a weaker result [25, Theorem 1.2] for general q .

What is needed to obtain good bounds in high rank for small q is an estimate for the number of unipotent characters of G and certain related groups of bounded degree. This follows in principle from the degree formulas for unipotent characters of classical groups in [28]. We begin with these computations.

Proposition 9.2 *There exists an absolute constant C' such that for every finite quasisimple group G of classical Lie type of rank r and every positive integer D , the number of unipotent characters of G of degree $\leq D$ is at most $D^{C'/r}$.*

Proof For every prime power q , we have by [23, Lemma 4.1(i), (iii)] that

$$\prod_{n=1}^{\infty} (1 - q^{-n}) > \frac{1}{4} \geq q^{-2} \tag{9.1}$$

and

$$\prod_{n=1}^{\infty} (1 + q^{-n}) < 2.4 < q^2. \tag{9.2}$$

(i) If G is of type A_r or 2A_r , then the unipotent characters of G are indexed by sets A of positive integers such that $\rho(A) = r + 1$ in the notation of §2. Denoting the elements of A by $\lambda_1 < \dots < \lambda_m$, we have

$$r + 1 = \sum_{i=1}^m (\lambda_i + 1 - i).$$

The terms $\mu_i := \lambda_i + 1 - i$ in this sum give a partition of $r + 1$, so in particular, $m \leq r + 1$.

Here, $G = \text{SL}_n^\varepsilon(q)$ with $n = r + 1$. The degree d_A of the character with given set A is the absolute value of

$$\frac{\prod_{1 \leq j < i \leq m} ((\varepsilon q)^{\lambda_i} - (\varepsilon q)^{\lambda_j}) \prod_{i=1}^r ((\varepsilon q)^i - 1)}{\prod_{i=1}^m \prod_{j=1}^{\lambda_i} ((\varepsilon q)^j - 1) \prod_{k=2}^{m-1} q^{\binom{k}{2}}}. \tag{9.3}$$

For any fixed j , we have from (9.1) that

$$\prod_{i=j+1}^m |(\varepsilon q)^{\lambda_i} - (\varepsilon q)^{\lambda_j}| > q^{-2} \cdot q^{\sum_{i=j+1}^m \lambda_i},$$

so

$$\prod_{j=1}^{m-1} \prod_{i=j+1}^m |(\varepsilon q)^{\lambda_i} - (\varepsilon q)^{\lambda_j}| > q^{2-2m} \cdot q^{\sum_{1 \leq j < i \leq m} \lambda_i}.$$

Treating the other factors of (9.3) in the same way (and using (9.2) for the products in the denominator), we obtain

$$d_A > q^{-4m} \cdot q^{\sum_{1 \leq j < i \leq m} \lambda_i + \binom{r+1}{2} - \sum_{i=1}^m \binom{\lambda_i+1}{2} - \sum_{k=2}^{m-1} \binom{k}{2}}.$$

As is well-known (see e.g. the proof of [18, Lemma 5.3], the exponent of the second factor on the right-hand side is

$$\frac{1}{2} \left(n^2 - \sum_{i=1}^m \mu_i^2 \right) \geq \frac{1}{2} \left(n^2 - \mu_m \sum_{i=1}^m \mu_i \right) = n(n - \mu_m)/2,$$

and so

$$d_A \geq q^{n(n-\mu_m)/2-4m} \geq q^{n(n-\mu_m)/2-4n}$$

Therefore,

$$n - \mu_m \leq \left(2 \frac{\log d_A}{\log q^n} + 8 \right).$$

As $\mu_m = \lambda_m + 1 - m$ is the largest part of the partition of n associated to A , the number of possibilities for A such that $d_A \leq D$ is at most

$$\sum_{i=1}^{\lfloor 2 \frac{\log D}{\log q^n} + 8 \rfloor} p(i),$$

where p denotes the partition function. As $p(i)$ is sub-exponential in i , when $D \geq q^{n/3}$ this number is $e^{O(\log D / \log q^n)} = D^{O(1)/n}$, yielding a uniform upper bound of the form $D^{C'/r}$ for the number of unipotent characters of degree $\leq D$. However, by [20], for $D \leq q^{n/3}$, there are no non-trivial irreducible characters of degree $\leq D$, and in particular no such unipotent characters.

(ii) The proof for the remaining classical groups follows the same pattern. Let $[X]$ denote an equivalence class of ordered symbols, and let $X = (X^0, X^1)$ be the representative such that $0 \notin X^\cap = X^0 \cap X^1$. Let the elements of X^0 and X^1 respectively form the increasing sequences of non-negative integers $\lambda_1^0 < \dots < \lambda_{|X^0|}^0$ and $\lambda_1^1 < \dots < \lambda_{|X^1|}^1$, so assuming X^0 and X^1 are both non-empty, we have $\lambda_1^0 + \lambda_1^1 > 0$. We note that X^0 and X^1 determine

(possibly improper) partitions $\{\lambda_i^0 + 1 - i\}_i$ and $\{\lambda_j^1 + 1 - j\}_i$ of the ranks $\rho(X^0)$ and $\rho(X^1)$ respectively. More precisely, if $\lambda_i^j > 0$ then $\{\lambda_i^j + 1 - i\}_i$ is a partition of $\rho(X^j)$, but if $\lambda_i^j = 0$ then $\{\lambda_i^j + 1 - i\}_i$ is a sequence of initial zeroes concatenated with a partition of $\rho(X^j)$. Let $n := |X^0| + |X^1|$, and let $\nu_1 \leq \nu_2 \leq \dots \leq \nu_n$ be the sequence obtained by first merging X^0 and X^1 and then sorting, without eliminating repetitions. Thus

$$\nu_1 < \nu_3 < \nu_5 < \dots, \tag{9.4}$$

and

$$0 < \nu_2 < \nu_4 < \dots. \tag{9.5}$$

By (2.1), the rank r of the symbol X is given by

$$r = \sum_i \lambda_i^0 + \sum_j \lambda_j^1 - \left\lfloor \frac{(|X^0| + |X^1| - 1)^2}{4} \right\rfloor = \sum_k \nu_k - \left\lfloor \frac{(n - 1)^2}{4} \right\rfloor = \sum_{k=1}^n (\nu_k - \lfloor (k - 1)/2 \rfloor). \tag{9.6}$$

We have $\nu_k \geq \lfloor (k - 1)/2 \rfloor$ for all k , with strict equality when k is even, so $r \geq \lfloor n/2 \rfloor$. Thus,

$$\begin{aligned} r &= \rho(X^0) + \binom{|X^0|}{2} + \rho(X^1) + \binom{|X^1|}{2} - \left\lfloor \frac{(|X^0| + |X^1| - 1)^2}{4} \right\rfloor \\ &= \rho(X^0) + \rho(X^1) + \begin{cases} (\text{def}(X)^2 - 1)/4, & 2 \nmid n \\ \text{def}(X)^2/4, & 2 | n \end{cases} \\ &\geq \rho(X^0) + \rho(X^1). \end{aligned} \tag{9.7}$$

For every prime power q and every symbol X , there is at least one associated unipotent character of at least one classical group G of rank r over the field \mathbb{F}_q . If $\text{def}(X)$ is odd, we obtain characters of $G = \text{Sp}_{2r}(\mathbb{F}_q)$ and of $G = \text{Spin}_{2r+1}(\mathbb{F}_q)$ in this way. If it is divisible by 4, there is a character of $G = \text{Spin}_{2r}^+(\mathbb{F}_q)$; otherwise, there is a character of $G = \text{Spin}_{2r}^-(\mathbb{F}_q)$. If $X^0 = X^1$, then there are two unipotent characters for $\text{Spin}_{2r}^+(\mathbb{F}_q)$ associated to X ; otherwise, there is only one for each possible G . Moreover, all unipotent characters for groups of type B, C , and D arise in this way for a unique equivalence class of unordered symbols.

The degree d_X of the unipotent character of G associated to the symbol X is (at least)

$$\frac{|G|_{p'} \prod_{1 \leq j < i \leq n^0} (q^{\lambda_i^0} - q^{\lambda_j^0}) \prod_{1 \leq j < i \leq n^1} (q^{\lambda_i^1} - q^{\lambda_j^1}) \prod_{i=1}^{n^0} \prod_{j=1}^{n^1} (q^{\lambda_i^0} + q^{\lambda_j^1})}{2^{\lfloor n/2 \rfloor} \prod_{i=1}^{n^0} \prod_{j=1}^{\lambda_i^0} (q^{2j} - 1) \prod_{i=1}^{n^1} \prod_{j=1}^{\lambda_i^1} (q^{2j} - 1) \prod_{k=1}^{\lfloor (n^0+n^1-2)/2 \rfloor} q^{\binom{n^0+n^1-2k}{2}}}$$

with $n^j := |X^j|$. In terms of the sequence ν_i , this takes the form

$$d_X \geq \frac{|G|_{p'} \prod_{1 \leq j < i \leq n} (q^{\nu_i} \pm q^{\nu_j})}{2^{\lfloor n/2 \rfloor} \prod_{i=1}^n \prod_{j=1}^{\nu_i} (q^{2j} - 1) \prod_{k=1}^{\lfloor (n-2)/2 \rfloor} q^{\binom{n-2k}{2}}}.$$

Note that

$$|G|_{p'} = \begin{cases} \prod_{i=1}^r (q^{2i} - 1) & \text{if } \text{def}(X) \equiv 1 \pmod{2}, \\ (q^r - 1) \prod_{i=1}^{r-1} (q^{2i} - 1) & \text{if } \text{def}(X) \equiv 0 \pmod{4}, \\ (q^r + 1) \prod_{i=1}^{r-1} (q^{2i} - 1) & \text{if } \text{def}(X) \equiv 2 \pmod{4}, \end{cases}$$

so $|G|_{p'} \geq q^{-2} \cdot q^{r^2+r}$ if n is odd and $|G|_{p'} \geq q^{-2} q^{r^2}$ if n is even, again by (9.1).

Reasoning as in case *A*, for cases *B* and *C* (i.e., $n = 2m + 1$ is odd) and using the fact that $n \leq 2r + 1$, we have

$$\begin{aligned} d_X &\geq q^{-2+2-2n-n/2} \cdot q^{r^2+r+\sum_{1 \leq j < i \leq n} v_i - \sum_{i=1}^n v_i(v_i+1) - \sum_{k=1}^{n-1} \binom{n-2k}{2}} \\ &\geq q^{-15r/2} \cdot q^{r^2+r+\sum_{1 \leq j < i \leq n} v_i - \sum_{i=1}^n v_i(v_i+1) - \sum_{k=1}^{n-1} \binom{n-2k}{2}}. \end{aligned}$$

We define

$$Y := \sum_{i=1}^n \left((i-1) - 2 \lfloor \frac{i-1}{2} \rfloor \right) v_i = v_2 + v_4 + \dots + v_{2m} \geq \sum_{i=1}^m i = \frac{m(m+1)}{2},$$

By (9.6), the exponent of the second term on the right-hand side is

$$\begin{aligned} &r^2 + \sum_{i=1}^n v_i - m^2 + \sum_{i=1}^n (i-1)v_i - \sum_{i=1}^n v_i(v_i+1) - \sum_{k=1}^m \binom{n-2k}{2} \\ &= r^2 - m^2 + \sum_{i=1}^n (i-1)v_i - \sum_{i=1}^n v_i^2 - \sum_{k=1}^m \binom{n-2k}{2} \\ &= r^2 - m^2 + Y - \sum_{i=1}^n \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right)^2 + \sum_{i=1}^n \left(\lfloor \frac{i-1}{2} \rfloor \right)^2 - \sum_{k=1}^m \binom{n-2k}{2} \\ &\geq r^2 - \sum_{i=1}^n \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right)^2 + m \\ &\geq r^2 - \sum_{i=1}^n \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right) \cdot \max_i \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right) \\ &= r^2 - r \cdot \max_i \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right). \end{aligned}$$

Here we have used the identity

$$\sum_{i=1}^{2m+1} \left(\lfloor \frac{i-1}{2} \rfloor \right)^2 - \sum_{k=1}^m \binom{2m+1-2k}{2} = 2 \sum_{i=1}^{m-1} i^2 + m^2 - \sum_{i=1}^{m-1} i(2i+1) = \frac{m(m+1)}{2}.$$

Thus,

$$\max_i \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right) \geq r - \frac{\log d_X}{\log q^r} - 7.5. \tag{9.8}$$

Since $|\text{Irr}(G)| \leq q^{C_1 r}$ for some absolute constant C_1 by [11], by enlarging C' (which then covers all small ranks), we may assume $\log d_X / \log q^r \leq r/2 - 8.5$, and so

$$\max_i \left(v_i - \lfloor \frac{i-1}{2} \rfloor \right) \geq \frac{r}{2} + 1. \tag{9.9}$$

By (9.4) and (9.5) and the integrality of the v_i , we have

$$v_{i+2} - \left\lfloor \frac{(i+2)-1}{2} \right\rfloor \geq v_i - \left\lfloor \frac{i-1}{2} \right\rfloor,$$

and so

$$\max_i \left(v_i - \left\lfloor \frac{i-1}{2} \right\rfloor \right) = \max \left(v_{n-1} - \left\lfloor \frac{n-2}{2} \right\rfloor, v_n - \left\lfloor \frac{n-1}{2} \right\rfloor \right).$$

Now, if $\max_i (v_i - \lfloor \frac{i-1}{2} \rfloor)$ is attained at $i = n - 1$, then

$$v_{n-1} - \left\lfloor \frac{n-2}{2} \right\rfloor - 1 \geq r/2 + 1$$

and

$$v_n - \left\lfloor \frac{n-1}{2} \right\rfloor \geq v_{n-1} - \left\lfloor \frac{n-2}{2} \right\rfloor - 1 \geq r/2$$

by (9.9), and this violates (9.6). Hence, $v_i - \lfloor \frac{i-1}{2} \rfloor$ achieves its maximum at only $i = n$, and $v_n > v_{n-1}$, again by (9.9); also,

$$r \geq v_n - \left\lfloor \frac{n-1}{2} \right\rfloor \geq r - \frac{\log d_X}{\log q^r} - 7.5.$$

Applying the Landazuri-Seitz bound as before, we may assume that $d_X \geq q^r$. The rank r' of the symbol, obtained from X by deleting the largest single term v_n , call it X' , is bounded above by $\log d_X / \log q^r + 7.5$, by (9.6) and (9.8). Applying (9.7) to $X' = ((X')^0, (X')^1)$, we see that

$$\rho((X')^0) + \rho((X')^1) \leq \log d_X / \log q^r + 7.5 \leq x + 7.5 \tag{9.10}$$

if $d_X \leq q^{rx}$ with $x \geq 0$. In fact, we can show that such symbols X satisfy

$$|X^0| + |X^1| < 3x + 22.5. \tag{9.11}$$

Indeed, without loss we may assume that $\mu_1 \geq 1$, so the sequence $\{\mu_j + 1 - j\}$ of $|T'|$ integers is a proper partition of $\rho((X')^1)$, and so $|(X')^1| \leq \rho((X')^1) \leq x + 7.5$ by (9.10). Applying (9.7) to X' we have $(|(X')^0| - |(X')^1|)^2 \leq 4r' + 1$, and so

$$|(X')^0| - |(X')^1| \leq \sqrt{4r' + 1} \leq \sqrt{4(x + 7.5) + 1} < x + 6.5.$$

Hence $|(X')^0| < 2x + 14$, and $|X^0| + |X^1| = |(X')^0| + |(X')^1| + 1 < 3x + 22.5$, as stated.

By (9.11), even when $\lambda_1^j = 0$, the number of zero entries in the sequence $\{\lambda_i^j + 1 - i\}$ is at most $|(X')^j| < 3x + 22.5$. Now, counting the number of (possibly improper) partitions $\{\lambda_i^j + 1 - i\}$ of $\rho((X')^j)$ and using (9.10), we see that the number of possibilities for the symbol X with $d_X < q^{rx}$ is bounded above by

$$(3x + 22.5) \sum_{i=1}^{\lfloor x+7.5 \rfloor} p(i)^2,$$

an exponential in x for $x \geq 1$, proving the proposition for types B_r and C_r .

(iii) For types D_r and 2D_r , that is, when $n = 2m$ is even, we have $n \leq 2r$, and

$$\begin{aligned} d_X &\geq q^{-2+2-2n-n/2} \cdot q^{r^2+\sum_{1 \leq j < i \leq n} v_i - \sum_{i=1}^n v_i(v_i+1) - \sum_{k=1}^{\frac{n-2}{2}} \binom{n-2k}{2}} \\ &\geq q^{-5r} \cdot q^{r^2+\sum_{1 \leq j < i \leq n} v_i - \sum_{i=1}^n v_i(v_i+1) - \sum_{k=1}^{\frac{n-2}{2}} \binom{n-2k}{2}}. \end{aligned}$$

The exponent of the second term on the right hand side is therefore

$$r^2 + \sum_{i=1}^n (i-1)v_i - \sum_{i=1}^n v_i(v_i+1) - \sum_{k=1}^{m-1} \binom{n-2k}{2}$$

$$\begin{aligned}
 &= r^2 + \sum_{i=1}^n (i - 2)v_i - \sum_{i=1}^n v_i^2 - \sum_{k=1}^{m-1} \binom{n - 2k}{2} \\
 &= r^2 - Y - \sum_{i=1}^n \left(v_i - \left\lfloor \frac{i - 1}{2} \right\rfloor \right)^2 + \sum_{i=1}^n \left(\left\lfloor \frac{i - 1}{2} \right\rfloor \right)^2 - \sum_{k=1}^{m-1} \binom{n - 2k}{2} \\
 &\geq r^2 - \sum_{i=1}^n \left(v_i - \left\lfloor \frac{i - 1}{2} \right\rfloor \right)^2 - r \\
 &\geq r^2 - \sum_{i=1}^n \left(v_i - \left\lfloor \frac{i - 1}{2} \right\rfloor \right) \cdot \max_i \left(v_i - \left\lfloor \frac{i - 1}{2} \right\rfloor \right) - r \\
 &= r^2 - r \cdot \max_i \left(v_i - \left\lfloor \frac{i - 1}{2} \right\rfloor \right) - r.
 \end{aligned}$$

Here we have used the inequality

$$\begin{aligned}
 Y &:= \sum_{i=1}^n (2 \lfloor \frac{i-1}{2} \rfloor - (i - 2)) v_i = v_1 + v_3 + \dots + v_{2m-1} \\
 &= \sum_{i=1}^n 2 \lfloor \frac{i-1}{2} \rfloor v_i + \binom{m}{2} \leq r + \binom{m}{2}
 \end{aligned}$$

and the identity

$$\sum_{i=1}^{2m} \left(\left\lfloor \frac{i - 1}{2} \right\rfloor \right)^2 - \sum_{k=1}^{m-1} \binom{2m - 2k}{2} = 2 \sum_{i=1}^{m-1} i^2 - \sum_{i=1}^{m-1} i(2i - 1) = \binom{m}{2}.$$

The argument finishes as before. □

Proof of Theorem 9.1 Let $G^* = (\mathbf{G}^*)^{F^*}$ denote the dual group of G . We partition the irreducible characters of G into rational Lusztig series $\mathcal{E}(s)$, indexed by conjugacy classes of semisimple conjugacy classes (s) of G^* . There is a bijection between the elements of $\mathcal{E}(s)$ and unipotent characters of $\mathbf{C}_{G^*}(s)$; this correspondence multiplies degrees by $|G^*|_{p'}/|\mathbf{C}_{G^*}(s)|_{p'}$.

We restate [25, Lemma 3.2] in a form more convenient for our purposes. Since the ratio n/r between the dimension n of the natural module and the rank of G is bounded between 1 and 3, and since the constants d and d' in [25] are absolute, if δ is greater than some absolute constant δ_0 , then the number of semisimple conjugacy classes (s) with $|G^*|_{p'}/|\mathbf{C}_{G^*}(s)|_{p'} \leq q^{\delta r}$ is less than $q^{\delta A}$ for some absolute constant A . Moreover, $\mathbf{C}_{G^*}(s)$ contains a factor, the *large factor*, which is classical of rank $r' \geq r - B\delta$ for some absolute constant B , and this large factor is $A_{r'}(q)$ or ${}^2A_{r'}(q)$ when \mathbf{G} is of type A .

For $D < q^{r/3}$, there is only one irreducible character, by [20]. Hence we may assume $D \geq q^{r/3}$. Enlarging C if necessary, we may assume that $D = q^{\delta r}$ with $\delta \geq \max(\delta_0, 1/2)$. By [11], $|\text{Irr}(G)| \leq q^{C_1 r}$ for some absolute constant C_1 , so the result follows if $\delta C \geq C_1 r$. Again enlarging C if necessary, we may assume without loss of generality that $\delta < r/2B$.

If χ is an irreducible character of degree $\leq D$, then it belongs to the Lusztig series $\mathcal{E}(s)$ for some s with $|G|_{p'}/|\mathbf{C}_{G^*}(s)|_{p'} \leq D$. The number of such semisimple classes s is bounded above by $q^{\delta A}$. Following the proof of [25, Lemma 3.4], note that for each s , $\mathbf{C}_{G^*}(s)$ contains a subgroup $\mathbf{C}_{G^*}(s)^\circ$ which is the group of F^* -fixed points of the connected reductive algebraic group $\mathbf{C}_{G^*}(s)^\circ$. If \mathbf{G} is of type $B, C,$ or D , the quotient group $\mathbf{C}_{G^*}(s)/\mathbf{C}_{G^*}(s)^\circ$ has order ≤ 4 . Suppose \mathbf{G} is of type A and $\mathbf{C}_{G^*}(s)^\circ$ is a proper subgroup of $\mathbf{C}_{G^*}(s)$. Lifting s to an element \hat{s} of $\text{GL}_n^E(q)$, we see that every eigenvalue of \hat{s} has multiplicity $\leq n/2$, but this contradicts the existence of the large factor of $\mathbf{C}_{G^*}(s)$ which is of type $A_{r'}(q)$ or ${}^2A_{r'}(q)$ with $r' > r/2$. So we have $\mathbf{C}_{G^*}(s) = \mathbf{C}_{G^*}(s)^\circ$ for type A . Thus, there are at most 4 unipotent characters

of $C_{G^*}(s)$ of degree $\leq D$ for each unipotent character of $C_{G^*}(s)^\circ$ of degree $\leq D$. Taking F^* -fixed points of the derived group of $C_{G^*}(s)^\circ$ we obtain a subgroup whose unipotent characters correspond to those of $C_{G^*}(s)$, and this subgroup is a product of classical groups whose ranks sum up to at most r and at least one of which, the large factor, has rank r' at least $r - B\delta \geq r/2$. The total number of unipotent characters of the product of all the factors other than the large factor can therefore be bounded above by $q^{C_1 B \delta}$ by [11]. The number of unipotent characters of the large factor of degree $\leq D = q^{\delta r}$ is bounded above by $D^{C'/r'} \leq q^{2\delta C'}$, by Proposition 9.2. Hence, the number of unipotent characters of degree $\leq D$ of $C_{G^*}(s)^\circ$ is bounded by $q^{(BC_1+2C')\delta}$, and the number for $C_{G^*}(s)$ is likewise bounded by an exponential in δ . Thus the number of characters of degree at most D is bounded by $q^{C_2\delta}$ for some absolute constant C_2 , and the theorem follows by taking $C \geq C_2$. \square

For later use, we prove the following related statement:

Proposition 9.3 *Let $n \geq 4$ and $j \geq 1$ be integers and let q be any prime power. Let $G = \mathbf{G}^F$ be one of the groups $\mathrm{Sp}_{2n}(q)$, $\mathrm{SO}_{2n+1}(q)$, or $\mathrm{SO}_{2n}^\epsilon(q)$, and suppose that $\chi \in \mathrm{Irr}(G)$ has degree*

$$\chi(1) \leq \min \left(q^{nj}, q^{(n^2-n)/2-4} \right).$$

Then there is an F -stable Levi subgroup $\mathbf{L} = \mathbf{L}_1 \times \mathbf{L}_2$ of \mathbf{G} , possibly equal to \mathbf{G} , such that the following statements hold:

- (i) \mathbf{L}_1 is F -stable, of the same type as \mathbf{G} , and of rank $n - m \geq n - 2j$.
- (ii) \mathbf{L}_2 is F -stable, of type GL_m with $m \leq 2j$.
- (iii) $\chi = \pm R_{\mathbf{L}}^{\mathbf{G}}(\varphi_1 \boxtimes \varphi_2)$, where $\varphi_1 \in \mathrm{Irr}(\mathbf{L}_1^F)$ is a unipotent or quadratic unipotent character, and $\varphi_2 \in \mathrm{Irr}(\mathbf{L}_2^F)$. Moreover, if $j \leq n/2$, then $\varphi_1(1) \leq q^{(n-m)j}$.

Proof View $G = \mathrm{Sp}(V)$ or $\mathrm{SO}(V)$, where $V = \mathbb{F}_q^{2n}, \mathbb{F}_q^{2n+1}$, or \mathbb{F}_q^{2n} . Then we can identify the dual group $G^* = (\mathbf{G}^*)^{F^*}$ with $\mathrm{SO}(V^*) \cong \mathrm{SO}_{2n+1}(q)$, $\mathrm{Sp}(V^*) \cong \mathrm{Sp}_{2n}(q)$, or $\mathrm{SO}(V^*) \cong \mathrm{SO}(V) \cong \mathrm{SO}_{2n}(q)$, respectively. Let $\mathcal{E}(G, (s))$ be the rational Lusztig series that contains χ , where $s \in G^*$ is semisimple. If $s^2 = \mathrm{id}_{V^*}$, then we are done by choosing $\mathbf{L} = \mathbf{G}$. Otherwise we can decompose $V^* = V_1^* \oplus V_2^*$, where $s^2 - \mathrm{id}_{V^*}$ is zero on V_1^* , and invertible on its orthogonal complement $V_2^* \neq 0$. Since s is semisimple, we see that $C_{G^*}(s)$ is contained in a proper F^* -stable Levi subgroup $\mathbf{L}^* = \mathbf{L}_1^* \times \mathbf{L}_2^*$ of \mathbf{G}^* , where $\mathbf{L}_2^* \cong \mathrm{GL}_m$, with $m := \dim(V_2^*)/2 \leq n$, and \mathbf{L}_1^* , of the same type as of \mathbf{G}^* , are both F^* -stable. Let $\mathbf{L} = \mathbf{L}_1 \times \mathbf{L}_2$ denote the Levi subgroup of \mathbf{G} dual to \mathbf{L}^* , where \mathbf{L}_1 is F -stable and of the same type as of \mathbf{G} , and $\mathbf{L}_2 \cong \mathrm{GL}_m$. By [8, Thm. 11.4.3], $\varepsilon_{\mathbf{G}} \varepsilon_{\mathbf{L}} R_{\mathbf{L}}^{\mathbf{G}}$ yields a bijection between $\mathcal{E}(\mathbf{L}^F, (s))$ and $\mathcal{E}(G, (s))$, which implies (iii); in particular,

$$\chi(1) = \frac{|G|_{p'}}{|\mathbf{L}^F|_{p'}} \varphi_1(1)\varphi_2(1), \tag{9.12}$$

if p denotes the unique prime divisor of q .

Using (9.1) and (9.2), one readily checks that

$$\frac{|G|_{p'}}{|\mathbf{L}^F|_{p'}} > \begin{cases} q^{(4nm-3m^2-m-8)/2}, & \mathbf{G} = \mathrm{SO}_{2n}, \\ q^{(4nm-3m^2+m-8)/2}, & \mathbf{G} = \mathrm{Sp}_{2n}, \mathrm{SO}_{2n+1}. \end{cases}$$

In particular, if $m = n$ then $\chi(1) > q^{(n^2-n)/2-4}$ by (9.12), a contradiction. Assume now that $m \leq n - 1$, but $m \geq 2j + 1$. Then $4n - 3m - 1 \geq n + 2$, whence

$$4nm - 3m^2 - m - 8 = m(4n - 3m - 1) - 8 \geq (2j + 1)(n + 2) - 8$$

$$= 2nj + n + 2(2j + 1) - 8 \geq 2nj + 1,$$

and so $\chi(1) > q^{nj}$, again a contradiction. Thus $m \leq 2j$.

To show $\varphi_1(1) \leq q^{(n-m)j}$, it suffices by (9.12) to check that $|G|_{p'}/|\mathbf{L}^F|_{p'} \geq q^{mj}$. This is obvious if $m = 0$. If $1 \leq m \leq 3$, then $j \leq n/2 \leq 2n - 6$ implies that $4nm - 3m^2 - m - 8 \leq 2mj$. If $m \geq 4$, then

$$4nm - 3m^2 - m - 8 = m(4n - 3m - 1) - 8 \geq m(n + 2) - 8 \geq mn \geq 2mj,$$

and so we are done. □

10 Applications to asymptotic variants of Thompson’s conjecture

10.1 Type A

Recall $\mathrm{SL}_n^\varepsilon(q)$ denotes $\mathrm{SL}_n(q)$ when $\varepsilon = +$, and $\mathrm{SU}_n(\mathbb{F}_{q^2})$ when $\varepsilon = -$, and similarly for $\mathrm{GL}_n^\varepsilon(q)$.

Theorem 10.1 *For all $k \in \mathbb{Z}_{\geq 1}$, there exists an explicit constant $B = B(k) > 0$ such that the following statement holds for all $n \in \mathbb{Z}_{\geq 1}$ and all prime powers q . Suppose $G = \mathrm{SL}_n^\varepsilon(q)$ for some $\varepsilon = \pm$ and $g \in G$ is a regular semisimple element whose characteristic polynomial on the natural module of G is a product of k pairwise distinct irreducible polynomials. Then $g^G \cdot g^G$ contains every element $x \in G$ of support $\geq B$.*

Proof (a) Embed G in $\tilde{G} := \mathrm{GL}_n^\varepsilon(q)$. Since the support of an element of \tilde{G} is at most n , by enlarging B , we are free to make $n \geq k$ as large as we wish.

Note that the element g is regular semisimple, and $T := \mathbf{C}_{\tilde{G}}(g)$ is a maximal torus, so of order at most $(q + 1)^n$. Moreover, the image of T under the determinant map is the same as that of \tilde{G} . Hence the conjugacy class of g in G is the same as its class in \tilde{G} . Let $x \in G$. To show that $x \in g^G \cdot g^G$, it suffices to prove that

$$\sum_{\chi \in \mathrm{Irr}(\tilde{G})} \frac{\chi(g)^2 \bar{\chi}(x)}{\chi(1)} \neq 0.$$

As $\det(g) = \det(x) = 1$, for every character χ of degree 1 we have $\chi(g) = \chi(g)^2 \bar{\chi}(x) = 1$. Therefore, it suffices to prove that

$$\sum_{\{\chi \in \mathrm{Irr}(\tilde{G}) \mid \chi(1) > 1\}} \frac{|\chi(g)|^2 |\chi(x)|}{\chi(1)} < q - \varepsilon. \tag{10.1}$$

(b) For any fixed $\varepsilon > 0$, choosing B sufficiently large, the contribution of characters $\chi \in \mathrm{Irr}(\tilde{G})$ satisfying $\chi(1) \geq q^{\varepsilon n^2}$ to (10.1) is $o(1)$. Indeed, consider any such character χ and any irreducible constituent ψ of $\chi|_G$. Since $\tilde{G}/G \cong C_{q-\varepsilon}$, by Clifford’s theorem we have $\chi|_G = \psi_1 + \dots + \psi_t$, where $\psi_1 = \psi, \dots, \psi_t$ are distinct \tilde{G} -conjugates of ψ , and $t \mid (q - \varepsilon)$. By [21, Theorem 5.5],

$$|\psi_i(x)| \leq \psi_i(1)^{1-\sigma B/n} = (\chi(1)/t)^{1-\sigma B/n}$$

for some absolute constant $\sigma > 0$, and so $|\chi(x)| \leq t(\chi(1)/t)^{1-\sigma B/n}$. As $\chi(1) \geq (q + 1)^2 \geq t^2$, we obtain

$$|\chi(x)/\chi(1)| \leq \chi(1)^{-\sigma B/2n} \leq q^{-\varepsilon \sigma B n/2}.$$

Since $|T| \leq (q + 1)^n < q^{2n}$, it follows that the contribution of all these characters to (10.1) is at most

$$q^{-\varepsilon\sigma Bn/2} \sum_{\chi} |\chi(g)|^2 \leq q^{-\varepsilon\sigma Bn/2} |T| < q^{2n(1-\varepsilon\sigma B/4)}$$

which is $o(1)$ when B is large enough.

(c) Now let j denote the level of $\chi \in \text{Irr}(\tilde{G})$, as defined in [18]. Assuming $\chi(1) > 1$, we have $j > 0$. If $j \geq n/2$, then $\chi(1) \geq q^{n^2/4-2}$ by [18, Theorem 1.2(ii)], and, as shown in (b), the contribution of all such characters to the left hand side of (10.1) is $o(1)$. Hence it remains to consider the characters χ with

$$j < n/2;$$

any such character is irreducible over G , see [18, Corollary 8.6]. Up to a linear factor, we may assume that χ has true level j . By [18, Theorem 3.9], any such character χ is of the form

$$\chi = \pm R_L^{\tilde{G}}(\varphi \boxtimes \psi),$$

where $L = \text{GL}_{n-m}^{\varepsilon}(q) \times \text{GL}_m^{\varepsilon}(q)$ is a (possibly non-proper) Levi subgroup of \tilde{G} with $0 \leq m < n$, $\varphi = \varphi^{\lambda}$ is the unipotent character of $\text{GL}_{n-m}^{\varepsilon}(q)$ labeled by a partition $\lambda \vdash (n-m)$ with largest part $\lambda_1 = n - j$, so, in particular,

$$m \leq j, \tag{10.2}$$

and $\psi \in \text{Irr}(\text{GL}_m^{\varepsilon}(q))$ when $m > 0$. Moreover, the total number of characters of \tilde{G} of true level j is $|\text{Irr}(\text{GL}_j^{\varepsilon}(q))|$, which is shown in [11, Propositions 3.5, 3.9] to be at most $9q^j$.

Since χ has level j , $\chi(1) \geq q^{j(n-j)-1} > q^{nj/3}$ by [18, Theorem 1.2(i)]. For these characters χ , $\text{supp}(x) \geq B$ implies by [21, Theorem 5.5] that

$$|\chi(x)|/\chi(1) < q^{-\sigma Bj/3}. \tag{10.3}$$

As g is regular semisimple, the Steinberg character St_G of G takes value ± 1 at x . Applying [8, Cor. 10.2.10] we have

$$\chi(g) = \pm(\text{St}_G \cdot \chi)(g) = \pm \text{Ind}_L^{\tilde{G}}(\text{St}_L \cdot \varphi)(g). \tag{10.4}$$

Note that if $V = \mathbb{F}_q^n$ denotes the natural module of \tilde{G} (endowed with a Hermitian form when $\varepsilon = -$), then the L -module V is a direct (orthogonal when $\varepsilon = -$) sum of two non-isomorphic irreducible modules $V_1 := \mathbb{F}_q^{n-m}$ and $V_2 := \mathbb{F}_q^m$, with $m \leq j < n/2$, see (10.2). In particular, if $y \in \mathbf{N}_G(L)$, then y preserves each of V_1 and V_2 , and thus $\mathbf{N}_G(L) = L$.

Now we count the number N of elements $y \in G$ such that $y^{-1}gy \in L$, i.e. $g \in yLy^{-1}$. Then g acts on each of the subspaces yV_1 and yV_2 . On the other hand, the decomposition $p_V(g) = \prod_{i=1}^k f_i(X)$ leads to a decomposition $V = \bigoplus_{i=1}^k U_i$, where $p_{U_i}(g) = f_i(X)$, and each U_i is a minimal $\langle g \rangle$ -invariant, non-degenerate if $\varepsilon = -$, subspace. Moreover, the $\langle g \rangle$ -modules U_i are pairwise non-isomorphic. Hence (yV_1, yV_2) is uniquely determined by choosing a subset of $\{U_1, \dots, U_k\}$ (so that yV_1 is the sum over this subset and yV_2 is the sum over the complement). Thus the total number of possibilities for $yLy^{-1} = \mathbf{N}_G(yV_1, yV_2)$ is at most 2^k . On the other hand, $yLy^{-1} = y'Ly'^{-1}$ if and only if $y^{-1}y' \in \mathbf{N}_G(L) = L$. Hence

$$N \leq 2^k |L|. \tag{10.5}$$

Suppose $y^{-1}gy = \text{diag}(g_1, g_2) \in L$, with $g_1 \in L_1 := \text{GL}_{n-m}^{\varepsilon}(q)$ and $g_2 \in L_2 := \text{GL}_m^{\varepsilon}(q)$. Let k_i denote the number of irreducible factors of the characteristic polynomial of

g_i on the natural module for L_i . Then $k_1 + k_2 \leq k$. Since φ is unipotent and g_1 is regular semisimple, we have

$$|\varphi(g_1)| \leq 2^{k_1-1} \cdot k_1!$$

by Corollary 7.5. On the other hand, when $m > 0$, (10.2) and Corollary 7.6 show that

$$|\psi(g_2)| \leq k_2! \cdot j^{k_2}.$$

As $y^{-1}gy$ is regular semisimple in L , $|\text{St}_L(y^{-1}gy)| = 1$. Hence

$$|(\text{St}_L \cdot (\varphi \boxtimes \psi))(y^{-1}gy)| \leq 2^{k-1} \cdot k! \cdot j^{k-1}.$$

It now follows from (10.4) and (10.5) that

$$|\chi(g)| = \left| \frac{1}{|L|} \sum_{y \in G: y^{-1}gy \in L} \pm(\text{St}_L \cdot \varphi)(y^{-1}gy) \right| \leq 2^{2k-1} \cdot k! \cdot j^{k-1}.$$

With (10.3), this shows that the total contribution of characters of a fixed true level $1 \leq j < n/2$ to (10.1) is at most

$$9q^j \cdot q^{-\sigma B j/3} \cdot (2^{2k-1} \cdot k! \cdot j^{k-1})^2 < A(k)/q_1^j,$$

where $A(k) = 9 \cdot (2^{2k-1} \cdot k!)^2$ and $q_1 := q^{\sigma B/3-k}$ (here we use the estimate $q^j \geq 2^j \geq j^2$). Note that

$$\sum_{j=1}^{\infty} 1/q_1^j = 1/(q_1 - 1) \tag{10.6}$$

is $o(1)$ when B is large enough. Hence, the total contribution of characters χ , of level at least 1 and less than $n/2$, to (10.1), is less than $(q - \varepsilon)A(k)/(q_1 - 1) = o(q - \varepsilon)$, and the theorem follows. □

10.2 Types BCD

Recall the involution $f \mapsto f^\vee$ on the set \mathcal{F}_q^* of monic irreducible polynomial $f \in \mathbb{F}_q[t]$ with $f(0) \neq 0$: if $\deg(f) = m$ then $f^\vee(t) = t^m f(1/t)/f(0)$ (equivalently, $\lambda \in \overline{\mathbb{F}_q}$ is a root of f^\vee if and only if $1/\lambda$ is a root of f). In the following theorem, the condition that the regular semisimple element $g \in G$ has pairwise distinct cycles implies that the characteristic polynomial of g on the natural $\mathbb{F}_q G$ -module is of the form

$$(t - 1)^a (t + 1)^b \prod_{i=1}^c f_i f_i^\vee \prod_{j=1}^d h_j,$$

where $a, b, c, d \geq 0$, $a, b \leq 2$, $f_i^\vee \neq f_i \in \mathcal{F}_q^*$, $h_j = h_j^\vee \in \mathcal{F}_q^*$, $\deg(f_i) = m_i$, $\deg(h_j) = n_j$, $m_1 > \dots > m_c$, and $n_1 > \dots > n_d$ (and $c + d \leq k \leq c + d + 2$ for the cycle length k). Note that if the irreducible factors of the characteristic polynomial of g have pairwise distinct degrees, then g has pairwise distinct cycles.

Theorem 10.2 *For all $k \in \mathbb{Z}_{\geq 1}$, there exists $B > 0$ such that the following statement holds for all $n \in \mathbb{Z}_{\geq 1}$ and all prime powers q . If $G = \text{Sp}_{2n}(q)$, $\text{SO}_{2n+1}(q)$, or $\text{SO}_{2n}^\pm(q)$, and $g \in G$ is a regular semisimple element with cycle length k and pairwise distinct cycles, then $g^G \cdot g^G$ contains every element $x \in [G, G]$ of support $\geq B$.*

Proof (a) Enlarging B , we are free to make $n \geq \max(k, 5)$ as large as we wish. Write $G = \mathbf{G}^F$ for a corresponding simple algebraic group of type Sp or SO . Then $\mathbf{C}_G(g)$ is a maximal torus, so, using the well-known structure of centralizers of semisimple elements in the finite group G , we see that $T := \mathbf{C}_G(g)$ has order at most $2(q + 1)^n$.

Let $x \in [G, G]$. Since the linear characters of G take value 1 at x , to show that $x \in g^G \cdot g^G$, it suffices to prove that

$$\sum_{\{\chi \in \mathrm{Irr}(G) \mid \chi(1) > 1\}} \frac{|\chi(g)|^2 |\chi(x)|}{\chi(1)} < 1 \leq |G/[G, G]|. \tag{10.7}$$

(b) For any fixed $\epsilon > 0$, choosing B sufficiently large, the contribution of characters $\chi \in \mathrm{Irr}(G)$ satisfying $\chi(1) \geq q^{\epsilon n^2}$ to (10.7) is $o(1)$. Indeed, for any such character χ , by [21, Theorem 5.5] we have

$$|\chi(x)/\chi(1)| \leq \chi(1)^{-\sigma B/n} \leq q^{-\epsilon \sigma B n}$$

for some absolute constant $\sigma > 0$. Since $|T| \leq 2(q + 1)^n < q^{2n}$, it follows that the contribution of all these characters to (10.7) is at most

$$q^{-\epsilon \sigma B n} \sum_{\chi} |\chi(g)|^2 \leq q^{-\epsilon \sigma B n} |T| < q^{2n(1-\epsilon \sigma B/2)}$$

which is $o(1)$ when B is large enough.

(c) Now we consider any $\chi \in \mathrm{Irr}(G)$ with $1 < \chi(1) < q^{(n^2-4n)/4}$. By the Landazuri-Seitz bound [20], we have $\chi(1) > q^{n/2}$. Let $j \in \mathbb{Z}_{\geq 1}$ be the unique integer such that $q^{n(j-1)} \leq \chi(1) < q^{nj}$, and note that

$$q^{nj/2} \leq \chi(1) < \min\left(q^{nj}, q^{(n^2-n)/2-4}\right), \quad j < n/4.$$

By Proposition 9.3, there is an F -stable Levi subgroup $\mathbf{L} = \mathbf{L}_1 \times \mathbf{L}_2$ of \mathbf{G} , possibly equal to \mathbf{G} , such that the following statements hold:

- (α) \mathbf{L}_1 is F -stable, of the same type as of \mathbf{G} , and of rank $n - m \geq n - 2j$.
- (β) \mathbf{L}_2 is F -stable, of type GL_m with $m \leq 2j$.
- (γ) $\chi = \pm R_{\mathbf{L}}^{\mathbf{G}}(\varphi_1 \boxtimes \varphi_2)$, where $\varphi_1 \in \mathrm{Irr}(\mathbf{L}_1^F)$ is a unipotent or quadratic unipotent character, and $\varphi_2 \in \mathrm{Irr}(\mathbf{L}_2^F)$.

Moreover, by Theorem 9.1, there is an absolute constant C such that the total number N_j of characters of G of degree $\leq q^{nj}$ is at most

$$N_j \leq q^{Cj}. \tag{10.8}$$

Since $\chi(1) \geq q^{nj/2}$, $\mathrm{supp}(x) \geq B$ implies by [21, Theorem 5.5] that

$$|\chi(x)|/\chi(1) < q^{-\sigma B j/2}. \tag{10.9}$$

Next we bound $|\chi(g)|$, again using (10.4). Let $V = \mathbb{F}_q^d$ denote the natural module of G endowed with a symplectic or quadratic form, $d = 2n$ or $2n + 1$, and let $L := \mathbf{L}^F$. Then the L -module V is an orthogonal sum of two non-degenerate L -invariant subspaces $V_1 := \mathbb{F}_q^{d-2m}$ and $V_2 := \mathbb{F}_q^{2m}$, with

$$2m \leq 4j < n \leq d - 2m.$$

Furthermore, V_1 is the natural, irreducible module of dimension $d - 2m$ for $L_1 := \mathbf{L}_1^F$, with \mathbf{L}_1 of the same type as of \mathbf{G} , and L_1 acts trivially on V_2 . Next, $L_2 := \mathbf{L}_2^F \cong \mathrm{GL}_m(q)$ or

$GU_m(q)$, with V_1 a minimal L_2 -invariant non-degenerate subspace, and L_2 acts trivially on V_1 . In particular, if $y \in \mathbf{N}_G(L)$, then y preserves each of V_1 and V_2 , and thus $\mathbf{N}_G(L) = L$.

Now we count the number N of elements $y \in G$ such that $y^{-1}gy \in L$, i.e. $g \in yLy^{-1}$. Then g acts on each of the subspaces yV_1 and yV_2 . On the other hand, since g has cycle length k with pairwise distinct cycles, V admits an orthogonal decomposition $V = \bigoplus_{i=1}^k U_i$, where each U_i is a minimal $\langle g \rangle$ -invariant non-degenerate subspace. Moreover, the $\langle g \rangle$ -modules U_i are pairwise non-isomorphic. Hence (yV_1, yV_2) is uniquely determined by choosing a subset of $\{U_1, \dots, U_k\}$ (so that yV_1 is the sum over this subset and yV_2 is the sum over the complement). Thus the total number of possibilities for $yLy^{-1} = \mathbf{N}_G(yV_1, yV_2)$ is at most 2^k . On the other hand, $yLy^{-1} = y'Ly'^{-1}$ if and only if $y^{-1}y' \in \mathbf{N}_G(L) = L$. Hence

$$N \leq 2^k |L|. \tag{10.10}$$

Suppose $y^{-1}gy = \text{diag}(g_1, g_2) \in L$, with $g_1 \in L_1$ and $g_2 \in L_2 = \text{GL}_m^\pm(q)$. Let k_i denote the cycle length of g_i . Then $k_1 + k_2 \leq k$. Since φ_1 is quadratic unipotent and g_1 is regular semisimple, we have

$$|\varphi_1(g_1)| \leq 2^{3k_1+4} \cdot k_1!$$

by Corollary 8.2. On the other hand, when $m > 0$, the statement (β) and Corollary 7.6 show that

$$|\varphi_2(g_2)| \leq k_2! \cdot (2j)^{k_2}.$$

As $y^{-1}gy$ is regular semisimple in L , $|\text{St}_L(y^{-1}gy)| = 1$. Hence

$$|(\text{St}_L \cdot (\varphi_1 \boxtimes \varphi_2))(y^{-1}gy)| \leq 2^{3k+4} \cdot k! \cdot j^{k-1}.$$

It now follows from (10.4) and (10.10) that

$$|\chi(g)| = \left| \frac{1}{|L|} \sum_{y \in G: y^{-1}gy \in L} \pm(\text{St}_L \cdot \varphi)(y^{-1}gy) \right| \leq 2^{4k+4} \cdot k! \cdot j^{k-1}.$$

With (10.9), this shows that the total contribution of characters of degree satisfying $q^{n(j-1)} \leq \chi(1) < q^{nj}$ to (10.7) is at most

$$q^{Cj} \cdot q^{-\sigma B j/2} \cdot (2^{4k+4} \cdot k! \cdot j^{k-1})^2 < A(k)/q_1^j,$$

where $A(k) = (2^{4k+4} \cdot k!)^2$ and $q_1 := q^{\sigma B/2 - C - k + 1}$ (here we again use $q^j \geq j^2$). Recalling (10.6), we conclude that the total contribution to (10.7) of characters χ , of degree at least 2 and less than $q^{(n^2 - 4n)/4}$, is less than $A(k)/(q_1 - 1) = o(1)$, and hence the theorem follows. □

Proof of Corollary 2. Recall that $\chi(x) = 1$ and $|\chi(g)|^2 = 1$ for any irreducible character $\chi \in \text{Irr}(\tilde{G})$ of degree 1. Hence, the inequalities (10.1) and (10.7) established in the proof of Theorem 10.1, respectively Theorem 10.2, show that

$$\sum_{\chi \in \text{Irr}(\tilde{G})} \frac{|\chi(g)|^2 \overline{\chi(x)}}{\chi(1)} > 0.$$

Hence the statement follows from (1.2). □

10.3 Another result for SL

For any positive interger k , let \underline{a} denote a fixed increasing sequence $a_1 < \dots < a_k$ of positive integers. By an \underline{a} -flag in an \mathbb{F}_q -vector space V , we mean a flag

$$V_1 \subset \dots \subset V_k \subset V$$

of \mathbb{F}_q -subspaces such that $\dim V_i = a_i$ for $1 \leq i \leq k$. The number of \underline{a} -flags is

$$F_{\underline{a}}(\dim V) := \frac{\prod_{j=1}^{a_{k+1}} (q^j - 1)}{\prod_{i=0}^k \prod_{j=1}^{a_{i+1} - a_i} (q^j - 1)},$$

where we define $a_0 := 0$ and $a_{k+1} := \dim V$. As

$$\prod_{j=1}^{\infty} (1 - q^{-j}) > \frac{1}{4}$$

by [23, Lemma 4.1], we have

$$\frac{q^{d_{\underline{a}}(N)}}{4} < F_{\underline{a}}(N) < 4^k q^{d_{\underline{a}}(N)}, \tag{10.11}$$

where

$$\begin{aligned} d_{\underline{a}}(N) &:= \sum_{0 \leq i < j \leq k} (a_{i+1} - a_i)(a_{j+1} - a_j) \\ &= a_k(N - a_k) + \sum_{0 \leq i < j \leq k-1} (a_{i+1} - a_i)(a_{j+1} - a_j). \end{aligned}$$

In particular, as N goes to infinity,

$$d_{\underline{a}} = a_k N + O(1), \tag{10.12}$$

where the implicit constant depends on \underline{a} . Moreover,

$$\frac{F_{\underline{a}}(N + 1)}{F_{\underline{a}}(N)} = \frac{q^{N+1} - 1}{q^{N+1-a_k} - 1} = q^{a_k} + q^{-N+O(1)}, \tag{10.13}$$

so

$$\lim_{N \rightarrow \infty} \frac{F_{\underline{a}}(N + 1)}{F_{\underline{a}}(N)} = q^{a_k}. \tag{10.14}$$

Lemma 10.3 *Let k and m be positive integers, and let \underline{a} be an increasing sequence of k positive integers. If N is sufficiently large in terms of m and \underline{a} , then for all $g \in \text{GL}_N(q)$ with $\text{supp}(g) = m$, the number of g -stable \underline{a} -flags in \mathbb{F}_q^N can be written*

$$q^{-a_k m} (1 + \epsilon) F_{\underline{a}}(N),$$

with $|\epsilon| < q^{-N/2}$.

Proof We may assume $N > 3m$, so the eigenvalue λ of multiplicity $N - m$ is unique and therefore lies in \mathbb{F}_q . Let $W_\lambda \subset \mathbb{F}_q^N$ denote the generalized λ -eigenspace of g and W^λ the direct sum of the generalized eigenspaces of g for all eigenvalues other than λ . Thus $\dim W^\lambda < N/3$.

If $V_1 \subset \dots \subset V_k$ is a g -stable \underline{a} -flag, V_k is determined by the decomposition

$$V_k = (V_k \cap W_\lambda) \oplus (V_k \cap W^\lambda).$$

If $\dim V_k \cap W_\lambda < a_k$, then by applying (10.11) to sequences of length 1, we see that the number of possibilities for V_k is less than

$$\begin{aligned} \sum_{i=0}^{a_k-1} F_i(\dim W_\lambda) F_{a_k-i}(\dim W^\lambda) &< 4^2 \sum_{i=0}^{a_k-1} q^{i(\dim W_\lambda-i)} q^{(a_k-i)(\dim W^\lambda-a_k+i)} \\ &= q^{(a_k-1)N-O(1)}, \end{aligned}$$

so the total number of possibilities for the whole flag is less than $q^{(a_k-1)N-O(1)}$.

If $\dim V_k \cap W_\lambda = a_k$, then $V_k \subset W_\lambda$. Let I_λ and K_λ denote the image of $\lambda - g$ on W_λ and the kernel of $\lambda - g$ respectively. Because V_k is g -stable, either $V_k \subset K_\lambda$ or $V_k \cap I_\lambda \neq \{0\}$. In the latter case, V_k is spanned by a non-zero vector in I_λ and a subspace of W_λ of dimension $a_k - 1$. As $\dim I_\lambda \leq k < N/3$, the number of possibilities for the whole flag is less than $q^{(a_k-1)N-O(1)}$.

Finally, we consider the number of possibilities when $V_k \subset K_\lambda$. As g acts on K_λ as scalar multiplication, all \underline{a} -flags with $V_k \subset K_\lambda$ are g -stable. The total number is

$$F_{\underline{a}}(\dim K_\lambda) = F_{\underline{a}}(N - m) = q^{a_k m} (1 - q^{-N+O(1)}) F_{\underline{a}}(N),$$

by (10.13). The lemma follows. □

The unipotent characters of $GL_N(q)$ are indexed by partitions $\lambda \vdash N$, and we say $\chi = \chi_\lambda$ has level $N - \lambda_1$, where the parts of λ are arranged from largest to smallest, see [18, §3]. Also recall that, for $\lambda, \mu \vdash N$, the Kostka number $K_{\lambda\mu}$ is the number of semistandard Young tableaux of shape λ and weight μ .

Lemma 10.4 *Assuming $\mu_1 \geq N/2$, $K_{\lambda\mu}$ depends only the partitions $(\lambda_2, \lambda_3, \dots)$ and (μ_2, μ_3, \dots) , obtained by removing the largest parts λ_1 and μ_1 from λ and μ , and not on the value of N .*

Proof In any semistandard Young tableaux of shape λ and weight μ , the first μ_1 entries of the first row must have filled with value 1, and the remaining boxes in the first row are all to the right of every box in the remaining rows. Therefore, such a tableau is determined by choosing from the μ_2 values 2, the μ_3 values 3, and so on, an arbitrary weakly increasing sequence for the $\lambda_1 - \mu_1$ remaining boxes in the first row, and from the values that remain, a semistandard Young tableau of shape $(\lambda_2, \lambda_3, \dots)$. The number of such choices depends only on $(\lambda_2, \lambda_3, \dots)$ and (μ_2, μ_3, \dots) , but not on N . □

Proposition 10.5 *Let m and n be fixed positive integers. If N is a positive integer sufficiently large in terms of m and n , χ is a unipotent character of $GL_N(q)$ of level n , and $g \in GL_N(q)$ has support m , then*

$$\left| \frac{q^{mn} \chi(g)}{\chi(1)} - 1 \right| < q^{-N/3}. \tag{10.15}$$

Proof As $K_{\lambda\mu}$ is the number of semistandard Young tableaux of shape λ and weight μ , we have $K_{\lambda\lambda} = 1$, and if $K_{\lambda\mu} \neq 0$, then λ dominates μ : $\lambda \geq \mu$. In particular, this implies that $\mu_1 \leq \lambda_1$.

For each $\mu \vdash N$, we define the increasing sequence \underline{a}_μ of positive integers such that the sequence $a_1 = a_1 - a_0, \dots, a_{k+1} - a_k = N - a_k$ gives the parts of μ in increasing order.

Let ϕ_μ denote the permutation character of $GL_N(q)$ acting on the set of \underline{a}_μ -flags in \mathbb{F}_q^N . Then by [1, Lemma 2.4],

$$\phi_\mu = \sum_{\lambda \geq \mu} K_{\lambda\mu} \chi_\lambda.$$

If $N \geq 2n$ and $\mu_1 \geq N - n$, then by Lemma 10.4, $K_{\lambda\mu}$ depends only on $(\lambda_2, \lambda_3, \dots)$ and (μ_2, μ_3, \dots) .

As every Kostka matrix K (for partitions of N) is unitriangular, we can invert and write $\chi = \chi_\lambda$ as a linear combination of permutation characters associated to ϕ_μ , where $\mu \geq \lambda$. We can therefore express each unipotent character of level n , including χ_λ , as a linear combination of permutation characters $\chi_{\underline{b}, N}$ associated to flags \underline{b} with maximal dimension $\leq n$, with coefficients which are entries in the inverse Kostka matrix K^{-1} .

Note that, for any fixed n , the set of partitions $\lambda \vdash N$ with $\lambda_1 \geq N - n$ depends only on n , but not on N, m , or q . The unitriangularity of K implies that the submatrix of K^{-1} , truncated to only partitions of N with the first part $\geq N - n$, is the inverse of the submatrix of K , truncated to the same set of partitions. Applying Lemma 10.4, we see that all entries of this truncated submatrix of K^{-1} are bounded by some constant $O(1)$ that depends only on n :

$$|(K^{-1})_{\lambda\mu}| \leq O(1) \tag{10.16}$$

whenever $\lambda, \mu \geq N - n$. Moreover,

$$\sum_{\mu \geq \lambda} (K^{-1})_{\lambda\mu} \phi_\mu(1) = \chi_\lambda(1). \tag{10.17}$$

Define ϵ_μ so that

$$\phi_\mu(g) = q^{(\mu_1 - N)m} (1 + \epsilon_\mu) \phi_\mu(1). \tag{10.18}$$

By Lemma 10.3, for fixed m and μ , $|\epsilon_\mu| < q^{-N/2}$ if N is sufficiently large and g is of support m . When $\mu_1 = \lambda_1$, we have $q^{(\mu_1 - N)m} = q^{-mn}$. On the other hand, if $\mu_1 > \lambda_1$, then by [18, Theorem 1.2(i)] we have

$$\phi_\mu(1) \leq q^{(n-1)N}.$$

Now, by (10.16), (10.17), and (10.18),

$$\begin{aligned} \chi_\lambda(g) &= \sum_{\mu \geq \lambda} (K^{-1})_{\lambda\mu} \phi_\mu(g) \\ &= \sum_{\mu \geq \lambda} (K^{-1})_{\lambda\mu} (1 + \epsilon_\mu) q^{-mn} \phi_\mu(1) \\ &\quad + \sum_{\substack{\mu \geq \lambda \\ \mu_1 > \lambda_1}} (K^{-1})_{\lambda\mu} (1 + \epsilon_\mu) (q^{-m(N-\mu_1)} - q^{-mn}) \phi_\mu(1) \\ &= q^{-mn} \sum_{\mu \geq \lambda} (K^{-1})_{\lambda\mu} (1 + \epsilon_\mu) \phi_\mu(1) + q^{(n-1)N+O(1)} \\ &= q^{-mn} \sum_{\mu \geq \lambda} (K^{-1})_{\lambda\mu} \phi_\mu(1) + q^{-mn} \sum_{\mu \geq \lambda} \epsilon_\mu |(K^{-1})_{\lambda\mu}| \phi_\mu(1) + q^{(n-1)N+O(1)}, \\ &= (1 + \epsilon) q^{-mn} \chi_\lambda(1) + q^{(n-1)N+O(1)}, \end{aligned} \tag{10.19}$$

where $|\epsilon| \leq \max_{\mu} |\epsilon_{\mu}| < q^{-N/2}$. On the other hand, $\chi_{\lambda}(1) > q^{nN-O(1)}$ by [18, Theorem 1.2(i)]. So for N sufficiently large, (10.19) implies (10.15). \square

Proposition 10.6 *Let $p \geq 3$ be a prime, and let \mathcal{X} denote the set of cuspidal characters of $GL_p(q)$, i.e. characters of the form $I_p^k[1]$ in the notation of [16]. Let $T < GL_p(q)$ denote the centralizer of a semisimple element with characteristic polynomial irreducible over \mathbb{F}_q and $T^1 := T \cap SL_p(q)$. Let z be a central element of $SL_p(q)$. If t is a generator of T^1 , then*

$$\sum_{\chi \in \mathcal{X}} \chi(t)^2 \chi(z) = p(1 - q)\chi(1).$$

Proof First we claim that for all non-negative integers a, b, c , if

$$c \equiv q^a + q^b \pmod{|T^1|}$$

then

$$\gcd(c(q - 1), |T|) = q - 1. \tag{10.20}$$

It is clear that $q - 1$ divides both factors. If a prime ℓ divides

$$\gcd(c(q - 1), |T|) = \gcd((q^a + q^b)(q - 1), q^p - 1),$$

then it divides $|T|$, so it cannot divide q . It must also divide either $q^a + q^b$ or $q - 1$ or both. If it divides $q - 1$ but not $q^a + q^b$, then the highest power of ℓ dividing $(q^a + q^b)(q - 1)$ is the same as the highest power dividing $q - 1$ and therefore the same as the highest power dividing $\gcd(c(q - 1), |T|)$. If it divides both, then $0 \equiv q^a + q^b \equiv 2 \pmod{\ell}$, so $\ell = 2$. However, $|T^1|$ is odd (as p is odd), so the highest power of 2 dividing $|T|$ is the highest power dividing $q - 1$ and therefore the highest dividing $\gcd(c(q - 1), |T|)$. If ℓ divides only $q^a + q^b$, then we may replace a and b by their remainders under division by p (since $\ell | (q^p - 1)$), and assume ℓ divides $q^a + q^b$ for $0 \leq a \leq b < p$ and therefore divides $1 + q^{b-a}$ with $0 \leq b - a < p$. We have already seen that the highest power of 2 dividing $q - 1$ is the same as the highest power dividing $\gcd(c(q - 1), |T|)$, so we may assume $b - a > 0$. Thus, the order of $q \pmod{\ell}$ divides $2(b - a)$ as well as p , so $q \equiv 1 \pmod{\ell}$, contrary to assumption.

Let t_0 be a generator of the cyclic group T with $t_0^{q-1} = t$, and let c be as above. If ϕ is a character of T and $\phi(t)^c = 1$, then (10.20) implies $\phi^{q-1}(t_0) = 1$ and so $\phi^{q-1} = 1_T$. Therefore, if $\phi^{q-1} \neq 1_T$, we have

$$\sum_{i=0}^{|T|-1} \phi(t^{ic}) = \sum_{i=0}^{|T|-1} \phi(t_0^{ic(q-1)}) = \sum_{i=0}^{|T|-1} (\phi(t^c))^i = 0.$$

If I denotes the set of $i \in \{0, \dots, |T| - 1\}$ such that i is not divisible by $|T^1|$, then

$$\sum_{i \in I} \phi^i(t^c) = \sum_{i \in I} \phi(t^{ic}) = \sum_{i=0}^{|T|-1} \phi(t^{ic}) - \sum_{j=0}^{q-2} 1 = 1 - q. \tag{10.21}$$

Now consider the action of $\mathbb{Z}/p\mathbb{Z}$ on the character group of T which is generated by the map $\psi \mapsto \psi^q$. All orbits are of length p except for the singletons $\{\psi\}$ for which $\psi^{q-1} = 1_T$. By [16, p. 431], the restriction of any character $\chi \in \mathcal{X}$ to T is of the form

$$\chi(t) = \phi(t) + \phi(t^q) + \dots + \phi(t^{q^{p-1}})$$

for some length p orbit $\{\phi, \phi^q, \dots, \phi^{q^{p-1}}\}$, and moreover different $\chi \in \mathcal{X}$ correspond to different orbits of length p . If $g = zu$ with u unipotent, then $\chi(g) = \phi(z)\chi(u)$. Note that this is well defined because $z \neq 1$ implies $q \equiv 1 \pmod{p}$ and $z^p = 1$, which implies

$$\phi^q(z) = \phi(z^{q-1})\phi(z) = \phi(z).$$

In particular, we have

$$\chi(z) = \phi(z)\chi(1) = \phi(t_0^k)\chi(1),$$

where k is some integer divisible by $|T^1|$.

Therefore $\chi(t)^2\chi(z)/\chi(1)$ is a sum of p^2 terms of the form $\phi(t^{q^a+q^b+k})$. Denoting by ϕ_0 a generator of the character group of T , we see that the set of $\phi \in \text{Irr}(T)$ with $\phi^{q-1} \neq 1_T$ is precisely $\{\phi_0^i \mid i \in I\}$. Now using (10.21) we have

$$\begin{aligned} \sum_{\chi \in \mathcal{X}} \frac{\chi(t)^2\chi(z)}{\chi(1)} &= \frac{1}{p} \sum_{\{\phi \mid \phi^{q-1} \neq 1_T\}} \sum_{a=0}^{p-1} \sum_{b=0}^{p-1} \phi(t^{q^a+q^b+k}) \\ &= \frac{1}{p} \sum_{a=0}^{p-1} \sum_{b=0}^{p-1} \sum_{\{\phi \mid \phi^{q-1} \neq 1_T\}} \phi^{q^a+q^b+k}(t) \\ &= \frac{1}{p} \sum_{a=0}^{p-1} \sum_{b=0}^{p-1} \sum_{i \in I} \phi_0^i(t^{q^a+q^b+k}) \\ &= \frac{1}{p} \sum_{a=0}^{p-1} \sum_{b=0}^{p-1} (1 - q) = p(1 - q). \end{aligned}$$

□

Theorem 10.7 *For all but finitely many ordered pairs (p, q) where $p \geq 3$ is prime and q is a prime power, the following statement holds. If t is a generator of the norm-1 subgroup $T_1 \cong C_{(q^p-1)/(q-1)}$ of $\mathbb{F}_{q^p}^\times$ then every non-central element of $\text{SL}_p(q)$ is a product of two conjugates of t .*

Proof Fixing an \mathbb{F}_q -basis of \mathbb{F}_{q^p} we can identify $\mathbb{F}_{q^p}^\times$ with the centralizer T in $G := \text{GL}_p(q)$ of any generator of \mathbb{F}_{q^p} . As p is prime, every non-central element of T generates \mathbb{F}_{q^p} as \mathbb{F}_q -algebra, so no such element is contained in a proper parabolic subgroup of G . Therefore, every Harish-Chandra induced character of G vanishes on every element of $T \setminus \mathbf{Z}(G)$; in particular at our element t . By [16, (12)], a primary (i.e. not Harish-Chandra induced) character of G can be non-zero at t if and only if it is of the form $I_1^k[p]$ or of the form $I_p^k[1]$. In the first case, it belongs to the set \mathcal{X} of Proposition 10.6. In the second case, it is the product of a unipotent character and a linear character. Since $\mathbf{C}_G(t) = T$, the conjugate classes of t in G and in $\text{SL}_p(q)$ are the same.

By Theorem 10.1, we may assume that our target element g has bounded support. By the Frobenius formula, $(q - 1)^{-1}$ times the number of representations of g as a product of two conjugates of t in G is

$$(q - 1)^{-1} \frac{|t^G|^2 |g^G|}{|G|} \sum_{\chi \in \text{Irr}(G)} \frac{\chi(t)^2 \bar{\chi}(g)}{\chi(1)}.$$

We divide this sum into a sum over χ which are unipotent characters times linear characters and a sum over $\chi \in \mathcal{X}$.

For the first, we note that t and g are both in $SL_p(\mathbb{F}_q)$, and all linear characters of G are trivial on this subgroup. So we can simply sum over unipotent characters and omit the factor $(q - 1)^{-1}$. The contribution of the trivial character to the sum

$$\sum_{\chi \in \text{Irr}(G)} \frac{\chi(t)^2 \bar{\chi}(g)}{\chi(1)}$$

is 1. For the other unipotent characters, by [16, Theorem 12], $\chi_\lambda(t)$ is given by the value at a p -cycle of the character of the symmetric group S_p associated to the partition $\lambda \vdash p$, and by the Murnaghan-Nakayama rule, this value is ± 1 if λ is of the form $1^n(p - n)^1$ and 0 otherwise. By the main theorem of [21], since $g \notin \mathbf{Z}(G)$, there exists an absolute constant $\epsilon > 0$ such that

$$|\chi(g)| \leq \chi(1)^{1-\epsilon/p} \tag{10.22}$$

for all $\chi \in \text{Irr}(G)$. By the dimension formula for primary characters of G [16, Lemma 7.4],

$$\chi_\lambda(1) \geq q^{n(p-n/2-1/2)} \geq q^{np/3}.$$

Therefore,

$$\frac{|\chi_{1^n(p-n)^1}(g)|}{\chi_{1^n(p-n)^1}(1)} \leq 2^{-n\epsilon/3}.$$

Choosing A to be a sufficiently large absolute constant, we have $\sum_{n \geq A} 2^{-n\epsilon/3} < \frac{1}{3}$, which guarantees

$$\left| \sum_{\chi = \chi_{1^n(p-n)^1}, n \geq A} \frac{\chi(t)^2 \bar{\chi}(g)}{\chi(1)} \right| \leq \frac{1}{3}.$$

On the other hand, if p is large enough compared to A , then by Proposition 10.5, for $1 \leq n < A$ we have $\chi_{1^n(p-n)^1}(g) > 0$, and

$$\sum_{1 \leq n < A} \frac{\chi_{1^n(p-n)^1}(g)}{\chi_{1^n(p-n)^1}(1)}$$

is the sum of a positive term γ and an error term less than $1/3$ in absolute value. Recalling $\chi(t)^2 = 1$ on all these $\chi = \chi_{1^n(p-n)^1}$, it follows that the total contribution to the sum from nontrivial unipotent characters is γ plus an error term less than $2/3$ in absolute value. Hence the total contribution to the sum from all unipotent characters is at least $\gamma + 1/3$. If p is bounded but q is large enough, then by Gluck’s bound [15], there is some absolute constant $C > 0$ such that

$$\left| \sum_{1 \leq n < A} \frac{\chi_{1^n(p-n)^1}(g)}{\chi_{1^n(p-n)^1}(1)} \right| < \frac{CA}{\sqrt{q}} \leq \frac{1}{3},$$

and this ensures that the total contribution to the sum from all unipotent characters is at least $1/3$.

As p is prime, for $\chi \in \mathcal{X}$, $\chi(g) = 0$ unless $g = zu$, where z is scalar and u is unipotent, or g is conjugate to an element of T . In the former case, $\chi(g) = c_u \frac{\chi(z)}{\chi(1)}$, where $c_u \in \mathbb{Z}$ depends only on u but not on the particular $\chi \in \mathcal{X}$. By Proposition 10.6,

$$\frac{1}{q-1} \left| \sum_{\chi \in \mathcal{X}} \frac{\chi(t)^2 \bar{\chi}(g)}{\chi(1)} \right| = \frac{p|c_u|}{\chi(1)}.$$

As g is not scalar, $|c_u| = |\chi(g)| \leq \chi(1)^{1-\epsilon/p}$ by (10.22). Since $\chi(1) = \prod_{i=1}^p (q^i - 1) > q^{p(p-1)/2}/4$, we get a uniform bound $|c_u|/\chi(1) \leq q^{-\beta p}$ for some $\beta > 0$. Thus, the contribution of the cuspidal characters is less than $1/6$ when either p or q is sufficiently large, implying the theorem. \square

Data availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Andrews, S., Thiem, N.: The combinatorics of GL_n generalized Gelfand–Graev characters. *J. Lond. Math. Soc.* (2) **95**(2), 475–499 (2017)
2. Arad, Z., Herzog, M. (eds.): Products of Conjugacy Classes in Groups. *Lecture Notes in Mathematics*, vol. 1112. Springer, New York (1985)
3. Asai, T.: Unipotent class functions of split special orthogonal groups SO_{2n}^+ over finite fields. *Commun. Algebra* **12**(5–6), 517–615 (1984)
4. Asai, T.: The unipotent class functions on the symplectic groups and the odd orthogonal groups over finite fields. *Commun. Algebra* **12**(5–6), 617–645 (1984)
5. Bonnafé, C.: Sur les caractères des groupes réductifs finis à centre non connexe: applications aux groupes spéciaux linéaires et unitaires. *Astérisque* **306** (2006)
6. Carter, R.W.: *Finite Groups of Lie Type*, Wiley Classics Library. Wiley, Chichester (1993)
7. Deligne, P., Lusztig, G.: Representations of reductive groups over finite fields. *Ann. Math.* **103**(1), 103–161 (1976)
8. Digne, F., Michel, J.: *Representations of Finite Groups of Lie Type*. London Mathematical Society Student Texts, vol. 95, 2nd edn. Cambridge University Press, Cambridge (2020)
9. Ellers, E.W., Gordeev, N.: On the conjectures of J. Thompson and O. Ore. *Trans. Am. Math. Soc.* **350**(9), 3657–3671 (1998)
10. Fong, P., Srinivasan, B.: The blocks of finite general linear and unitary groups. *Invent. Math.* **69**, 109–153 (1982)
11. Fulman, J., Guralnick, R.M.: Bounds on the number and sizes of conjugacy classes in finite Chevalley groups with applications to derangements. *Trans. Am. Math. Soc.* **364**, 3023–3070 (2012)
12. Geck, M.: *An Introduction to Algebraic Geometry and Algebraic Groups*, Oxford Graduate Texts in Mathematics, vol. 10. Oxford University Press, Oxford (2003)
13. Geck, M., Malle, G.: *The Character Theory of Finite Groups of Lie Type: A Guided Tour*, Cambridge Studies in Advanced Mathematics, vol. 187. Cambridge University Press, Cambridge (2020)
14. Geck, M., Pfeiffer, G.: *Characters of Finite Coxeter Groups and Iwahori–Hecke Algebras*. London Mathematical Society Monographs. New Series, vol. 21. The Clarendon Press, Oxford University Press, New York (2000)
15. Gluck, D.: Sharper character value estimates for groups of Lie type. *J. Algebra* **174**(1), 229–266 (1995)
16. Green, J.: The characters of the finite general linear groups. *Trans. Am. Math. Soc.* **80**, 402–447 (1955)
17. Guralnick, R.M., Malle, G.: Simple groups admit Beauville structures. *J. Lond. Math. Soc.* **85**, 694–721 (2012)
18. Guralnick, R.M., Larsen, M., Tiep, P.H.: Character levels and character bounds. *Forum Math. Pi* **8**, e2 (2020)
19. Isaacs, I.M.: *Character Theory of Finite Groups*, Corrected Reprint of the 1976 Original [Academic Press, New York]. AMS Chelsea Publishing, Providence (2006)
20. Landazuri, V., Seitz, G.M.: On the minimal degrees of projective representations of the finite Chevalley groups. *J. Algebra* **32**, 418–443 (1974)

21. Larsen, M., Tiep, P.H.: Uniform characters bounds for finite classical groups, (submitted)
22. Larsen, M., Shalev, A.: Word maps and Waring type problems. *J. Am. Math. Soc.* **22**(2), 437–466 (2009)
23. Larsen, M., Malle, G., Tiep, P.H.: The largest irreducible representations of simple groups. *Proc. Lond. Math. Soc.* **106**, 65–96 (2013)
24. Lassueur, C., Malle, G.: Simple endotrivial modules for linear, unitary and exceptional groups. *Math. Z.* **280**(3–4), 1047–1074 (2015)
25. Liebeck, M.W., Shalev, A.: Character degrees and random walks in finite groups of Lie type. *Proc. Lond. Math. Soc.* (3) **90**(1), 61–86 (2005)
26. Liebeck, M.W., O'Brien, E.A., Shalev, A., Tiep, P.H.: The Ore conjecture. *J. Eur. Math. Soc. (JEMS)* **12**(4), 939–1008 (2010)
27. Lübeck, F., Malle, G.: A Murnaghan–Nakayama rule for values of unipotent characters in classical groups. *Represent. Theory Ser.* **20**, 139–161 (2016)
28. Lusztig, G.: *Characters of Reductive Groups Over a Finite Field*. *Annals of Mathematics Studies*, vol. 107. Princeton University Press, Princeton (1984)
29. Lusztig, G.: Character sheaves. IV. *Adv. Math.* **59**(1), 1–63 (1986)
30. Malle, G.: The decomposition of Lusztig induction in classical groups. *J. Algebra Ser.* **558**, 611–619 (2020)
31. Olsson, J.B.: *Combinatorics and Representations of Finite Groups*. *Vorlesungen aus dem Fachbereich Mathematik der Universität GH Essen [Lecture Notes in Mathematics at the University of Essen]*, 20. Universität Essen, Fachbereich Mathematik, Essen (1993)
32. Ore, O.: Some remarks on commutators. *Proc. Am. Math. Soc.* **2**, 307–314 (1951)
33. Stembridge, J.R.: On the eigenvalues of representations of reflection groups and wreath products. *Pac. J. Math.* **140**(2), 353–396 (1989)
34. Waldspurger, J. L.: Une conjecture de Lusztig pour les groupes classiques. *Mém. Soc. Math. Fr. (N.S.)* **96** (2004)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.