

Staffan Uhlén · Jonas Lindblom · Anna Kindlundh  
Paul Mugisha · Fred Nyberg

## Nandrolone treatment decreases the level of rat kidney $\alpha_{1B}$ -adrenoceptors

Published online: 29 October 2003

© Springer-Verlag 2003

### Naunyn-Schmiedeberg's Arch Pharmacol (2003) 368:91–98

In Table 1, two values were inadvertently transposed. The correct table is given here:

**Table 1**

	$\alpha_{1A}$ pK <sub>d</sub> [ <sup>3</sup> H]-prazosin	$\alpha_{1B}$ pK <sub>d</sub> [ <sup>3</sup> H]-prazosin	$\alpha_{1A}$ (fmol/mg)	$\alpha_{1B}$ (fmol/mg)	$\alpha_{1B}$ (% in individual experiments)	$\alpha_{1A}$ pK <sub>i</sub> 5-MU	$\alpha_{1B}$ pK <sub>i</sub> 5-MU	(n)
t-test	<i>p</i> =0.46	<i>p</i> =0.35	<i>p</i> =0.19	<i>p</i> <0.0001	<i>p</i> <0.0001	<i>p</i> =0.21	<i>p</i> =0.24	
control	9.40±0.07	9.63±0.09	44.4±3.2	14.8±1.8	25.0±2.8	8.63±0.08	6.66±0.17	12
nandrolone treated	9.41±0.07	9.58±0.12	48.6±3.6	6.5±0.8	12.4±1.6	8.53±0.08	6.41±0.30	13

The online version of the original article can be found at  
<http://dx.doi.org/10.1007/s00210-003-0763-y>

S. Uhlén · A. Kindlundh · P. Mugisha · F. Nyberg  
Department of Pharmaceutical Biosciences, Uppsala University,  
Box 591 BioMedical Centre, 751 24 Uppsala, Sweden

J. Lindblom (✉)  
Department of Neuroscience, Division of Pharmacology,  
Uppsala University, Box 593 BioMedical Centre,  
751 24 Uppsala, Sweden,  
Fax: +46-18-511540,  
e-mail: Jonas.Lindblom@neuro.uu.se