



# Sesamol protects against liver fibrosis induced in rats by modulating lysophosphatidic acid receptor expression and TGF- $\beta$ /Smad3 signaling pathway

Nesma A. Abd Elrazik<sup>1</sup> · Mohamed El-Mesery<sup>1</sup> · Mamdouh M. El-Shishtawy<sup>1</sup>

Received: 26 January 2022 / Accepted: 20 May 2022 / Published online: 1 June 2022  
© The Author(s) 2022

## Abstract

The present study aimed to investigate the hepatoprotective effect of sesamol (SML), a nutritional phenolic compound obtained from sesame seeds, in liver fibrosis induced by thioacetamide (TAA) in rats and to explore the underlying mechanisms. Thirty-two male Sprague–Dawley rats were equally divided into four groups: control, TAA, TAA + SML 50 mg/kg, and TAA + SML 100 mg/kg groups. Liver functions and hepatic contents of glutathione (GSH) and malondialdehyde (MDA) were measured colorimetrically. Gene expressions of lysophosphatidic acid receptor (LPA)-1 and -3, connective tissue growth factor (CTGF), transforming growth factor (TGF)- $\beta$ 1, small mothers against decapentaplegic (Smad)-3 and -7,  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA), and cytokeratin 19 (CK19) were analyzed by qRT-PCR. Moreover, phosphorylated Smad3 (pSmad3) was quantified by ELISA. Additionally, TGF- $\beta$ 1,  $\alpha$ -SMA, CK19, and vascular endothelial growth factor (VEGF) protein concentrations were semi-quantitatively analyzed by immunostaining of liver sections. SML treatment markedly improved liver index and liver functions. Moreover, SML protected against liver fibrosis in a dose-dependent manner as indicated by down-regulation of LPA1, LPA3, CTGF, TGF- $\beta$ 1/Smad3, and  $\alpha$ -SMA expressions and a decrease in pSmad3 level, as well as an up-regulation of Smad7 expression. In addition, SML suppressed ductular reaction hinted by the decrease in CK19 expression. These results reveal the anti-fibrotic effect of SML against liver fibrosis that might be attributed to down-regulation of LPA1/3 expressions, inhibition of TGF- $\beta$ 1/Smad3 pathway, and ductular reaction.

**Keywords** Ductular reaction · Liver fibrosis · LPA · Sesamol · TGF- $\beta$ 1/Smad3

## Abbreviations

$\alpha$ -SMA	$\alpha$ -Smooth muscle actin
CTGF	Connective tissue growth factor
CK19	Cytokeratin 19
CMC	Carboxy methyl cellulose
GAPDH	Glyceraldehyde 3-phosphate dehydrogenase
ECM	Extracellular matrix
GSH	Glutathione
H&E	Hematoxylin and eosin
HPCs	Hepatic progenitor cells
HSCs	Hepatic stellate cells
LPA	Lysophosphatidic acid receptor
MDA	Malondialdehyde
ROS	Reactive oxygen species

Smad	Small mothers against decapentaplegic
SML	Sesamol
TAA	Thioacetamide
TGF- $\beta$ 1	Transforming growth factor- $\beta$ 1
VEGF	Vascular endothelial growth factor

## Introduction

Liver fibrosis is a complex inflammatory process that may progress to cirrhosis and hepatocellular carcinoma (HCC). Hepatic fibrosis demonstrates the wound healing process in response to liver injury which is characterized by an elevation in extracellular matrix (ECM) deposition around the sinusoidal cell layer in the space of Disse. The imbalance between ECM synthesis and degradation leads to an increase in the fibrotic matrix (Zhang et al. 2016; Roehlen et al. 2020).

Thioacetamide (TAA), a thiosulfur compound, is a hepatotoxic agent that is used in research to induce liver

✉ Mamdouh M. El-Shishtawy  
mshisht@mans.edu.eg

<sup>1</sup> Department of Biochemistry, Faculty of Pharmacy,  
Mansoura University, P.O. Box, Mansoura 35516, Egypt

fibrosis and HCC by recurrent administration in experimental animals (Helmy et al. 2018; Ebrahim et al. 2018; Helmy et al. 2019; Shaker et al. 2021). TAA is metabolized to TAA sulfoxide and TAA disulfoxide metabolites that covalently bind to cellular macromolecules leading to an increase in reactive oxygen species (ROS) generation and acute centrilobular liver necrosis (Eraky et al. 2018; Nevzorova et al. 2020).

Lysophosphatidic acid (LPA) is a growth factor-like phospholipid that regulates various processes in the cell including cell motility, cell survival, cell proliferation, and cellular differentiation. LPA is produced from phospholipids by various enzymatic pathways. It works on specific G-protein coupled receptors called LPA receptors (LPARs). The signaling of LPARs leads to many actions on physiological, developmental, and pathological processes like liver fibrosis and liver cancer. It has been elucidated that fibrosis is firmly affected by elevated concentration of LPA and some of its receptors as LPAR1 and LPAR3 (Simo et al. 2014; Kaffe et al. 2017; Eraky et al. 2018; Xiang et al. 2020). Moreover, LPA signaling is involved in the fibrotic process through activating transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) which is considered as the cornerstone of fibrosis development (Li et al. 2017; Eraky et al. 2018).

The hepatic stellate cell (HSC) activation presents a substantial role in the initiation and development of liver fibrosis. Fibrogenic cytokines like TGF- $\beta$ 1 activate quiescent HSCs and promote trans-differentiation into myofibroblasts during liver injury. This step is characterized by the expression of  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA), eventually leading to ECM accumulation (Higashi et al. 2017). TGF- $\beta$ 1 stimulates phosphorylation of small mothers against decapentaplegic (Smad)-2 and -3 which in turn triggers overexpression of pro-fibrotic genes such as collagen,  $\alpha$ -SMA, and connective tissue growth factor (CTGF). TGF- $\beta$ 1/Smad pathway dysregulation is an important mechanism of pathogenesis in fibrosis of liver (Hu et al. 2018; Mu et al. 2018).

Ductular reaction is the proliferation of reactive bile ducts stimulated via liver injury. Cytokeratin 19 (CK19) is expressed normally in the lining of hepatobiliary tracts and considered as a helpful marker for the occurrence of ductular reaction which is closely related to liver fibrosis and damage (Jain et al. 2010; Sato et al. 2019).

Sesamol (SML), an active phenolic lignan and a prominent fragrance component in sesame oil, exhibits antioxidant, free radical scavenging ability (Gao et al. 2017) and anti-inflammatory effect (Abd Elrazik et al. 2021). Moreover, it has been demonstrated to display wide array of biological activities such as cardioprotective (Jayaraj et al. 2020), anti-mutagenic (Kaur and Saini 2000), radioprotective (Majdaeen et al. 2020), neuroprotective (Sachdeva et al. 2015), and anti-platelet activity (Chang et al. 2010). SML was reported to ameliorate cyclophosphamide-induced hepatotoxicity (Jnaneshwari et al. 2014), but the molecular

mechanisms of anti-fibrotic effect of SML are not well known.

Therefore, the current study aimed to investigate the probable protective effects of SML against liver fibrosis induced in rats by TAA. Additionally, we also aimed to evaluate SML possible mechanisms of action.

## Materials and methods

### Chemicals

TAA and SML were purchased from Sigma-Aldrich (St. Louis, MO, USA). Other chemicals and reagents consumed during this study were of high analytical grade and purchased from El-Gomhouria Co. For Trading Drugs, Chemicals & Medical Supplies (Cairo, Egypt).

### Animals

Adult male Sprague–Dawley rats weighing 180–200 g (8–10 weeks old) were obtained from VACSERA CO, Egypt. The animals were allowed to standard rat pellet food and water ad libitum and housed under controlled condition of  $25 \pm 2$  °C room temperature, 45–55% humidity, and 12-h dark/light cycle. This study protocol was approved by “Research Ethics Committee” Faculty of Pharmacy, Mansoura University, Mansoura City, Egypt (approval code: 2021–401), which is in accordance with “Guide for the Care and Use of Laboratory Animals, 8th edition” (National Research Council (US); Committee for the Update of the Guide for the Care and Use of Laboratory Animals, 2011).

### Experimental design

Rats were equally divided into four groups (8 animals/group):

**Control group:** rats received no treatment, only 0.2-mL sterile saline by intraperitoneal (i.p.) injection twice weekly for 8 weeks. **TAA group:** rats were injected by TAA (200 mg/kg dissolved in saline, i.p., twice weekly for 8 weeks) (Eraky et al. 2018). **SML 50 group:** rats received TAA (200 mg/kg/i.p.) twice weekly concurrently with SML (50 mg/kg/day) by oral gavage suspended in carboxymethyl cellulose (CMC) for 8 weeks (Hemalatha et al. 2013). **SML 100 group:** rats received TAA (200 mg/kg/i.p.) twice weekly concurrently with SML (100 mg/kg/day) by oral gavage suspended in CMC for 8 weeks (Hemalatha et al. 2013).

Based on previous studies and our preliminary experiments, it has been demonstrated that CMC had no statistically significant effects or treatment-related adverse effects and considered safe for all animal species (Helmy et al. 2019; Zeyada et al. 2020).

After 8 weeks, rats were weighted, anesthetized by thiopental (40 mg/kg, i.p.), and blood was gathered through the retro-orbital puncture and then centrifuged at 4000 rpm for serum separation and assessment of biochemical parameters. Then, all animals were sacrificed and their livers were isolated and weighted for calculation of the liver index (liver weight to body weight %) (Yogalakshmi et al. 2010). Isolated livers were divided into three portions. The first one was immediately transferred in RNA later, then stored at 4 °C overnight, and kept at –80 °C for qRT-PCR. The second portion was homogenized in (10% w/v) phosphate buffered saline (pH 7.4) for oxidative stress marker assessment and ELISA experiment. The third portion was fixed in 10% neutral buffered formalin for histopathological and immunohistochemical analyses.

### Serum biochemical analysis

Serum was used to measure aspartate aminotransferase (AST) and alanine aminotransferase (ALT) activities (Spectrum Diagnostics, Egypt), in addition to measuring levels of albumin, total protein (BioMed Diagnostic, Egypt), and bilirubin (Diamond Diagnostics, Egypt) following the manufacturer's instructions.

### Determination of oxidative stress biomarkers

Hepatic tissue supernatants were used to assess the tissue content of malondialdehyde (MDA) and glutathione (GSH) following the manufacturer's instructions (Biodiagnostic, Giza, Egypt).

### Histopathological analysis of hepatic tissue

Paraffin-embedded blocks of hepatic tissue were dissected into 5- $\mu$ m thickness segments, then stained with hematoxylin and eosin (H&E) stain for necroinflammatory scores assessment according to Ishak's activity index (Ishak et al. 1995).

For analysis of collagen content in the fibrotic areas, sections were stained with Masson's trichrome stain and the fibrosis intensity was demonstrated as a percentage of the fibrotic (stained) area to the total area via analyzing 30 random fields in the same slide (James et al. 1990).

### Immunohistochemistry

Immunohistochemical analysis was carried out on 5- $\mu$ m-thick deparaffinized liver sections incubated overnight at 4 °C with polyclonal antibodies for TGF- $\beta$ 1 (Fine Test, China), CK19 (Zeta Corporation, USA),  $\alpha$ -SMA, and vascular endothelial growth factor (VEGF) (Servicebio, China). Sections were treated with secondary antibodies conjugated with horseradish peroxidase and visualized by 2%

diaminobenzidine reagent. Then, slides were counterstained with hematoxylin and examined by a light microscope.

### Determination of hepatic phosphorylated Smad3 (pSmad3) protein level by ELISA

Hepatic pSmad3 protein level was measured in the liver homogenates by commercially available ELISA kit (Bioassay, England) following manufacturer's instruction.

### Gene expression analysis using quantitative real-time polymerase chain reaction (qRT-PCR)

RNA was isolated from liver tissues using Qiagen RNeasy Plus Mini kit (Qiagen®, USA). The concentration and purity of the extracted RNA were measured spectrophotometrically by NanoDrop System (Thermo Fisher Scientific Inc.). After that, RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher Scientific, USA) was used to reverse transcript 2  $\mu$ g of RNA to cDNA. Real-time PCR was carried out using HERA SYBR® Green qRT-PCR Kit (Willowfort, England) on Piko Real-PCR System (Thermo Fisher Scientific Inc.). The primer sequences used in this work are listed in Table 1. The mRNA expressions of these genes were normalized against

**Table 1** Sequences of primers used in qRT-PCR experiments

Name	Primer sequence
GAPDH — forward	5'-TCCCATTCTTCCACCTTTGA-3'
GAPDH — reverse	5'-CCACCACCTGTGTGCTGTAG-3'
LPAR1 — forward	5'-TTTCACAGCCCCAGTTCACA-3'
LPAR1 — reverse	5'-GCTTGCTCACAGTGTTCAT-3'
LPAR3 — forward	5'-GCAAGGGTGGAGGTGTAGAA-3'
LPAR3 — reverse	5'-GGTCTAAACTCGCCATCACG-3'
CTGF — forward	5'-CGAGTCCTTCCAAAGCAGTT-3'
CTGF — reverse	5'-ATCACACACCCACTCCTCAC-3'
TGF- $\beta$ 1 — forward	5'-CCGCAACAACGCAATCATGA-3'
TGF- $\beta$ 1 — reverse	5'-GCACTGCTTCCCGAATGTCT-3'
Smad3 — forward	5'-AGACACCAGTGCTACCTCCA-3'
Smad3 — reverse	5'-CCAGCGGGGAAGTTAGTGTT-3'
Smad7 — forward	5'-TTTTTCCCCCACCCTTCCAAC-3'
Smad7 — reverse	5'-AAACACACCACCTTCTCGCACC-3'
$\alpha$ -SMA — forward	5'-ACCATCGGGAATGAACGCTT-3'
$\alpha$ -SMA — reverse	5'-CTGTCAGCAATGCCTGGGTA-3'
CK19 — forward	5'-CGCATCGTGTCTCATCCTC-3'
CK19 — reverse	5'-GCCCACTAAAACCTTCCACC-3'

GAPDH glyceraldehyde 3-phosphate dehydrogenase, LPAR1 lysophosphatidic acid receptor1, LPAR3 lysophosphatidic acid receptor3, CTGF connective tissue growth factor, TGF- $\beta$ 1 transforming growth factor  $\beta$ 1, Smad3 small mothers against decapentaplegic3, Smad7 small mothers against decapentaplegic7,  $\alpha$ -SMA alpha smooth muscle actin, CK19 cytokeratin 19

glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as a housekeeping gene and quantified via the  $2^{-\Delta\Delta C_t}$  method.

## Statistical analysis

The experimental data were expressed as mean  $\pm$  SEM. Data analysis and graphing were done by GraphPad Prism 6.01 (GraphPad Software, San Diego, CA, USA). One-way analysis of variance (ANOVA) followed by Tukey's post hoc test was used for statistical evaluations. For the analysis of histopathology scores, non-parametric Kruskal–Wallis test followed by Dunn's multiple comparison post hoc test were used.

## Results

### SML reduces liver index

Liver index significantly increased in TAA-intoxicated rats by 1.9-fold ( $P < 0.0001$ ) compared with control group, while oral administration of either 50 or 100 mg/kg SML significantly reduced liver index by 23.7% ( $P < 0.0001$ )

and 38.5% ( $P < 0.0001$ ), respectively, compared with TAA group (Table 2).

### SML attenuates TAA-induced liver dysfunction and histopathological alterations in rats

TAA group showed a marked elevation in serum ALT and AST activities and total bilirubin level compared with control group ( $P < 0.0001$ ). In contrary, albumin and total protein levels were significantly reduced in TAA group compared with control group ( $P < 0.0001$ ). However, treating rats with either 50 or 100 mg/kg SML in the presence of TAA caused a significant improvement of liver function compared with TAA group (Table 2).

Histopathological examination of liver tissues stained with H&E showed a normal architecture of liver tissues in control group, while TAA group showed a significant inflammation and necrosis compared with control group ( $P < 0.0001$ ). On the contrary, both doses of SML treatment caused a reduction in necroinflammatory scores (Fig. 1).


**Table 2** Effect of 50 and 100 mg/kg sesamol (SML) on liver index and liver function tests

Groups	Control	TAA	SML 50	SML 100
Liver index	2.1 $\pm$ 0.03	3.88 $\pm$ 0.04 <sup>++++</sup>	3.02 $\pm$ 0.06 <sup>++++, ****</sup>	2.43 $\pm$ 0.03 <sup>++++, ****, SSSS</sup>
ALT (IU/L)	42.25 $\pm$ 4.18	192.62 $\pm$ 2.41 <sup>++++</sup>	123.5 $\pm$ 3.7 <sup>++++, ****</sup>	91.5 $\pm$ 1.45 <sup>++++, ****, SSSS</sup>
AST (IU/L)	122.87 $\pm$ 3.36	416.5 $\pm$ 8.24 <sup>++++</sup>	267.62 $\pm$ 3.34 <sup>++++, ****</sup>	219.12 $\pm$ 3.45 <sup>++++, ****, SSSS</sup>
Total bilirubin (mg/dL)	0.14 $\pm$ 0.01	0.66 $\pm$ 0.01 <sup>++++</sup>	0.37 $\pm$ 0.02 <sup>++++, ****</sup>	0.27 $\pm$ 0.01 <sup>++++, ****, SSSS</sup>
Albumin (g/dL)	3.98 $\pm$ 0.03	2.75 $\pm$ 0.06 <sup>++++</sup>	3.08 $\pm$ 0.03 <sup>++++, ****</sup>	3.48 $\pm$ 0.04 <sup>++++, ****, SSSS</sup>
Total protein (g/dL)	8.56 $\pm$ 0.11	6.21 $\pm$ 0.07 <sup>++++</sup>	6.66 $\pm$ 0.05 <sup>++++, **</sup>	7.05 $\pm$ 0.04 <sup>++++, ****, SS</sup>

ALT alanine aminotransferase, AST aspartate aminotransferase, TAA thioacetamide

Values are expressed as mean  $\pm$  SEM ( $n = 8$ ). <sup>+</sup>Significance against control group (<sup>+++</sup> $P < 0.001$ , <sup>++++</sup> $P < 0.0001$ ), <sup>\*</sup>significance against TAA group (<sup>\*\*</sup> $P < 0.01$ , <sup>\*\*\*\*</sup> $P < 0.0001$ ), and <sup>§</sup>significance against SML 50 group (<sup>§§</sup> $P < 0.01$ , <sup>§§§§</sup> $P < 0.0001$ )

**Fig. 1** Sesamol (SML) attenuates TAA-induced histopathological alterations in rats. **a** Histopathological changes of liver sections stained with H&E. Blue arrows indicate necroinflammatory lesions and black arrows indicate fibrosis ( $\times$ : 100, bar 100). **b** The necroinflammatory score. Values are expressed as median ( $n = 8$ ). TAA thioacetamide. <sup>+</sup>Significance against control group (<sup>++</sup> $P < 0.01$ , <sup>++++</sup> $P < 0.0001$ ). <sup>\*</sup>Significance against TAA group (<sup>\*</sup> $P < 0.05$ )



### SML attenuates TAA-induced liver fibrosis

In cross-sectional images of liver tissues stained with Masson's trichrome, control group showed no collagen deposition, while TAA caused a remarkable increase in collagen deposition and fibrosis percentage in liver tissue ( $P < 0.0001$ ). Rats treated with SML 50 or 100 mg/kg revealed a marked reduction in collagen deposition and fibrosis percentage compared with TAA-intoxicated rats ( $P < 0.0001$ ) (Fig. 2).

### SML improves the anti-oxidant capacity in rats

As shown in Fig. 3, TAA group showed 3.7-fold ( $P < 0.0001$ ) significant increase in hepatic MDA content associated with 4.7-fold ( $P < 0.0001$ ) marked decrease in hepatic GSH content compared with control group. Interestingly, 50- or 100-mg/kg SML administration exhibited a dose-dependent decrease in MDA ( $P < 0.0001$ ) as well as a dose-dependent elevation in GSH content ( $P < 0.01$ ,  $P < 0.0001$ , respectively) compared with TAA group.

### SML down-regulates gene expressions of LPAR1, LPAR3, and CTGF


As shown in Fig. 4, TAA group showed a marked up-regulation in gene expressions of LPAR1, LPAR3, and CTGF compared with control group ( $P < 0.0001$ ). However, treatment with SML 50 or 100 mg/kg caused a dose-dependent significant down-regulation in the mRNA expressions of LPAR1, LPAR3, and CTGF compared with TAA group ( $P < 0.0001$ ).

### SML reduces TGF- $\beta$ 1 gene expression and its protein concentration

As shown in Fig. 5, remarkable increase in TGF- $\beta$ 1 mRNA expression and its protein concentration was detected in TAA group, compared with control group ( $P < 0.0001$ ). After 8 weeks of daily administration of either 50 or 100 mg/kg SML, gene expression and protein concentration of TGF- $\beta$ 1 ( $P < 0.0001$ ) were reduced significantly compared with TAA group in a dose-dependent manner.


**Fig. 2** Sesamol (SML) attenuates TAA-induced liver fibrosis.


**a** Liver sections stained with Masson's trichrome. Black arrows indicate collagen deposition ( $\times$ : 100, bar 100). **b** % fibrosis. TAA thioacetamide. Values are expressed as mean  $\pm$  SEM ( $n = 6$ ). <sup>+</sup>Significance against control group ( $++++P < 0.0001$ ), <sup>\*</sup>significance against TAA group ( $****P < 0.0001$ ), and <sup>§</sup>significance against SML 50 group ( $.\$P < 0.01$ )



**Fig. 3** Sesamol (SML) improves the anti-oxidant capacity in rats.

**a** Malondialdehyde (MDA). **b** Glutathione (GSH). TAA thioacetamide. Values are expressed as mean  $\pm$  SEM ( $n = 8$ ). <sup>+</sup>Significance against control group ( $++++P < 0.0001$ ), <sup>\*</sup>significance against TAA group ( $**P < 0.01$ ,  $****P < 0.0001$ ), and <sup>§</sup>significance against SML 50 group ( $.\$P < 0.0001$ )





**Fig. 4** Sesamol (SML) down-regulates gene expressions of lysophosphatidic acid receptor1 (LPAR1), LPAR3, and connective tissue growth factor (CTGF). The mRNA expression of (a) LPAR1, (b) LPAR3, and (c) CTGF. TAA thioacetamide. Values are expressed as

mean  $\pm$  SEM ( $n=6$ ). +Significance against control group ( $^+P<0.05$ ,  $++++P<0.0001$ ), \*significance against TAA group ( $****P<0.0001$ ), and \$significance against SML 50 group ( $^{\$}P<0.05$ ,  $$$$P<0.001$ ,  $$$$$P<0.0001$ )

### SML down-regulates Smad3 gene expression and pSmad3 protein concentration and up-regulates Smad7 gene expression

TAA group showed about 4.3-fold ( $P<0.0001$ ) significant elevation in Smad3 gene expression as well as 6.5-fold ( $P<0.0001$ ) marked increase in pSmad3 protein level compared with control group. However, SML 50- and 100-mg/kg treatments showed a dose-dependent marked decrease in the mRNA expression of Smad3 and pSmad3 protein level compared with TAA group ( $P<0.0001$ ) (Fig. 6a, b).


Gene expression of Smad7 significantly down-regulated by 70% ( $P<0.001$ ) in TAA group compared with control group. On the contrary, treatment with SML either 50 or

100 mg/kg revealed 2.6- ( $P<0.01$ ) and 6.3-fold ( $P<0.0001$ ) marked increase in Smad7 mRNA expression, respectively, compared with TAA group in a dose-dependent manner (Fig. 6c).

### SML suppresses $\alpha$ -SMA gene expression and its protein concentration

TAA significantly increased  $\alpha$ -SMA gene expression and its protein concentration compared with control group ( $P<0.0001$ ). Administration of SML either 50 or 100 mg/kg caused a dose-dependent marked decrease in the mRNA expression and protein concentration of  $\alpha$ -SMA compared with TAA group ( $P<0.0001$ ) (Fig. 7).

**Fig. 5** Sesamol (SML) reduces transforming growth factor  $\beta 1$  (TGF- $\beta 1$ ) gene expression and its protein concentration. **a** The mRNA expression of TGF- $\beta 1$ . **b** and **c** The protein concentration of TGF- $\beta 1$  using immunohistochemistry. Bars represent the number of positive cells per 10 high-power field in sections stained with anti-TGF- $\beta 1$  antibodies; arrows indicate cytoplasmic protein expression of TGF- $\beta 1$  that appears in hepatocytes around areas of fibrosis ( $\times$ : 400, bar 50). TAA thioacetamide. Values are expressed as mean  $\pm$  SEM ( $n=6$ ). \*Significance against control group ( $+++P<0.001$ ,  $++++P<0.0001$ ), \*significance against TAA group ( $****P<0.0001$ ), and  $^{\$}$ significance against SML 50 group ( $^{\$}P<0.01$ ,  $^{\$}$$$P<0.0001$ )



### SML decreases CK19 gene expression and its protein concentration


TAA caused a remarkable up-regulation in gene expression of CK19 compared with control group ( $P<0.0001$ ). Treatment with 50 or 100 mg/kg SML showed a dose-dependent significant down-regulation in CK19-mRNA expression compared with TAA group ( $P<0.0001$ ) (Fig. 8a). Immunohistochemical analysis revealed positively stained bile ducts in control group and a significant increase and expansion of CK19-positive staining in portal areas in TAA group compared with control group ( $P<0.0001$ ) due to proliferation of bile ductules and macrophage infiltrates the fibrous septa. Interestingly, both SML groups showed a marked decrease in CK19 protein concentration compared with TAA group ( $P<0.0001$ ) (Fig. 8b, c).

### SML reduces VEGF protein concentration

Protein concentration of VEGF markedly increased in TAA group compared with control group ( $P<0.0001$ ), while treatment with SML 50 or 100 mg/kg exhibited a significant reduction in VEGF protein concentration ( $P<0.0001$ ) compared with TAA group in a dose-dependent manner (Fig. 9).

## Discussion

Liver fibrosis is a threatening health problem leading to development of cirrhosis and liver cancer (Zhang et al. 2020). Induction of liver fibrosis by TAA is considered as a well-established model for evaluation of potential



**Fig. 6** Sesamol (SML) down-regulates small mothers against decapentaplegic3 (Smad3) gene expression and phosphorylated Smad3 (pSmad3) protein concentration and up-regulates Smad7 gene expression. **a** The mRNA expression of Smad3 ( $n=6$ ). **b** The protein level of pSmad3 ( $n=8$ ). **c** The mRNA expression of Smad7 ( $n=6$ ). TAA

thioacetamide. Values are expressed as mean  $\pm$  SEM. <sup>+</sup>Significance against control group ( $+++P<0.001$ ,  $++++P<0.0001$ ), <sup>\*</sup>significance against TAA group ( $**P<0.01$ ,  $****P<0.0001$ ), and <sup>§</sup>significance against SML 50 group ( $§P<0.05$ ,  $§§§P<0.001$ ,  $§§§§P<0.0001$ )


anti-fibrotic drugs as it produces hepatic disease in rodents resemble to those observed in humans (Yanguas et al. 2016).

In our study, TAA group showed a marked increase in liver index due to ECM accumulation in liver leading to an increase in relative liver weight. Serum ALT and AST activities were elevated in TAA group due to centrilobular necrosis caused by TAA that was accompanied by hepatic enzyme leakage in the serum. Additionally, serum total bilirubin was increased upon TAA injection due to impairment of liver detoxifying function. Moreover, TAA caused a remarkable decrease in serum albumin and total protein due to attenuation of synthetic function of the

hepatocytes; these results are in accordance with many studies (Czechowska et al. 2015; El-Mihi et al. 2017). SML administration caused improvement and restoration of hepatic functions, particularly in rats treated with the higher dose of SML (100 mg/kg). These results were further reinforced via the histopathological examination of liver tissues which indicated the ability of SML to protect against the pathological alterations in the liver architecture and collagen deposition after exposure to TAA.


Oxidative stress has been reported as a main molecular mechanism implicated in liver fibrosis induced by TAA. Releasing ROS and toxic metabolites after the metabolism





**Fig. 7** Sesamol (SML) suppresses alpha smooth muscle actin (α-SMA) gene expression and its protein concentration. **a** The mRNA expression of α-SMA. **b** and **c** The protein concentration of α-SMA using immunohistochemistry. Bars represent the number of positive cells per 10 high-power field in sections stained with anti-α-SMA


antibodies; arrows indicate positively stained cells (×: 400, bar 50). TAA thioacetamide. Values are expressed as mean ± SEM (n=6). +Significance against control group (++P<0.01, +++P<0.0001), \*significance against TAA group (\*\*\*\*P<0.0001), and §significance against SML 50 group (§P<0.05)




**Fig. 8** Sesamol (SML) decreases cytokeratin 19 (CK19) gene expression and its protein concentration. **a** The mRNA expression of CK19. **b** and **c** The protein concentration of CK19 using immunohistochemistry. Bars represent the number of positive cells per 10 high-power field in sections stained with anti-CK19 antibodies; black arrows indicate positively stained bile ducts and yellow arrows indicate

macrophage that infiltrates the fibrous septa ( $\times$ : 400, bar 50). TAA thioacetamide. Values are expressed as mean  $\pm$  SEM ( $n=6$ ). +Significance against control group (+++ $P<0.001$ , ++++ $P<0.0001$ ), \*significance against TAA group (\*\*\*\* $P<0.0001$ ), and  $^{\text{€}}$ significance against SML 50 group ( $^{\text{€}}$  $P<0.05$ )

(a)



(b)



**Fig. 9** Sesamol (SML) reduces vascular endothelial growth factor (VEGF) protein concentration. **a** and **b** The protein concentration of VEGF using immunohistochemistry. Bars represent the number of positive cells per 10 high-power field in sections stained with anti-VEGF antibodies; arrows indicate positively stained cells ( $\times$ : 400, bar 50). TAA thioacetamide. Values are expressed as mean  $\pm$  SEM ( $n=6$ ). \*Significance against control group ( $^{++}P<0.01$ ,  $^{++++}P<0.0001$ ), \*significance against TAA group ( $^{****}P<0.0001$ ), and  $^{\$}$ significance against SML 50 group ( $^{\$}P<0.05$ )

of TAA in liver resulted in hepatocyte necrosis, lipid peroxidation of cell membrane leading to MDA release, and imbalance of anti-oxidant defense system due to depletion of GSH (Tsai et al. 2010; Sukalingam et al. 2018). SML, particularly high-dose, revealed anti-oxidant properties as it abolished MDA accumulation and GSH reduction. This is in accordance with the reported anti-oxidant activity of SML that might be attributed to the up-regulation in the anti-oxidant enzymes such as catalase and superoxide dismutase (Gao et al. 2017; Majdalawieh and Mansour 2019; Abd Elrazik et al. 2021).

Treatment of rats with TAA increases autotaxin expression which in turn hydrolyzes lysophosphatidyl choline (LPC) into LPA (Lebda et al. 2018; Kaffe et al. 2019). LPA is a pro-fibrogenic factor in liver that is derived through autotaxin from hepatocytes and acts in a paracrine way in order to induce activation and proliferation of adjacent HSCs via binding to LPARs (1 and 3) (Kaffe et al. 2017; Eraky et al. 2018). It has been reported that LPA-LPAR1/LPAR3 signaling is included in fibrosis of liver and other tissues like renal and lung fibrosis (Pradère et al. 2008; Shea and Tager 2012; Kaffe et al. 2019). Our results showed significant elevation in LPAR1/LPAR3 expression in TAA-intoxicated rats. So, LPAR1/LPAR3 down-regulation could provide therapeutic benefits against hepatic fibrosis. According to our results, SML treatment showed marked down-regulation of LPAR1 and LPAR3 gene expressions in a dose-dependent manner proposing that SML might exert its anti-proliferative and anti-fibrogenic effects via targeting LPAR1/LPAR3 signaling pathway. These results are in agreement with previous study which correlated the protective effect against hepatic fibrosis with the down-regulation of LPARs (Eraky et al. 2018). To our knowledge, this study is the first one to investigate the role of SML in LPAR1/LPAR3 expression as a possible mechanism of action of its protective role against TAA-induced liver fibrosis.

CTGF is an important downstream molecule in the LPA-LPAR1/LPAR3 and TGF- $\beta$ 1 signaling pathways. CTGF up-regulation is considered as a central pathway participating in activation of HSCs during hepatic fibrosis (Cabello-Verrugio et al. 2011; Gan et al. 2011; Weiskirchen 2011; Ramazani et al. 2018). Induction of liver fibrosis by TAA causes a marked elevation in CTGF expression and that was in agreement with previous studies (Park et al. 2016; Algandaby et al. 2017; Eraky et al. 2018). SML, especially the high dose, markedly attenuated the increase of CTGF expression in TAA rats that might explain its anti-fibrotic effect.

TGF- $\beta$ 1 is a chief moderator in the development of liver fibrosis and a promising target to treat fibrosis. It is produced by hepatocytes, sinusoidal endothelial cells, and Kupffer cells in response to liver injury (Kajdaniuk et al. 2013; El-Mowafy et al. 2021). Additionally, LPARs are involved in the induction of TGF- $\beta$  expression (Li et al. 2017). TGF- $\beta$ 1 is an important factor in the activation of HSCs. Binding of active TGF- $\beta$ 1 to its receptors results in phosphorylation of cytoplasmic mediators Smad2 and Smad3 and consequently a Smad complex is formed with Smad4. Afterward, Smad complex translocates into the nucleus to control the transcription of the target genes such as  $\alpha$ -SMA and CTGF (Liu and Desai 2015). Activated HSCs express large amounts of  $\alpha$ -SMA which is a unique marker for HSCs. Smad7 negatively regulates TGF- $\beta$ 1/Smad3 signaling via binding to TGF- $\beta$  type I receptors and blocking Smad2/3 phosphorylation (Hu et al. 2018). Fibrotic cascade is triggered by

activation of TGF- $\beta$ 1/Smad3 pathway or loss of inhibitory Smad such as Smad7 (Hu et al. 2018; Dewidar et al. 2019). Our study showed stimulation of TGF- $\beta$ 1/Smad3 pathway in TAA group as indicated by an increase in TGF- $\beta$ 1, Smad3, pSmad3,  $\alpha$ -SMA expressions, and down-regulation of Smad7 expression. On the other hand, the enhanced TGF- $\beta$ 1, Smad3, pSmad3, and  $\alpha$ -SMA in TAA rats were dose-dependently attenuated after SML treatment in addition to elevation in the inhibitory Smad7. These results suggest that SML inhibits HSC activation by targeting TGF- $\beta$ 1/Smad3 pathway.

Although the activated HSC is the key effector cell type in liver fibrosis (Trautwein et al. 2015), there is another cell that can remarkably participate in fibrosis, including hepatic progenitor cells (HPCs) (Yovchev et al. 2008; Wang et al. 2009). Ductular reaction is the appearance of proliferating bile ductular structures upon liver injury (Sato et al. 2019). These proliferating ductules are assumed to emerge from HPCs (Falkowski et al. 2003; Roskams 2003; Roskams et al. 2004). A strong correlation between fibrosis extent and ductular reaction has been documented in previous studies (Clouston et al. 2005; Wood et al. 2014). The present study revealed that the gene expression and protein concentration of CK19, marker of HPCs and ductular reaction, was remarkably increased in the TAA group, but reduced after treatment with SML suggesting its ability to ameliorate ductular reaction.

Inflammation and hypoxia are important elements in angiogenesis. Liver tissue impacted by fibrosis exhibits a persistent inflammation and low level of oxygen due to ECM accumulation in liver parenchyma and disrupted blood vessels, leading to rise in major angiogenic factor (VEGF) expression and neovascularization (Elpek 2015; Zadorozhna et al. 2020). Bile ductules are continuously associated with microvessels, and ductule number matches with accompanied microvessel density, which gives an indication about the link between ductular reaction and microvessels in addition to angiogenesis (Gouw et al. 2006). Our study showed a marked elevation in VEGF protein concentration in TAA rats that was in accordance with other studies (Nakamura et al. 2014; Algandaby et al. 2017; Elfarawy et al. 2021). However, we found that treating TAA rats with SML significantly decreased the protein concentration of VEGF that may indicate its possible anti-angiogenic activity.

## Conclusion

Sesamol alleviated hepatic fibrosis induced by TAA in rats. The hepatoprotective effect of SML may be attributed to down-regulation of LPAR1/3 expression, inhibition of TGF- $\beta$ 1/Smad3 signaling, up-regulation of Smad7 expression, and eventually decreasing ductular reaction and angiogenesis.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00210-022-02259-7>.

**Author contribution** N. A.: conceptualization; performed the practical work, and data acquisition; formal analysis; writing the manuscript. M. E.: conceptualization; methodology; supervision; data interpretation and manuscript revision. M. M.: study concept and design; methodology supervision; data interpretation; critically reviewed the manuscript and provided valuable input.

All authors read and approved the manuscript and all data were generated in-house and that no paper mill was used.

**Funding** Open access funding provided by The Science, Technology & Innovation Funding Authority (STDF) in cooperation with The Egyptian Knowledge Bank (EKB).

**Data availability** All data generated or analyzed during this study are included in this article (and its supplementary information files).

## Declarations

**Ethics approval** This study protocol was approved by “Research Ethics Committee” Faculty of Pharmacy, Mansoura University, Mansoura City, Egypt (approval code: 2021–401), which is in accordance with “Guide for the Care and Use of Laboratory Animals, 8th edition” (National Research Council (US); Committee for the Update of the Guide for the Care and Use of Laboratory Animals, 2011).

**Consent to participate** Not applicable.

**Consent for publication** Not applicable.

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Abd Elrazik NA, El-Mesery M, El-Karef A, Eissa LA, El Gayar AM (2022) Sesamol upregulates death receptors and acts as a chemosensitizer in solid Ehrlich carcinoma model in mice. *Nutr Cancer* 74:250–264
- Algandaby MM, Breikaa RM, Eid BG, Neamatallah TA, Abdel-Naim AB, Ashour OM (2017) Icariin protects against thioacetamide-induced liver fibrosis in rats: implication of anti-angiogenic and anti-autophagic properties. *Pharmacol Rep* 69:616–624
- Cabello-Verrugio C, Córdova G, Vial C, Zúñiga LM, Brandan E (2011) Connective tissue growth factor induction by lysophosphatidic acid requires transactivation of transforming growth factor type  $\beta$  receptors and the JNK pathway. *Cell Signal* 23:449–457

- Chang CC, Lu WJ, Chiang CW, Jayakumar T, Ong ET, Hsiao G, Fong TH, Chou DS, Sheu JR (2010) Potent antiplatelet activity of sesamol in an in vitro and in vivo model: pivotal roles of cyclic AMP and p38 mitogen-activated protein kinase. *J Nutr Biochem* 21:1214–1221
- Clouston AD, Powell EE, Walsh MJ, Richardson MM, Demetris AJ, Jonsson JR (2005) Fibrosis correlates with a ductular reaction in hepatitis C: roles of impaired replication, progenitor cells and steatosis. *Hepatology* 41:809–818
- Czechowska G, Celinski K, Korolczuk A, Wojcicka G, Dudka J, Bojarska A, Reiter RJ (2015) Protective effects of melatonin against thioacetamide-induced liver fibrosis in rats. *J Physiol Pharmacol* 66:567–579
- Devidar B, Meyer C, Dooley S (2019) TGF- $\beta$  in hepatic stellate cell activation and liver fibrogenesis—updated 2019. *Cells* 8:1419
- Ebrahim RA, El-Mesery M, El-Karef A, Eissa LA (2018) Vitamin D potentiates anti-tumor activity of 5-fluorouracil via modulating caspase-3 and TGF- $\beta$ 1 expression in hepatocellular carcinoma-induced in rats. *Can J Physiol Pharmacol* 96:1218–1225
- El-Mihi KA, Kenawy HI, El-Karef A, Elsherbiny NM, Eissa LA (2017) Naringin attenuates thioacetamide-induced liver fibrosis in rats through modulation of the PI3K/Akt pathway. *Life Sci* 187:50–57
- El-Mowafy M, Elgaml A, El-Mesery M, Sultan S, Ahmed TA, Gomaa AI, Aly M, Mottawea W (2021) Changes of gut-microbiota-liver axis in hepatitis C virus infection. *Biology* 10:55
- Elnfarawy AA, Nashy AE, Abozaid AM, Komber IF, Elweshahy RH, Abdelrahman RS (2021) Vinpocetine attenuates thioacetamide-induced liver fibrosis in rats. *Hum Exp Toxicol* 40:355–368
- Elpek GÖ (2015) Angiogenesis and liver fibrosis. *World J Hepatol* 7:377
- Eraky SM, El-Mesery M, El-Karef A, Eissa LA, El-Gayar AM (2018) Silymarin and caffeine combination ameliorates experimentally-induced hepatic fibrosis through down-regulation of LPAR1 expression. *Biomed Pharmacother* 101:49–57
- Falkowski O, An HJ, Ianus IA, Chiriboga L, Yee H, West AB, Theise ND (2003) Regeneration of hepatocyte 'buds' in cirrhosis from intrabiliary stem cells. *J Hepatol* 39:357–364
- Gan L, Xue JX, Li X, Liu DS, Ge Y, Ni PY, Deng L, Lu Y, Jiang W (2011) Blockade of lysophosphatidic acid receptors LPAR1/3 ameliorates lung fibrosis induced by irradiation. *Biochem Biophys Res Commun* 409:7–13
- Gao XJ, Xie GN, Liu L, Fu ZJ, Zhang ZW, Teng LZ (2017) Sesamol attenuates oxidative stress, apoptosis and inflammation in focal cerebral ischemia/reperfusion injury. *Exp Ther Med* 14:841–847
- Gouw AS, van den Heuvel MC, Boot M, Slooff MJ, Poppema S, de Jong KP (2006) Dynamics of the vascular profile of the finer branches of the biliary tree in normal and diseased human livers. *J Hepatol* 45:393–400
- Helmy SA, El-Mesery M, El-Karef A, Eissa LA, El Gayar AM (2018) Chloroquine upregulates TRAIL/TRAILR2 expression and potentiates doxorubicin anti-tumor activity in thioacetamide-induced hepatocellular carcinoma model. *Chem Biol Interact* 279:84–94
- Helmy SA, El-Mesery M, El-Karef A, Eissa LA, El Gayar AM (2019) Thymoquinone upregulates TRAIL/TRAILR2 expression and attenuates hepatocellular carcinoma in vivo model. *Life Sci* 233:116673
- Hemalatha G, Pugalendi KV, Saravanan R (2013) Modulatory effect of sesamol on DOCA-salt-induced oxidative stress in uninephrectomized hypertensive rats. *Mol Cell Biochem* 379:255–265
- Higashi T, Friedman SL, Hoshida Y (2017) Hepatic stellate cells as key target in liver fibrosis. *Adv Drug Deliv Rev* 121:27–42
- Hu H-H, Chen D-Q, Wang Y-N, Feng Y-L, Cao G, Vaziri ND, Zhao Y-Y (2018) New insights into TGF- $\beta$ /Smad signaling in tissue fibrosis. *Chem Biol Interact* 292:76–83
- Ishak K, Baptista A, Bianchi L, Callea F, De Groote J, Gudat F, Denk H, Desmet V, Korb G, MacSween RN et al (1995) Histological grading and staging of chronic hepatitis. *J Hepatol* 22:696–699
- Jain R, Fischer S, Serra S, Chetty R (2010) The use of cytokeratin 19 (CK19) immunohistochemistry in lesions of the pancreas, gastrointestinal tract, and liver. *Appl Immunohistochem Mol Morphol* 18:9–15
- James J, Bosch KS, Aronson DC, Houtkooper JM (1990) Sirius red histophotometry and spectrophotometry of sections in the assessment of the collagen content of liver tissue and its application in growing rat liver. *Liver* 10:1–5
- Jayaraj P, Narasimhulu CA, Rajagopalan S, Parthasarathy S, Desikan R (2020) Sesamol: a powerful functional food ingredient from sesame oil for cardioprotection. *Food Funct* 11:1198–1210
- Jnaneshwari S, Hemshekhar M, Thushara R, Shanmuga Sundaram M, Sebastin Santhosh M, Sunitha K, Shankar R, Kemparaju K, Girish K (2014) Sesamol ameliorates cyclophosphamide-induced hepatotoxicity by modulating oxidative stress and inflammatory mediators. *Anti-Cancer Agents Med Chem* 14:975–983
- Kaffe E, Katsifa A, Xylourgidis N, Ninou I, Zannikou M, Harokopos V, Foka P, Dimitriadis A, Evangelou K, Moulas AN (2017) Hepatocyte autotaxin expression promotes liver fibrosis and cancer. *Hepatology* 65:1369–1383
- Kaffe E, Magkrioti C, Aidinis V (2019) Deregulated lysophosphatidic acid metabolism and signaling in liver cancer. *Cancers (Basel)* 11:1626
- Kajdaniuk D, Marek B, Borgiel-Marek H, Kos-Kudła B (2013) Transforming growth factor beta1 (TGFbeta1) in physiology and pathology. *Endokrynol Pol* 64:384–396
- Kaur IP, Saini A (2000) Sesamol exhibits antimutagenic activity against oxygen species mediated mutagenicity. *Mutat Res Genet Toxicol Environ Mutagen* 470:71–76
- Lebda MA, Sadek KM, Abouzed TK, Tohamy HG, El-Sayed YS (2018) Melatonin mitigates thioacetamide-induced hepatic fibrosis via antioxidant activity and modulation of proinflammatory cytokines and fibrogenic genes. *Life Sci* 192:136–143
- Li HY, Oh YS, Choi J-W, Jung JY, Jun H-S (2017) Blocking lysophosphatidic acid receptor 1 signaling inhibits diabetic nephropathy in db/db mice. *Kidney Int* 91:1362–1373
- Liu R-M, Desai LP (2015) Reciprocal regulation of TGF- $\beta$  and reactive oxygen species: a perverse cycle for fibrosis. *Redox Biol* 6:565–577
- Majdalawieh AF, Mansour ZR (2019) Sesamol, a major lignan in sesame seeds (*Sesamum indicum*): anti-cancer properties and mechanisms of action. *Eur J Pharmacol* 855:75–89
- Majdaeen M, Banaei A, Abedi-Firouzjah R, Gorji KE, Ataei G, Momeni F, Zamani H (2020) Investigating the radioprotective effect of sesamol oral consumption against gamma irradiation in mice by micronucleus and alkaline comet assays. *Appl Radiat Isot* 159:109091
- Mu M, Zuo S, Wu RM, Deng KS, Lu S, Zhu JJ, Zou GL, Yang J, Cheng ML, Zhao XK (2018) Ferulic acid attenuates liver fibrosis and hepatic stellate cell activation via inhibition of TGF- $\beta$ /Smad signaling pathway. *Drug Des Devel Ther* 12:4107–4115
- Nakamura I, Zakharia K, Banini BA, Mikhail DS, Kim TH, Yang JD, Moser CD, Shaleh HM, Thornburgh SR, Walters I (2014) Brivanib attenuates hepatic fibrosis in vivo and stellate cell activation in vitro by inhibition of FGF, VEGF and PDGF Signaling. *PLoS One* 9:e92273
- Nevzorova YA, Boyer-Diaz Z, Cubero FJ, Gracia-Sancho J (2020) Animal models for liver disease—a practical approach for translational research. *J Hepatol* 73:423–440
- Park HJ, Kim HG, Wang JH, Choi MK, Han JM, Lee JS, Son CG (2016) Comparison of TGF- $\beta$ , PDGF, and CTGF in hepatic fibrosis models using DMN, CCl4, and TAA. *Drug Chem Toxicol* 39:111–118

- Pradère JP, Gonzalez J, Klein J, Valet P, Grès S, Salant D, Bascands JL, Saulnier-Blache JS, Schanstra JP (2008) Lysophosphatidic acid and renal fibrosis. *Biochim Biophys Acta* 1781:582–587
- Ramazani Y, Knops N, Elmonem MA, Nguyen TQ, Arcolino FO, van den Heuvel L, Levtchenko E, Kuypers D, Goldschmeding R (2018) Connective tissue growth factor (CTGF) from basics to clinics. *Matrix Biol* 68:44–66
- Roehlen N, Crouchet E, Baumert TF (2020) Liver fibrosis: mechanistic concepts and therapeutic perspectives. *Cells* 9:875
- Roskams T (2003) Progenitor cell involvement in cirrhotic human liver diseases: from controversy to consensus. *J Hepatol* 39:431–434
- Roskams TA, Theise ND, Balabaud C, Bhagat G, Bhathal PS, Bioulac-Sage P, Brunt EM, Crawford JM, Crosby HA, Desmet V (2004) Nomenclature of the finer branches of the biliary tree: canals, ductules, and ductular reactions in human livers. *Hepatology* 39:1739–1745
- Sachdeva AK, Misra S, Kaur IP, Chopra K (2015) Neuroprotective potential of sesamol and its loaded solid lipid nanoparticles in ICV-STZ-induced cognitive deficits: behavioral and biochemical evidence. *Eur J Pharmacol* 747:132–140
- Sato K, Marzioni M, Meng F, Francis H, Glaser S, Alpini G (2019) Ductular reaction in liver diseases: pathological mechanisms and translational significances. *Hepatology* 69:420–430
- Shaker ME, Eisa NH, Elgaml A, El-Mesery A, El-Shafey M, El-Dosoky M, El-Mowafy M, El-Mesery M (2021) Ingestion of mannose ameliorates thioacetamide-induced intrahepatic oxidative stress, inflammation and fibrosis in rats. *Life Sci* 286:120040
- Shea BS, Tager AM (2012) Role of the lysophospholipid mediators lysophosphatidic acid and sphingosine 1-phosphate in lung fibrosis. *Proc Am Thorac Soc* 9:102–110
- Simo KA, Niemeyer DJ, Hanna EM, Swet JH, Thompson KJ, Sindram D, Iannitti DA, Eheim AL, Sokolov E, Zuckerman V, McKillop IH (2014) Altered lysophosphatidic acid (LPA) receptor expression during hepatic regeneration in a mouse model of partial hepatectomy. *HPB (oxford)* 16:534–542
- Sukalingam K, Ganesan K, Xu B (2018) Protective effect of aqueous extract from the leaves of *Justicia tranquebariensis* against thioacetamide-induced oxidative stress and hepatic fibrosis in rats. *Antioxidants* 7:78
- Trautwein C, Friedman SL, Schuppan D, Pinzani M (2015) Hepatic fibrosis: concept to treatment. *J Hepatol* 62:S15–24
- Tsai MK, Lin YL, Huang YT (2010) Effects of salvianolic acids on oxidative stress and hepatic fibrosis in rats. *Toxicol Appl Pharmacol* 242:155–164
- Wang P, Liu T, Cong M, Wu X, Bai Y, Yin C, An W, Wang B, Jia J, You H (2009) Expression of extracellular matrix genes in cultured hepatic oval cells: an origin of hepatic stellate cells through transforming growth factor beta? *Liver Int* 29:575–584
- Weiskirchen R (2011) CCN proteins in normal and injured liver. *Front Biosci (Landmark Ed)* 16:1939–1961
- Wood MJ, Gadd VL, Powell LW, Ramm GA, Clouston AD (2014) Ductular reaction in hereditary hemochromatosis: the link between hepatocyte senescence and fibrosis progression. *Hepatology* 59:848–857
- Xiang H, Lu Y, Shao M, Wu T (2020) Lysophosphatidic acid receptors: biochemical and clinical implications in different diseases. *J Cancer* 11:3519–3535
- Yanguas SC, Cogliati B, Willebrords J, Maes M, Colle I, van den Bossche B, de Oliveira CPMS, Andraus W, Alves VA, Leclercq I (2016) Experimental models of liver fibrosis. *Arch Toxicol* 90:1025–1048
- Yogalakshmi B, Viswanathan P, Anuradha CV (2010) Investigation of antioxidant, anti-inflammatory and DNA-protective properties of eugenol in thioacetamide-induced liver injury in rats. *Toxicology* 268:204–212
- Yovchev MI, Grozdanov PN, Zhou H, Racherla H, Guha C, Dabeva MD (2008) Identification of adult hepatic progenitor cells capable of repopulating injured rat liver. *Hepatology* 47:636–647
- Zadorozhna M, Di Gioia S, Conese M, Mangieri D (2020) Neovascularization is a key feature of liver fibrosis progression: anti-angiogenesis as an innovative way of liver fibrosis treatment. *Mol Biol Rep* 47:2279–2288
- Zeyada MS, Abdel-Rahman N, El-Karef A, Yahia S, El-Sherbiny IM, Eissa LA (2020) Niclosamide-loaded polymeric micelles ameliorate hepatocellular carcinoma in vivo through targeting Wnt and Notch pathways. *Life Sci* 261:118458
- Zhang CY, Yuan WG, He P, Lei JH, Wang CX (2016) Liver fibrosis and hepatic stellate cells: etiology, pathological hallmarks and therapeutic targets. *World J Gastroenterol* 22:10512–10522
- Zhang YN, Fowler KJ, Ozturk A, Potu CK, Louie AL, Montes V, Henderson WC, Wang K, Andre MP, Samir AE (2020) Liver fibrosis imaging: a clinical review of ultrasound and magnetic resonance elastography. *J Magn Reson Imaging* 51:25–42

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.