

Erratum

S. Beuchler* · R. Schneider* · C. Schwab**

Multiresolution weighted norm equivalences and applications

Published online: 15 November 2005
© Springer-Verlag 2005

Numer. Math (2004)
Digital Object Identifier (DOI) 10.1007/s00211-003-0491-8
Published online: April 8, 2004 – © Springer-Verlag 2004

Four equations in statements and proofs of Theorems 3.2 and 3.3 were printed with errors. The text below contains the correct equations.

Theorem 3.2 *The infinite matrix $M = ((\psi_k^l, \psi_{k'}^{l'})_w)_{(k,l);(k',l')}$ is bounded in l_2 .*

Proof We decompose the matrix M into $M = M_1 + M_2$ where the coefficients in M_2 are $(\psi_k^l, \psi_{k'}^{l'})_w$ iff $0 \in \text{supp } \psi_k^l \cap \text{supp } \psi_{k'}^{l'}$ and M_1 does not contain the interaction of wavelets which are both located at the point zero. By applying Theorem 3.1, Lemma 3.7 and the Schur Lemma to M_1 we have $\|M_1\|_2 \leq c$. From Lemma

* Supported by the DFG-Sonderforschungsbereich 393 “Numerische Simulation auf massiv parallelen Rechnern”.

** Supported by the TMR-project “Wavelets and Multiscale Methods in Numerical Simulation” of the European Union and by the swiss Government under Grant No. BBW 97.404.

S. Beuchler
Johann Radon Institute for Computational and Applied Mathematics, Austrian Academy of Science, Altenberger Strasse 69, 4040 Linz, Austria
E-mail: sven.beuchler@oeaw.ac.at

R. Schneider
Scientific Computing, Institute of Computer Science, Christian Albrecht University Kiel, Christian-Albrechts-Platz 4, 24098 Kiel, Germany
E-mail: rs@numerik.uni-kiel.de

C. Schwab (✉)
Seminar für Angewandte Mathematik, Eidgenössische Technische Hochschule, 8092 Zürich, Switzerland
E-mail: schwab@math.ethz.ch

3.8 we have $\| M_2 \|_1 \leq c$ and $\| M_2 \|_\infty \leq c$ which shows $\| M_2 \|_2 \leq c$. Hence, the assertion is proven. \square

We show now the equivalence of the L_w^2 norm of a function

$$u = \sum_{l=l_0}^{\infty} \sum_k u_k^l \psi_k^l \in L_w^2((0, 1))$$

with its discrete l_w^2 norm of the coefficients $(u_k^l)_{(k,l)} \in \mathbb{R}$, i.e.

$$\| \|u_k^l\| \| \|_w^2 := \sum_l \sum_k w^2(2^{-l}k) |u_k^l|^2.$$

Theorem 3.3 *Let us assume that Assumptions 3.1 and 3.2 are valid. For any function $u = \sum_{l=l_0}^{\infty} \sum_k u_k^l \psi_k^l \in L_w^2((0, 1))$ holds*

$$\| u \|_w^2 \approx \| \|u_k^l\| \| \|_w^2.$$

Proof From Theorem 3.2 we conclude

$$\begin{aligned} \| u \|_w^2 &= \sum_{l,l'} \sum_{k,k'} u_k^l u_{k'}^{l'} w(2^{-l}k) w(2^{-l'}k') (\psi_k^l, \psi_{k'}^{l'})_w \\ &\leq \| M \|_2 \sum_l \sum_k \left(|u_k^l| w(2^{-l}k) \right)^2 \leq c \| \|u_k^l\| \| \|_w^2. \end{aligned}$$

To prove the lower estimate we consider the dual system

$$\tilde{v} = \sum_l \sum_k \tilde{v}_k^l \tilde{\psi}_k^l = G(\tilde{v}_k^l)$$

in the dual space $L_{w^{-1}}^2((0, 1))$. We denote by \tilde{M} the mass matrix of the dual wavelet basis $\tilde{\psi}_k^l$ with respect to the $L_{w^{-1}}^2((0, 1))$ innerproduct. Then, by the same arguments

$$\| \tilde{v} \|_{w^{-1}}^2 \leq \| \tilde{M} \|_2 \| \| \tilde{v}_k^l \| \|_{w^{-1}}^2.$$

This means $G : l_{w^{-1}}^2 \rightarrow L_{w^{-1}}^2((0, 1))$ is bounded. Therefore, the adjoint operator $G^* : L_w^2((0, 1)) \rightarrow l_w^2$ is bounded, too. G^* is explicitly given by

$$G^* u := \left(\langle u, \tilde{\psi}_k^l \rangle \right)_{l,k} = (u_k^l)_{l,k}$$

which proves the lower bound. \square