

Chris C. Streeter · John Hennen · Yong Ke ·
J. Eric Jensen · Ofra Sarid-Segal · Leanne E. Nassar ·
Clifford Knapp · Angela A. Meyer · Tae Kwak ·
Perry F. Renshaw · Domenic A. Ciraulo


Prefrontal GABA levels in cocaine-dependent subjects increase with pramipexole and venlafaxine treatment

Published online: 7 March 2006
© Springer-Verlag 2006

Psychopharmacology (2005) 182:516–526

Unfortunately, Figs. 2 and 3 were published with errors. The correct figures are given here.

Fig. 2 Individual prefrontal lobe GABA levels in mmol/kg showing differences between before treatment (Scan 1) and after treatment (Scan 2) by group



The online version of the original article can be found at <http://dx.doi.org/10.1007/s00213-005-0121-5>.

C. C. Streeter (✉) · O. Sarid-Segal · C. Knapp · T. Kwak ·
P. F. Renshaw · D. A. Ciraulo
Division of Psychiatry, Boston University School of Medicine,
85 E. Newton St. M912E,
Boston, MA, USA
e-mail: streeter@bu.edu
Tel.: +1-617-6388008
Fax: +1-617-6388007

C. C. Streeter · O. Sarid-Segal · A. A. Meyer · D. A. Ciraulo
Boston VA Healthcare System,
Boston, MA, USA

C. C. Streeter · J. Hennen · Y. Ke · J. E. Jensen · L. E. Nassar ·
P. F. Renshaw
McLean Hospital,
Belmont, MA, USA

C. C. Streeter · J. Hennen · Y. Ke · J. E. Jensen · P. F. Renshaw ·
D. A. Ciraulo
Department of Psychiatry, Harvard Medical School,
Boston, MA, USA




Fig. 3 Changes in brain GABA levels by treatment group. *The overall statistical model for the effect of drug treatment on brain GABA levels together with the baseline GABA adjustment was strongly significant ($p=0.002$). **GABA change-from-baseline in the pramipexole treatment group differed significantly from GABA change in the placebo group, ($p=0.031$)