The effects of systemic, intrastriatal, and intrapallidal injections of caffeine and systemic injections of A_{2A} and A_1 antagonists on forepaw stepping in the unilateral 6-OHDA-lesioned rat

John E. Kelsey · Nicole A. Langelier · Brad S. Oriel ·
Catherine Reedy

Published online: 23 October 2008
© Springer-Verlag 2008

Erratum to: Psychopharmacology
DOI 10.1007/s00213-008-1319-0

The original version of this article unfortunately contained a mistake. In the second to the last sentence of the Results section of the Abstract, the sentence should read: “The selective A_{2A} antagonist SCH-58261 (2 mg/kg) also increased stepping by 13% and enhanced the therapeutic effect of L-DOPA, whereas the selective A_1 antagonist 8-cyclopentyltheophylline (3–7 mg/kg) and A_1 agonist N^6 -cyclopentyladenosine (0.03–0.2 mg/kg) had no effect.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00213-008-1319-0>.

J. E. Kelsey (✉) · N. A. Langelier · B. S. Oriel
Program in Neuroscience, Bates College,
Lewiston, ME 04240, USA
e-mail: jkelsey@bates.edu

J. E. Kelsey · C. Reedy
Department of Psychology, Bates College,
Lewiston, ME 04240, USA