

## Erratum to: A preliminary double-blind, placebo-controlled randomized study of baclofen effects in alcoholic smokers

Lorenzo Leggio<sup>1,2,3</sup> · William H. Zywiak<sup>4,5,6</sup> · Steven M. Edwards<sup>7</sup> · Jennifer W. Tidey<sup>6</sup> · Robert M. Swift<sup>6,8,9</sup> · George A. Kenna<sup>6</sup>

Published online: 26 March 2015  
© Springer-Verlag Berlin Heidelberg 2015

**Erratum to: Psychopharmacology**  
**DOI 10.1007/s00213-014-3652-9**

The original version of this manuscript contained a mistake in the text. The sentence 'BACL significantly reduced the % days of abstinence from alcohol-tobacco co-use (BACL, 12.1±2.0 vs. PLA, 3.5±2.2 (M±SE); F(1,197.6)=8.27, p=0.004; Fig. 1a).' should read 'BACL significantly increased the % days of abstinence from alcohol-tobacco co-use (BACL, 12.1±2.0 vs. PLA, 3.5±2.2 (M±SE); F(1,197.6)=8.27, p=0.004; Fig. 1a).'

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00213-014-3652-9>.

✉ Lorenzo Leggio  
lorenzo.leggio@nih.gov

- <sup>1</sup> Section on Clinical Psychoneuroendocrinology and Neuropsychopharmacology, Laboratory of Clinical and Translational Studies, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, 10 Center Drive (10CRC/15330) MSC 1108; Room 1-5429, Bethesda, MD 20892-1108, USA
- <sup>2</sup> Intramural Research Program, National Institute on Drug Abuse, National Institutes of Health, Baltimore, MD, USA
- <sup>3</sup> Department of Behavioral and Social Sciences, Center for Alcohol and Addiction Studies, Brown University, Providence, RI, USA
- <sup>4</sup> Decision Sciences Institute, P.I.R.E, Pawtucket, RI, USA
- <sup>5</sup> Butler Hospital, Providence, RI, USA
- <sup>6</sup> Department of Psychiatry and Human Behavior, Center for Alcohol and Addiction Studies, Brown University, Providence, RI, USA
- <sup>7</sup> Department of Psychology, University of Nebraska-Lincoln, Lincoln, NE, USA
- <sup>8</sup> Veterans Affairs Medical Center, Providence, RI, USA
- <sup>9</sup> Roger Williams Medical Center, Providence, RI, USA