



## Correction to: Fluorescence anisotropy study of radiation-induced DNA damage clustering based on FRET

Ken Akamatsu<sup>1</sup> · Naoya Shikazono<sup>1</sup> · Takeshi Saito<sup>2</sup>

Received: 9 March 2021 / Accepted: 9 March 2021 / Published online: 13 March 2021  
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

**Correction to: Analytical and Bioanalytical Chemistry**  
<https://doi.org/10.1007/s00216-020-03082-w>

The authors would like to call the reader's attention to the fact that, unfortunately, there was an error regarding figure 3 in this manuscript; please find the correct figure below. The apparent base separation in a cluster produced by 2.0 MeV/u  $^4\text{He}^{2+}$ , 0.52 MeV/u  $^4\text{He}^{2+}$ , and 0.37 MeV/u  $^{12}\text{C}^{5+}$  beams demonstrated in the “Abstract” and the “Results and Discussion” sections are changed to be ca. 21, 19, and 18 bp, respectively.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s00216-020-03082-w>

---

✉ Ken Akamatsu  
akamatsu.ken@qst.go.jp

<sup>1</sup> DNA Damage Chemistry Research Group, Institute for Quantum Life Science, National Institutes for Quantum and Radiological Science and Technology (QST), 8-1-7 Umemidai, Kizugawa, Kyoto 619-0215, Japan

<sup>2</sup> Division of Radiation Life Science, Institute for Integrated Radiation and Nuclear Science, Kyoto University, 2 Asashiro-Nishi, Kumatori, Sennan, Osaka 590-0494, Japan

