



The influence of weather conditions on bioactive compound content in sorghum grain

Anna Przybylska-Balcerek¹ · Jakub Frankowski² · Kinga Stuper-Szablewska¹

Received: 12 July 2019 / Revised: 14 October 2019 / Accepted: 19 October 2019 / Published online: 5 November 2019
© The Author(s) 2019

Abstract

Sorghum is the fifth most important cereal in the world in terms of the cropped area. It is mainly grown for feeding animals and it is also used in the food industry. Sorghum grain is generally a rich source of antioxidants such as polyphenols and carotenoids. For this reason, it is considered as a good source of bioactive food components and it has health-promoting properties. Sorghum is a gluten-free cereal grown in many regions worldwide, primarily in the tropical and subtropical regions. However, new hybrids and forms of sorghum are capable to produce seeds also in temperate climate. The aim of this study was to conduct the influence of weather conditions on bioactive compound content in sorghum grain. The quantitative analysis of selected bioactive compounds, such as phenolic acids, flavonoids, carotenoids, and phytosterols, was carried out. The tested material comprised grain of two varieties: ‘Sweet Susana’ and ‘Sweet Caroline’, which have different color of grain: red and white. The research material was obtained from growing seasons 2016–2018. Quantitative analysis of free phenolic acids, total carotenoids, and total phytosterols was performed by ultra-performance liquid chromatography (UPLC) after prior basic hydrolysis followed by acid. An ultra-efficient liquid chromatograph coupled with an absorption-based detector (UPLC-PDA) was used for these analyses. The results showed the variability of the content of bioactive compounds depending on weather conditions.

Keywords Sorghum grain · Phenolic compounds · Flavonoids · Carotenoids · Temperate climate

Introduction

Sorghum (*Sorghum* Moench), which belongs to the family of *Poaceae*, is an annual plant with C4 photosynthesis cycle [1, 2]. It is the fifth most important cereal in the world in terms of the cropped area [3, 4], because it is considered to be one of high nutritional value food crops [5–9]. The varieties of primary economic importance are *Sorghum bicolor* L. and *Sorghum sudanense* L., as well as numerous hybrids [10–12], which are also valuable animal fodder [13–16].

Furthermore, sorghum is used in various branches of industry [17, 18]. For example, its biomass can be useful for producing energy [19–22]. Stalks and leaves have high calorific value [23] and they are efficient solid biofuels after forming in bales [24]. Moreover, stems and grain are a feedstock for bioethanol production [23, 25–28]. In addition, biogas could also be produced from the biomass processing residue [29, 30]. Therefore, every piece of sorghum biomass can be effectively used for various purposes [23].

In Europe, various subspecies of sorghum have been grown sporadically so far, mainly for the production of animal feed or biofuels, due to the lack of adaptation of the varieties to the prevailing climate. However, intensive breeding in recent years has allowed the registration of many different hybrids and sorghum forms, which are capable of producing seeds in a temperate climate [31–33]. Continuous global warming and further development of genetic engineering make sorghum cultivation possible and profitable also in Poland, where sorghum was mainly used by the feed industry. But now, the grain can be produced successfully [23, 34].

✉ Anna Przybylska-Balcerek
ania_przybylska18@wp.pl

¹ The Department of Chemistry, The Faculty of Wood Technology, Poznań University of Life Sciences, ul. Wojska Polskiego 75, 60-101 Poznań, Poland

² The Department of Breeding and Agricultural Technology for Fibrous and Energy Plants, Institute of Natural Fibres and Medicinal Plants, ul. Wojska Polskiego 71b, 60-630 Poznań, Poland

In terms of chemical composition, sorghum is essentially similar to maize and millet [5, 34–40]. The main ingredient of sorghum is starch, sugars, protein, and lipids [23, 34–40]. Among micro and macroelements, the largest share is potassium, phosphorus, magnesium, calcium, and small amounts of iron and other chemical elements, but it also depends on many agrotechnical factors and cultivar [4, 41, 42]. It is also a good source of bioactive nutrients, e.g., vitamins, micro- and macronutrients, as well as non-food ingredients, e.g., phenolic acids, flavonoids, carotenoids, and phytosterols [5, 8, 34], with strong antioxidant properties.

Phenolic acids are aromatic organic compounds classified to the group of polyphenols [43–45]. They can be found in many plant species [45–49]. They are distinguished based on the carbon skeleton structure into two groups: derivatives of benzoic acid and derivatives of cinnamic acid [50, 51]. Moreover, sorghum grain also contains flavonoids. They are natural pigments responsible for proper functioning of the enzymatic system in plant cells. Their structure is based on three rings, within which modifications lead to the formation of various flavonoid compounds, e.g., flavonols, flavanone, and flavones [34, 52–55]. Another group of bioactive compounds found in sorghum grain comprises carotenoids. They may be found in the acyclic, monocyclic, or bicyclic forms. Their color ranging from yellow to red is dependent on the number of double bonds. In addition to the content of a wide range of phenolic compounds, sorghum seeds are also a rich source of phytosterols. We are known about 200 types of sterols marked in plants. Plant sterols are structural and functional analogs of cholesterol synthesized by plants. These compounds are part of plant cell membranes. These are 28- or 29-carbon polycyclic alcohols [56, 57]. Antioxidative properties of bioactive compounds are primarily based on antioxidant activity, which is intensified during the stress factors on the plant. Bioactive compounds such as phenolic acids, flavonoids, carotenoids, and phytosterols in sorghum inhibit the action of free radicals protecting plant cells, particularly chloroplasts against harmful effects of reactive oxygen species (ROS) [58–62].

Due to the presence of potentially valuable chemical components in sorghum grain, it can be successfully used as a substitute for conventional cereals in the production of functional food [34]. Notwithstanding, there is a need to investigate the concentration, composition, and value of its nutrients [12]. Concentrations of bioactive compounds in sorghum grain depend, e.g., on the cultivar, atmospheric conditions and on biotic stressors. Moreover, the beneficial effect of bioactive compounds on physiological functions in plant and consumer depends on their antioxidant action [63–65]. Literature reports on the contents of bioactive compounds in sorghum grain are only fragmentary [34]. In view of the above, studies were conducted on red and white grain

of sorghum grown in Poland to conduct quantitative analyses of selected bioactive compounds.

Therefore, the aim of this study was to conduct the influence of weather conditions on bioactive compound content in sorghum grain. The quantitative analysis of selected bioactive compounds: antioxidant activity applying an improved 2,2-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS^{•+}) radical cation decolorization, total phenolic acids content, and carotenoids consisted in alkaline and acid hydrolysis, followed by analyses using an ultra-performance liquid chromatograph coupled with a photodiode array detector (UPLC-PDA) were carried out. As a result, the possibility of using sorghum grain for food purposes cultivated in temperate climate was demonstrated.

Raw material

The research material was the grain of two cultivars of sorghum (*Sweet Susana—red* and *Sweet Caroline—white*) cultivated in 2016, 2017, and 2018. The experimental plots were located at the Sary Sielec Experimental Station in Poland (51° 39' N, 17° 10' E), which belongs to the Institute of Natural Fibers and Medicinal Plants (Instytut Włókien Naturalnych i Roślin Zielarskich). The experiments were carried out in the area with low precipitation (< 550 mm/year), on a soil of medium agricultural usefulness soil and good abundance of nutrients. Every year, the experimental plots were located next to the last position, but not in the same place to avoid the monoculture.

Sorghum was sown after triticale at the following fertilization level (in kg·ha⁻¹): 100 N, 60 P₂O₅, 120 K₂O, and 30 MgO. Soil cultivation from autumn to sowing seeds in first days of May was carried out in accordance with the principles of good agricultural practice. Neither before sowing nor during the growing season was any chemical plant protection products used. The weeds were removed mechanically at the initial stage of growth. Sorghum was harvested in the last days of September.

Testing methods

The weather conditions were measured by collecting the following data: daily participation, the lowest, the highest, and medium daily temperature. The measurements were made in a covered weather station located 2 m above the ground.

Grain samples of 100 g were collected for chemical analyses. Sorghum grain was ground in a WZ-40 laboratory mill.

Total polyphenolic contents

Samples of 50 g were collected for analyses of polyphenols. Samples were ground using a laboratory mill (WZ-1). Total phenolic compounds were extracted with 80% MeOH.

Samples of 10 g after flooding with 100 ml MeOH were placed in an ultrasound bath for 30 min; next, the precipitate was collected to distillation flasks and the extraction process was repeated three times. Next, the combined extracts were evaporated to dryness in an evaporator. Phenolic compounds were transferred quantitatively to a vial using MeOH and dried in a stream of nitrogen. The amount of 0.5 ml deionized water and 0.125 ml Folin–Ciocalteu reagent (Fluka) was added to 0.125 ml extract, and after 6 min, the mixture was supplemented with 1.25 ml 7% aqueous Na₂CO₃ solution and 1 ml deionized water. After 90 min, absorbance will be read at a wavelength of 760 nm in relation to water (Helios spectrophotometer Thermo Electron Corp.). Results were expressed in mg gallic acid/kg d.m. sample [66, 67].

ABTS⁺ method

For ABTS⁺ generation from ABTS salt, 3 mM of K₂S₂O₈ was reacted with 8 mM ABTS salt in distilled, deionized water for 16 h at room temperature in the dark. The ABTS⁺ solution was then diluted with pH 7.4 phosphate buffer solution containing 150 mM NaCl (PBS) to obtain an initial absorbance of 1.5 at 730 nm. Fresh ABTS⁺ solution was prepared for each analysis. Reaction kinetics was determined over a 2 h period with readings every 15 min. Reactions were complete in 30 min. Samples and standards (100 µm) were reacted with the ABTS⁺ solution (2900 µm) for 30 min. Trolox was used as a standard. Results were expressed in ABTS⁺ (µmol/TROLOX/kg) d.m. sample [66, 67].

Determination of carotenoids

Carotenoid isolation and quantification were performed by Acquity UPLC (Waters, USA) in grain samples by the saponification method. Carotenoids extracts were obtained from ground beans (0.4 mg) which were triturated with a mixture of acetone and petroleum ether (1: 1). Then, after separation of the plant tissue, the acetone and the hydrophilic fraction were removed from the extract by washing with water. As a result, the ether extract was obtained with a mixture of carotenoid pigments. The extract thus prepared was concentrated in a vacuum evaporator at 35 °C until an oily residue was obtained, then digested in 2 ml of methanol (Merck), and subjected to chromatographic analysis. Total carotenoids were determined using Acquity UPLC (Waters, USA) with a Waters Acquity PDA detector (Waters, USA). Chromatographic separation was performed on a Acquity UPLC[®] BEH C18 column (100 mm × 2.1 mm, particle size 1.7 µm) (Waters, Ireland). Elution was carried out using solvent A—methanol, B—water, and tert-butyl methyl ether (TBME). A gradient was applied at a flow of 0.4 ml/min. The column and samples were thermostated, the column temperature was 30° C, and the test temperature was 10° C.

During the analysis, the solutions were degassed in a Waters device. The injection volume was 10 µl. The registration was carried out at a wavelength $\lambda = 445$ nm [67, 68].

Chemical analysis of phytosterols

Samples containing 100 mg of ground grains were placed into 17-ml culture tubes, suspended in 2 ml of methanol, treated with 0.5 ml of 2 M aqueous sodium hydroxide, and tightly sealed. The culture tubes were then placed within 250-ml plastic bottles, tightly sealed and placed inside a microwave oven (Model AVM 401/1WH, Whirlpool, Sweden) operating at 2450 MHz and 900 W maximum output. Samples were irradiated (370 W) for 20 s and after about 5 min for an additional 20 s. After 15 min, the contents of culture tubes were neutralized with 1 M aqueous hydrochloric acid, 2 ml MeOH were added, and extraction with pentane (3 × 4 ml) was carried out within the culture tubes. The combined pentane extracts were evaporated to dryness in a nitrogen stream. Before analysis, samples were dissolved in 4 ml of MeOH, filtered through 13-mm syringe filters with a 0.5 mm pore diameter (Fluoropore Membrane Filters, Millipore, Ireland), and evaporated to dryness in a N₂ stream. The sample extract was dissolved in 1 ml of MeOH and 50 µl were analyzed by Acquity H class UPLC system equipped with a Waters Acquity PDA detector (Waters, USA). Chromatographic separation was performed on a Acquity UPLC[®] BEH C18 column (100 mm × 2.1 mm, particle size 1.7 µm) (Waters, Ireland) eluted with methanol/acetonitrile/water (85:10:5) at a flow rate of 0.4 ml/min. Total sterols was detected with a Waters Acquity PDA detector (Waters, USA) set at 282 nm [69].

Statistical analysis

The results of the chemical analyses were analyzed statistically by means of the STATISTICA v 8.0 software. Tukey's multiple comparison procedure was used to compare the contents of individual metabolites in the samples. Identical letters in rows or columns denote the lack of differences at a significance level $\alpha = 0.05$.

Results and discussion

Due to the fact that sorghum was sown in the first days of May and collected in the last days of September (in the period without risk of frost); 5 months were included in the climate date (Table 1 and Fig. 1). Compared to the long-term average temperature (1960–2000) for this region, the monthly temperatures in the spring and summer 2016–2018 were generally higher. Nevertheless, average temperatures within 10 days fluctuated significantly,

Table 1 Climate data in the decades of the months May–September in the triennium 2016–2018

Month	Decade	2016		2017		2018	
		Average temp. (°C)	Precipitation (mm)	Average temp. (°C)	Precipitation (mm)	Average temp. (°C)	Precipitation (mm)
May	I	11.1	25.7	7.9	19.1	14.0	14.7
	II	12.3	3.2	14.2	0.0	14.4	60.9
	III	17.3	48.7	16.5	15.8	17.1	1.2
June	I	18.0	28.1	16.2	10.4	20.0	13.5
	II	16.6	23.3	18.2	15.1	18.7	3.3
	III	20.1	12.2	19.8	28.8	14.8	50.5
July	I	17.5	15.5	17.8	21.8	19.8	0.0
	II	17.8	101.0	18.9	41.4	18.2	92.0
	III	21.8	2.0	19.8	29.5	23.9	0.1
August	I	17.7	41.3	21.5	29.3	25.1	0.0
	II	17.4	0.8	19.5	36.5	21.4	5.0
	III	18.7	11.0	18.5	0.2	17.9	13.6
September	I	19.1	4.0	14.5	44.3	17.6	3.1
	II	17.6	4.8	12.8	8.8	17.8	8.5
	III	12.4	0.2	11.2	22.7	12.3	37.0




Fig. 1 The temperature and precipitation during sorghum vegetation period

which is typical of central Poland. The inflow of humid and relatively cool masses of air from the Atlantic Ocean intertwines with the continental wind, which promises sunny and hot weather in the summer. The hottest part of the year in Poland is above all the turn of July and August. During these days, the daytime temperature exceeds 30° C and fluctuates around 20 at dawn. After a few days of heat, the weather changes suddenly and the temperature drops. Rainfall is not evenly distributed and is mainly caused by local storms. From time to time, as in the second decade of July 2016 and 2018, there were several days of intense rainfall. This situation undoubtedly influenced the growth of sorghum, because the puddles lasted for a long time on the surface of the plot. However, Wielkopolska, where the Experimental Station is located, is known as an area with water shortages. In summer, there are periods of drought each year. Weather conditions in the summer season do not have such a significant impact on bioactive compounds in sorghum grains, such as insolation and temperature in September, when the kernels ripen. The harvest time is usually much colder than the plant growth time. The most important thing is to do it before the beginning of even a light night frost, which immediately damages sorghum. On the basis of this study, unstable atmospheric conditions were observed during the growing season of sorghum. This could affect the quality and quantity of crops. Therefore, during these experiments, attention was paid to the influence of stress factors on the yield quality. The determinant of the severity of symptoms of oxidative stress was the concentration of selected bioactive compounds and the measurement of their antioxidant activity. Comparing the data from Table 1 with Table 2, it was found that both the air temperature and the amount of rainfall are important during the vegetation period of the plant. Sorghum is a thermophilic plant; therefore, the vegetation period of this plant in a temperate climate was from May to the end of September. It is important that at this time, optimal conditions for the correct yielding of the plant are met. The present study was conducted in a 3-year cycle, in which significant differences in the intensity of oxidative stress (biotic and abiotic) were noted. In the defense reaction to oxidative stress in plant cells, antioxidant compounds are formed that capture free radicals. The concentration of bioactive compounds was different for the two tested sorghum varieties, in individual years. In 2016, the average air temperature throughout the growing season was even, and the amount of rainfall was associated with sudden downpours. In 2017, the temperature and precipitation were quite stable. Another period is 2018. While high temperatures in the commune were dominated by long-lasting drought and rare rains. Each of the vegetation periods discussed was specific, but the sorghum grew under all experimental conditions. In this research,

Table 2 Contents of bioactive compounds in sorghum grain [mg/kg]

Chemical compound	Vegetation period	Sorghum grain	
		<i>Sweet Susana—red</i>	<i>Sweet Caroline—white</i>
ABTS ⁺ (μ molTROLOX/kg)	2016	789a	921b
	2017	659a	893b
	2018	703a	916b
TPC (mg GAE/100 g)	2016	549a	822b
	2017	496a	913b
	2018	548a	879b
Carotenoids (mg/kg)	2016	124a	428b
	2017	116a	571b
	2018	132a	499a
Phytosterols (mg/kg)	2016	26a	29a
	2017	27a	30a
	2018	23a	41a

ABTS⁺(μ molTROLOX/kg): antioxidant activity applying an improved 2,2-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid; TPC (mg GAE/100 g): total phenolic compounds

a, b: These same letters in rows mean no significant statistical differences, $p \leq 0.05$

temperature and rainfall were not connected with level of bioactive compound in sorghum. This is new information about these sorghum varieties grown in Poland.

As part of the present study, 18 samples of white sorghum grains (*Sweet Susana*) and 18 samples of white sorghum grain (*Sweet Caroline*) were analyzed. Based on the obtained results, the presence of selected bioactive compounds was found in all samples and significant differences in the content of these compounds were found. The amount and type of antioxidants in plants depends on the species, variety, place of growth, insolation, and the period in which the crops were harvested [3, 21]. The plants rich in bioactive compounds include fruits, vegetables, herbs, grain, as well as sorghum [70]. As part of this study, the results of free phenolic acid content, total carotenoid concentration as well as antioxidant activity were obtained. Based on the results presented in Table 2, it was found that the antioxidant activity of ABTS⁺ bioactive compounds was a higher sample of *Sweet Caroline* compared to *Sweet Susana*. The highest antioxidant activity was found in *Sweet Caroline* seeds in 2016 and 2018, but there were not significant statistical differences between years. ABTS⁺'s antioxidant activity was lower by 15 and 23%, respectively, in *Sweet Susana*. Other studies were conducted by Xiong et al. [71]; they tested the activity of ABTS⁺ and DPPH extracts and raw sorghum grains. Studies have shown that the content of total phenolic acids and other polyphenols was significantly higher in extracts than in raw sorghum. However, the antioxidant activity has not changed, suggesting that the concentration of

phenolic compounds in white sorghum may affect the level of antioxidant activity. In contrast, in colored sorghum, other compounds may contribute to higher antioxidant activity. It is also worth mentioning that the antioxidant activity of ABTS⁺ bioactive compounds in sorghum is higher compared to ABTS⁺ grains of wheat and triticale cultivated in temperate climates [72].

In this work, the concentration of TPC in two types of sorghum was also determined. Phenolic acids are responsible for preventing reactions of oxidation, cell degradation, and plant death. The production of phenolic acids depends on the intensity of oxidative stress caused by biotic and abiotic factors. Based on the experiments carried out under the same atmospheric conditions, a higher concentration of TPC in Sweet Caroline was found compared to Sweet Susana in each crop year. The average TPC concentrations varied, respectively, in 2016 by 33%, in 2017, by 45%, and in 2018 by 37%. In addition to TPC, other bioactive compounds, such as carotenoids, are also found in sorghum grains. Carotenoids perform very important functions in the plant world. They are responsible for the stability of lipid membranes; they participate in the accumulation of light in the process of photosynthesis, as well as protection against the process of photooxidation induced by reactive oxygen species formed during the chlorophyll excitation in the process of photosynthesis. The antioxidant activity of carotenoids on lipid membranes depends on their orientation, location, and organization in membranes. The antiradical activity of carotenoids is based on two mechanisms, such as: the first involves electron transfer and the second on the formation

of adducts with peroxide radicals. During this study, it was found that Sweet Caroline is a richer source of carotenoids than Sweet Susana. At Sweet Caroline in 2016, 71% more carotenoids were determined compared to the second tested variety. In 2017, the difference was 80%, and in 2018, it was 73%.

Statistical analyses show that there was no correlation between years. John et al., based on their studies, noted a higher concentration of carotenoids, especially in brown beans. They also found that the main bioactive compounds in sorghum beans are phenolic acids, flavonoids, and tannins [73, 74]. Similar studies were conducted by Przybylska-Balcerek et al. in 2019 [68], and they found the presence of many phenolic acids, flavonoids, and carotenoids. The presence of bioactive compounds, including polyphenols and their antioxidant activity, was also touched by Morais Cardoso et al., Dykes and Rooney, as well as Stefoska-Needham et al., the assertion that polyphenols are present in sorghum grain [8, 74, 75].

Discriminant analysis

The discriminant analysis was used to identify among the analyzed bioactive compounds, those which content characterizes sorghum red and white (Figs. 2, 3, 4). On this basis, it was found all identified bioactive compounds are characterized by the highest discriminatory power for the entire population of analyzed samples. In the case of sorghum Sweet Caroline, it was found more concentration analyzed compounds.




Fig. 2 Principal component analysis (PCA) and discriminant analysis (DA) of various types of sorghum-bioactive compounds—2016




Fig. 3 Principal component analysis (PCA) and discriminant analysis (DA) of various types of sorghum-bioactive compounds—2017




Fig. 4 Principal component analysis (PCA) and discriminant analysis (DA) of various types of sorghum-bioactive compounds—2018

Based on the results of chemical analyses and their statistical analysis, it was stated that contents of bioactive compounds are a factor differentiating individual forms sorghum.

Our research shows that Sweet Carolina sorghum is a more rich source of bioactive compounds compared to Sweet Susana. It was a surprise that in the Sweet Carioline variety,

the level of carotenoids responsible for the yellow–brown color of the grain was much higher than in the brown Sweet Susane variety. Similar studies have been carried out by Choi et al. who, during their research, also found the presence of bioactive compounds in sorghum and divided them into five groups depending on the color of the grain [76]. Choi et al.

found that the color of the grain was associated with the content of TPC, but not with the content of anthocyanins. In addition, the antioxidant activity in the ABTS test was similar to the TPC content. On the basis of the tests, it was found that the amount of antioxidants differs from sorghum. The current literature on sorghum is quite extensive. There are many scientific reports on the cultivation of sorghum in the equatorial and tropical climate zone. In South America and in Africa, sorghum is widely used for food, feed, and energy. Our experiments prove that in conditions of a moderate climate, it is also possible to grow sorghum. Based on the results obtained, it was found that during 3 years of sorghum cultivation in Poland, the results are reproducible. Antioxidant activity and the presence of bioactive compounds are characteristic and stable for a given variety.

Concluding remarks

The study showed confirmed that the content of bioactive compounds in sorghum grain depends on the variety and fluctuate naturally as weather conditions change during the vegetation period. It was also investigated that white sorghum grain of *Sweet Caroline* is a richer source of phenolic acids and carotenoids. The color of the thin seed coat of caryopsis does not reflect the wealth of carotenoid compounds contained within. Moreover, the analysis of antioxidant activity applying an improved ABTS radical cation decolorization showed that for this variety, the capacity is also higher. Notwithstanding the foregoing, the content of bioactive compounds was confirmed in both tested sorghum varieties. Therefore, they may be a good source of grain considered as a functional food, since it combines nutritive and medicinal properties. On the basis of these results, the content of bioactive compounds was confirmed in sorghum grain cultivated in temperate climate.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Compliance with ethics requirement This article does not contain any studies with human or animal subjects.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Awika JM, Rooney LW (2004) Sorghum phytochemicals and their potential impact on human health. *Phytochem* 65:1199–1221
- Long SP, Marshall-Colon A, Zhu XG (2015) Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. *Cell* 161:56–66
- Che P, Anand A, Wu E, Sander JD, Simon MK, Zhu W, Sigmund AL, Zastrow-Hayes G, Miller M, Liu D, Lawit SJ, Zhao Z-Y, Albertsen MC, Jones TJ (2018) Developing a flexible, high-efficiency agrobacterium-mediated Sorghum transformation system with broad application. *Plant Biotechnol* 16:1388–1395
- Dicko MH, Gruppen H, Traoré AS, Voragen AG, Van Berkel WJ (2006) Sorghum grain as human food in Africa: relevance of content of starch and amylase activities. *Afr J Biotechnol* 5:384–395
- Dykes L, Rooney LW (2006) Sorghum and millet phenols and antioxidants. *J Cereal Sci* 44:236–251
- Proietti I, Frazzoli C, Mantovani A (2015) Exploiting nutritional value of staple foods in the world's semi-arid areas: risks, benefits, challenges and opportunities of sorghum. *Healthcare* 3:172–193
- Macedo WR, Araújo DK, Santos VM, Castro PRDC, Fernandes GM (2017) Plant growth regulators on sweet sorghum: physiological and nutritional value analysis. *Comunicata Scientiae* 8:170–174
- de Moraes Cardoso L, Pinheiro SS, Martino HSD, Pinheiro-Sant'Ana HM (2017) Sorghum (*Sorghum bicolor* L.): nutrients, bioactive compounds, and potential impact on human health. *Crit Rev Food Sci Nutr* 57:372–390
- Usman MA, Bolade MK, James S (2016) Functional properties of weaning food blends from selected sorghum (*Sorghum bicolor* (L.) Moench) varieties and soybean (*Glycine max*). *Afr J Food Sci* 10:112–121
- Kangama CO, Rumei X (2005) Introduction of sorghum (*Sorghum bicolor* L. Moench) into China. *Afr J Biotechnol* 4:575–579
- Vierling RA, Xiaang Z, Joshi CP, Gilbert ML, Nguyen HT (1994) Genetic diversity among elite Sorghum lines revealed by restriction fragment length polymorphisms and random amplified polymorphic DNAs. *Theor Appl Genet* 87:816–820
- Moreau RA, Harron AF, Powell MJ, Hoyt JL (2016) A comparison of the levels of oil, carotenoids, and lipolytic enzyme activities in modern lines and hybrids of grain sorghum. *J Am Oil Chem Soc* 93:569–573
- Pino F, Heinrichs AJ (2017) Sorghum forage in precision-fed dairy heifer diets. *J Dairy Sci* 100:224–235
- Getachew G, Putnam DH, De Ben CM, De Peters EJ (2016) Potential of sorghum as an alternative to corn forage. *Am J Plant Sci* 7:1106–1121
- Iqbal MA, Iqbal A (2015) Overviewing forage shortage for dairy animals and suitability of forage sorghum for ensiling. *Global Veterinaria* 14:173–177
- Vinutha KS, Kumar GA, Blümmel M, Srinivasa Rao P (2017) Evaluation of yield and forage quality in main and ratoon crops of different sorghum lines. *Trop Grassl-Forrajage Tropicales* 5:40–49
- Qi G, Li N, Sun XS, Wang D (2016) Overview of sorghum industrial utilization. In: Ciampitti I, Prasad V (eds) *Sorghum: state of the art and future perspectives*. ASA and CSSA, Madison. <https://doi.org/10.2134/agronmonogr58.2014.0070>
- Rooney W (2018). Achieving sustainable cultivation of sorghum. Volume 2: Sorghum utilization around the world. Achieving sustainable cultivation of sorghum. Volume 2: Sorghum utilization around the world, pp 243
- Pannacci E, Bartolini S (2016) Evaluation of sorghum hybrids for biomass production in central Italy. *Biomass Bioenerg* 88:135–141

20. Rooney W (2014) Sorghum. Cellulosic energy cropping systems 7:109–129
21. Barbanti L, Grandi S, Vecchi A, Venturi G (2006) Sweet and fibre sorghum (*Sorghum bicolor* (L.) Moench), energy crops in the frame of environmental protection from excessive nitrogen loads. *Eur J Agron* 25:30–39
22. Mullet J, Morishige D, McCormick R, Truong S, Hilley J, McKinley B, Anderson R, Olson SN, Rooney W (2014) Energy Sorghum—a genetic model for the design of C4 grass bioenergy crops. *J Exp Bot* 65:3479–3489
23. Szambelan K, Nowak J, Frankowski J, Szwengiel A, Jeleń H, Burczyk H (2018) The comprehensive analysis of sorghum cultivated in Poland for energy purposes: separate hydrolysis and fermentation and simultaneous saccharification and fermentation methods and their impact on bioethanol effectiveness and volatile by-products from the grain and the energy potential of sorghum straw. *Bioresour Technol* 250:750–757
24. Serra P, Colauzzi M, Amaducci S (2017) Biomass sorghum production risk assessment analysis: a case study on electricity production in the Po Valley. *Biomass Bioenerg* 96:75–86
25. Cole MR, Eggleston G, Petrie E, Uchimiya SM, Dalley C (2017) Cultivar and maturity effects on the quality attributes and ethanol potential of sweet sorghum. *Biomass Bioenerg* 96:183–192
26. Sebayang AH, Masjuki HH, Ong HC, Dharma S, Silitonga AS, Kusumo F, Milano J (2017) Optimization of bioethanol production from sorghum grains using artificial neural networks integrated with ant colony. *Ind Crops Prod* 97:146–155
27. Larnaudie V, Rochón E, Ferrari MD, Lareo C (2016) Energy evaluation of fuel bioethanol production from sweet sorghum using very high gravity (VHG) conditions. *Renew Energ* 88:280–287
28. Almodares A, Hadi MR (2009) Production of bioethanol from sweet sorghum: a review. *Afr J Agric Res* 9:772–780
29. Cao W, Sun C, Li X, Qiu J, Liu R (2017) Methane production enhancement from products of alkaline hydrogen peroxide pretreated sweet sorghum bagasse. *RSC Adv* 10:5701–5707
30. Zhang Z, Zhang G, Li W, Li C, Xu G (2016) Enhanced biogas production from sorghum stem by co-digestion with cow manure. *Int J Hydrogen Energ* 41:9153–9158
31. Zegada-Lizarazu W, Fernando Luna D, Monti A (2016) Differential characteristics of photochemical acclimation to cold in two contrasting sweet sorghum hybrids. *Physiol Plant* 157:479–489
32. Windpassinger S, Friedt W, Frauen M, Snowdon R, Wittkop B (2015) Designing adapted sorghum silage types with an enhanced energy density for biogas generation in temperate Europe. *Biomass Bioenerg* 81:496–504
33. Hermuth J, Kosova K (2017) Characterization of the first Czech sorghum variety Ruzrok tested in the Czech Republic. *Czech J Gen Plant Breed* 53:37–44
34. Przybylska-Balcerak A, Frankowski J, Stuper-Szablewska K (2018) Bioactive compounds in sorghum. *Eur Food Res Technol*. <https://doi.org/10.1007/s00217-018-3207-0>
35. Stefoska-Needham A, Beck EJ, Johnson SK, Tapsell LC (2015) Sorghum: an underutilized cereal whole grain with the potential to assist in the prevention of chronic disease. *Food Rev Int* 31:401–437
36. Marston K, Khouryieh H, Aramouni F (2016) Effect of heat treatment of sorghum flour on the functional properties of gluten-free bread and cake. *Food Sci Technol* 65:637–644
37. Althwab S, Carr TP, Weller CL, Dweikat IM, Schlegel V (2015) Advances in grain sorghum and its co-products as a human health promoting dietary system. *Food Res Int* 77:349–359
38. Trappey EF, Khouryieh H, Aramouni F, Herald T (2015) Effect of sorghum flour composition and particle size on quality properties of gluten-free bread. *Food Sci Technol Int* 21:188–202
39. Bin T (2007) Properties of grain sorghum and its prospects of utilization in food industry. *Cereal Feed Ind* 7–9
40. Frankowski J (2017) Nutritional and therapeutic properties of sorghum (*Sorghum* Moench). *Post Fitoter* 18:209–214
41. Wu G, Bennett SJ, Bornman JF, Clarke MW, Fang Z, Johnson SK (2017) Phenolic profile and content of sorghum grains under different irrigation managements. *Food Res Int* 97:347–355
42. Costa RF, Pires DADA, Moura MMA, Sales ECJD, Rodrigues JAS, Rigueira JPS (2016) Agronomic characteristics of sorghum genotypes and nutritional values of silage. *Acta Sci Anim Sci* 38:127–133
43. Robbins RJ (2003) Phenolic acids in foods: an overview of analytical methodology. *J Agric Food Chem* 51:2866–2887
44. Stalikas CD (2007) Extraction, separation, and detection methods for phenolic acids and flavonoids. *J Sep Sci* 30:3268–3295
45. Croft KD (1998) The chemistry and biological effects of flavonoids and phenolic acids. *Ann NY Acad Sci* 854:435–442
46. Stuper-Szablewska K, Kurasiak-Popowska D, Nawracała J, Perkowski J (2014) Comparison of phenolic acid contents in various wheat genotypes. *Przem Chem* 93:2274–2278
47. Ghasemzadeh A, Ghasemzadeh N (2011) Flavonoids and phenolic acids: role and biochemical activity in plants and human. *J Med Plants Res* 31:6697–6703
48. Cai Y, Luo Q, Sun M, Corke H (2004) Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci* 74:2157–2184
49. Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M (1999) Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem* 47:3954–3962
50. Natella F, Nardini M, Di Felice M, Scaccini C (1999) Benzoic and cinnamic acid derivatives as antioxidants: structure-activity relation. *J Agric Food Chem* 47:1453–1459
51. Murphy JB, Stutte CA (1978) Analysis for substituted benzoic and cinnamic acids using high-pressure liquid chromatography. *Anal Biochem* 86:220–228
52. Taleon V, Dykes L, Rooney WL, Rooney LW (2012) Effect of genotype and environment on flavonoid concentration and profile of black sorghum grains. *J Cereal Sci* 56:470–475
53. Dykes L, Peterson GC, Rooney WL, Rooney LW (2011) Flavonoid composition of lemon-yellow sorghum genotypes. *Food Chem* 128:173–179
54. Svensson L, Sekwati-Monang B, Lutz DL, Schieber A, Ganzle MG (2010) Phenolic acids and flavonoids in nonfermented and fermented red sorghum (*Sorghum bicolor* (L.) Moench). *J Agric Food Chem* 58:9214–9220
55. Dykes L, Seitz LM, Rooney WL, Rooney LW (2009) Flavonoid composition of red sorghum genotypes. *Food Chem* 116:313–317
56. Moreau RA, Whitaker BD, Hicks KB (2002) Phytosterols, phytostanols, and their conjugates in foods: structural diversity, quantitative analysis, and health-promoting uses. *Prog Lipid Res* 41:457–500
57. Lagarda MJ, Garcia-Llatas G, Farre R (2006) Analysis of phytosterols in foods. *J Pharm Biom Anal* 41(5):1486–1496
58. Kean EG, Bordenave N, Ejeta G, Hamaker BR, Ferruzzi MG (2011) Carotenoid bioaccessibility from whole grain and decorticated yellow endosperm sorghum porridge. *J Cereal Sci* 54:450–459
59. Farre G, Bai C, Twyman RM, Capell T, Christou P, Zhu C (2011) Nutritious crops producing multiple carotenoids—a metabolic balancing act. *Trends Plant Sci* 16:532–540
60. Kean EG, Ejeta G, Hamaker BR, Ferruzzi MG (2007) Characterization of carotenoid pigments in mature and developing kernels of selected yellow-endosperm sorghum varieties. *J Agric Food Chem* 55:2619–2626

61. Blessin CW (1962) Carotenoids of corn and sorghum. I. Analytical procedure. *Cereal Chem* 39:236–242
62. Gill SS, Tuteja N (2010) Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Biochem* 48:909–930
63. Wu G, Johnson SK, Bornman JF, Bennett SJ, Fang Z (2017) Changes in whole grain polyphenols and antioxidant activity of six sorghum genotypes under different irrigation treatments. *Food Chem* 214:199–207
64. Tobias AV, Arnold FH (2006) Biosynthesis of novel carotenoid families based on unnatural carbon backbones: a model for diversification of natural product pathways. *BBA* 1761:235–246
65. Costa PHAD, Neto ADDA, Bezerra MA, Prisco JT, Gomes-Filho E (2005) Antioxidant-enzymatic system of two sorghum genotypes differing in salt tolerance. *Braz J Plant Physiol* 17:353–362
66. Przybylska A, Stuper-Szablewska K, Matysiak A, Perkowski J (2018) Optimization of extraction conditions total phenolic compounds and wheat grain antioxidant activity. *ABID* 1(2018):5–12
67. Mróz P, Wilczek K, Żak M, Zielińska-Pisklak M (2012) Chromatographic methods for isolation and identification of phenolic acids. *Biul Wydz Farm WUM* 6:40–48
68. Kurasiak-Popowska D, Ryńska B, Stuper-Szablewska K (2019) Analysis of distribution of selected bioactive compounds in *Camelina sativa* from seeds to pomace and oil. *Agronomy* 9(4):168
69. Góral T, Stuper-Szablewska K, Przybylska A, Matysiak A, Perkowski J (2017) Impact of inoculation with *Fusarium* fungi on beta-sitosterol content in grain of different winter wheat cultivars. *Biul IHAR* 281:27–35
70. Przybylska-Balcerek A, Kurasiak-Popowska D, Stuper-Szablewska K (2019) Blue Corner—a new source of antioxidants. *KOSMOS Problemy Nauk Biologicznych* 68:161–166
71. Xiong Y, Zhang P, Luo J, Johnson S, Fang Z (2019) Effect of processing on the phenolic contents, antioxidant activity and volatile compounds of sorghum grain tea. *J Cereal Sci* 85:6–14
72. Przybylska-Balcerek A, Góral T, Kurasiak-Popowska D, Stuper-Szablewska K (2019) Phenolic acids in various genotypes of cereals grown in Poland in the years 2017–2018. *Acad J Med Plants* 7:092–099
73. John RN, Anyango JO (2011) Sorghum flour and flour products: production, nutritional quality, and fortification. Academic Press, pp 127–139
74. Dykes L, Rooney LW (2006) Sorghum and millet phenols and antioxidants. *J Cereal Sci* 44:236–251
75. Stefoska-Needham A, Beck E, Johnson SK, Tapsell LC (2015) Sorghum: an underutilized cereal whole grain with the potential to assist in the prevention of chronic disease. *Food Rev Int* 31(4):401–437
76. Choi SC, Kim JM, Lee YG, Kim C (2019) Antioxidant activity and contents of total phenolic compounds and anthocyanins according to grain colour in several varieties of Sorghum bicolor (L.) Moench. *Cereal Res Commun* 47(2):228–238

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.