

Erratum

Generalized Kneser coloring theorems with combinatorial proofs

Günter M. Ziegler*

TU Berlin, Inst. Mathematics, MA 6-2, 10623 Berlin, Germany
(e-mail: ziegler@math.tu-berlin.de)

Invent. math., <http://dx.doi.org/10.1007/s002220100188>

Oblatum 31-V-2005 & 6-VI-2005
Published online: 27 October 2005 – © Springer-Verlag 2005

In [5], we presented a lower bound for the chromatic numbers of hypergraphs $\text{KG}_s^r \mathcal{S}$, “generalized r -uniform Kneser hypergraphs with intersection multiplicities s .“ It generalized previous lower bounds by Kříž [1, 2] for the case $s = (1, \dots, 1)$ without intersection multiplicities, and by Sarkaria [4] for $\mathcal{S} = \binom{[n]}{k}$.

The following two problems that arise for intersection multiplicities $s_i > 1$ were noticed by Carsten Lange resp. by Karsten Vogel:

1. In the presence of intersection multiplicities, there are two different versions of a “Kneser hypergraph,” depending on whether one admits hypergraph edges that are multisets rather than sets. It is shown in [3] that the chromatic numbers are substantially different for the two concepts of hypergraphs. The lower bounds of Sarkaria [4] and of [5, Thm. 5.1] apply only to the multiset version. The coloring and upper bound of [5, Lemma 3.1] is also valid for the multiset version. The coloring of [5, Example 7.2] refers to the set version.
2. The reductions to the case of prime r in the proof for [5, Thm. 5.1] works only if the intersection multiplicities are strictly smaller than the largest prime factor of r . (Specifically, a problem arises in the reduction of [5, pp. 679–680] if the auxiliary set system \mathcal{T} is empty.) Currently we have no valid proof for the lower bound result in the other cases. This also applies to the special case $\mathcal{S} = \binom{[n]}{k}$ of Sarkaria [4].

Details will be presented in [3].

* Partially supported by Deutsche Forschungs-Gemeinschaft (DFG)

References

1. Kříž, I.: Equivariant cohomology and lower bounds for chromatic numbers. *Trans. Am. Math. Soc.* **33**, 567–577 (1992)
2. Kříž, I.: A correction to “Equivariant cohomology and lower bounds for chromatic numbers”. *Trans. Am. Math. Soc.* **352**, 1951–1952 (2000)
3. Lange, C., Ziegler, G.M.: On generalized Kneser hypergraph colorings. Preprint, March 2005, 8 pages; www.arXiv.org/math.CO/0504607
4. Sarkaria, K.S.: A generalized Kneser conjecture. *J. Combin. Theory, Ser. B* **49**, 236–240 (1990)
5. Ziegler, G.M.: Generalized Kneser coloring theorems with combinatorial proofs. *Invent. Math.* **147**, 671–691 (2002)