



## Correction to: Effects of intrinsic and extrinsic factors on reproduction of an ecologically significant invasive bryozoan: implications for invasion success

Danielle Denley<sup>1</sup> · Anna Metaxas<sup>1</sup>

Published online: 2 December 2017  
© Springer-Verlag GmbH Germany, part of Springer Nature 2017

**Correction to: Mar Biol (2017) 164:145**  
<https://doi.org/10.1007/s00227-017-3172-3>

Fig. 2 was published with incorrect values for June 2012 and May 2014. The corrected Figure 2 is provided here.


---

The online version of the original article can be found under  
<https://doi.org/10.1007/s00227-017-3172-3>.

---

✉ Danielle Denley  
danielle.denley@dal.ca

<sup>1</sup> Department of Oceanography, Dalhousie University,  
Halifax, NS B3H 4R2, Canada



**Fig. 2** Potential fecundity (mean + SE,  $n = 1\text{--}211$ ) of colonies of *Membranipora membranacea* collected on the 4 most numerically abundant algal substrates in Nova Scotia (SL, *Saccharina latissima*; LD, *Laminaria digitata*; AC, *Agarum clathratum*; Fu, *Fucus evanescens*) at The Lodge, Sandy Cove, and Paddy's Head from June 2012 to August 2013; and at The Lodge from July 2014 to November 2015

and Sandy Cove from May 2014 to November 2015. Zeros indicate colonies were not fecund (fecundity = 0). NA indicates colony fecundity could not be accurately measured (June 2012, see "Materials and methods") or colonies were not present on the algal substrate (May 2014)