Correction to: Multiple zooplankton species alter the stoichiometric interactions between producer and consumer levels

Christoph Plum¹ · Helmut Hillebrand^{1,2,3}

Published online: 1 February 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Marine Biology (2019) 166:163
<https://doi.org/10.1007/s00227-019-3609-y>

Unfortunately, in the print version of this article the below appendix Figs. 5, 6 and 7 and appendix Table 4 were wrongly placed in the body of the article.

Appendix

See Figs. 5, 6 and 7 and Table 4.

The original article can be found online at <https://doi.org/10.1007/s00227-019-3609-y>.

✉ Christoph Plum
c.plum@uni-oldenburg.de

¹ University of Oldenburg, Institute for Chemistry and Biology of the Marine Environment (ICBM), Schleusenstr. 1, 26382 Wilhelmshaven, Germany

² Helmholtz Institute for Functional Marine Biodiversity (HIFMB) at the University of Oldenburg, Oldenburg, Germany

³ Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research [AWI], Bremerhaven, Germany

Fig. 5 Algal biomass (algal C $\mu\text{mol L}^{-1}$), RUE and NUE in response to the factorial combination of light and nitrogen availability (mean $\pm$ SE) before grazing (left panel) and after grazing (right panel). HL, high light; LL, low light; N+, nitrogen enriched; N-, nitrogen limited

Fig. 6 Algal elemental ratios (molar C:N, C:P and N:P) in response to the factorial combination of light and nitrogen availability (mean $\pm$ SE) before grazing (left panel) and after grazing (right panel). For abbreviations, see “Fig. 5 in Appendix”

Fig. 7 *Skeletonema costatum* (Ske) was proportionally the most abundant species in the algal assemblages (except for HL +N) with up to 70% of the total biovolume. Both, *Rhodomonas salina* (Rho) and *Tetraselmis* sp. (Tet) remained below 20% of total biovolume. The proportions were more equally distributed at HL and +N which

was mainly based on the dominance of *Tetraselmis* sp. (up to 50% of total biovolume) and higher proportions of the subdominant species *Rhodomonas salina* (up to 25% of total biovolume). The biovolume of *Chaetoceros danicus* (Chae) contributed less than 1% to the total algal biovolume under all resource conditions

Table 4 Phytoplankton response to light, nitrogen and the interaction of both before grazing tested with a two-factorial ANOVA

Response	Factor	df	F	p
Biomass	Light	1	497.67	< 0.001
	Nitrogen	1	430.4	< 0.001
	Light × nitrogen	1	434.85	< 0.001
RUE	Light	1	3.1108	0.087
	Nitrogen	1	67.7372	< 0.001
	Light × nitrogen	1	69.6467	< 0.001
NUE	Light	1	3.6414	0.066
	Nitrogen	1	47.0745	< 0.001
	Light × nitrogen	1	74.4110	< 0.001
C:N	Light	1	114.78	< 0.001
	Nitrogen	1	116.48	< 0.001
	Light × nitrogen	1	50.13	< 0.001
C:P	Light	1	118.42	< 0.001
	Nitrogen	1	114.44	< 0.001
	Light × nitrogen	1	184.67	< 0.001
N:P	Light	1	113.23	< 0.001
	Nitrogen	1	212.84	< 0.001
	Light × nitrogen	1	276.83	< 0.001

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.