



Correction to: Prey size spectra and predator to prey size ratios of southern ocean salps

Christian K. Fender¹  · Moira Décima² · Andres Gutierrez-Rodriguez³ · Karen E. Selph⁴ · Natalia Yingling¹ · Michael R. Stukel^{1,5}

Published online: 20 May 2023

This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2023

Correction to: Marine Biology (2023) 170:40
<https://doi.org/10.1007/s00227-023-04187-3>

In this article the Fig. 3 has been corrected. The correct figure and values should have appeared as shown below.

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00227-023-04187-3>.

✉ Christian K. Fender
ckf18b@fsu.edu

¹ Department of Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee, FL, USA

² Integrative Oceanography Division, Scripps Institution of Oceanography, La Jolla, CA, USA

³ National Institute of Water and Atmospheric Research, Wellington, New Zealand

⁴ Department of Oceanography, University of Hawaii, Honolulu, HI, USA

⁵ Center for Ocean-Atmospheric Prediction Studies, Florida State University, Tallahassee, FL, USA




Fig. 3 Contribution of particles of various size classes to the total biomass present in the water column (outlined by black boxes) compared to that of the guts of each of the 58 salps sampled averaged by species (as indicated on the x-axis) per cycle. WC water column, SF

Salpa fusiformis, ST *Salpa thompsoni*, TV *Thetys vagina*, SZ *Soestia zonaria*, PC *Pegea confoederata*, IM *Ihlea magalhanica*, TD *Thalia democratica*. Horizontal dashed lines extend the WC demarcations for easier comparison to salp species