

D. N. Bateman · J. Chick · A. M. Good · C. A. Kelly  
G. Masterton

## Are selective serotonin re-uptake inhibitors associated with an increased risk of self-harm by antidepressant overdose?

Published online: 3 February 2005  
© Springer-Verlag 2005

---

### Eur J Clin Pharmacol (2004) 60:221–224

It has come to our attention that an error was made in one calculation in respect of the above paper. As a result the likelihood ratio published is incorrect. The correct likelihood ratios for admissions involving venlafaxine

shown in Table 1 (and also in the abstract) is 0.97 (0.81–1.16), not 1.33 (1.13–1.55).

This error does not alter the main findings of the manuscript or any of the overall conclusions that we reached.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00228-004-0748-x>

---

D. N. Bateman (✉) · A. M. Good · C. A. Kelly  
Scottish Poisons Information Bureau (NPIS Edinburgh),  
Royal Infirmary of Edinburgh, 51 Little France Crescent,  
EH10 4SA Edinburgh, Scotland  
E-mail: [spib@luht.scot.nhs.uk](mailto:spib@luht.scot.nhs.uk)  
Tel.: +131-242-1383  
Fax: +131-242-1387

G. Masterton  
Department of Psychological Medicine,  
Royal Infirmary of Edinburgh, EH16 4SA  
Edinburgh, Scotland

J. Chick  
Royal Edinburgh Hospital,  
EH10 5HF Edinburgh, Scotland