Correction to: Protective effect of flumazenil infusion in severe acute benzodiazepine toxicity: a pilot randomized trial

Akram Sadat Razavizadeh¹ · Nasim Zamani^{2,3} · Pardis Ziaeefer⁴ · Sara Ebrahimi⁴ · Hossein Hassanian-Moghaddam^{2,3}

Published online: 26 February 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: *European Journal of Clinical Pharmacology*
<https://doi.org/10.1007/s00228-020-03031-7>

There were two errors in the abstract and fig. 1. The first sentence of abstract should appear as follows: Sixty severely poisoned patients (need for intubation due to loss of consciousness) exposed by sole benzodiazepines referred to Loghman Hakim hospital between May 2018 and August 2019 were considered to be included in the current study. Some numbers noted in fig. 1 should be corrected in the flow-chart and be appeared as below:

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00228-020-03031-7>

✉ Hossein Hassanian-Moghaddam
hassanian@sbmu.ac.ir

¹ Imam Ali Hospital, Alborz University of Medical Sciences, Karaj, Iran

² Social Determinants of Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³ Department of Clinical Toxicology, Loghman Hakim Hospital, School of Medicine, Shahid Beheshti University of Medical Sciences, South Karegar Street, Tehran, Iran

⁴ School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Fig. 1 Flowchart of recruiting patients and their respiratory impairments in flumazenil vs. conservative treatment. 0 h in case group defines the initiation time of flumazenil infusion after receiving flumazenil loading dose. 0 h in control group defines the time after receiving flumazenil loading dose