Wolfgang Ebeling · Atsushi Takahashi

Graded matrix factorizations of size two and reduction

Received: 15 July 2021 / Accepted: 7 January 2022 / Published online: 29 January 2022

Abstract. We associate a complete intersection singularity to a graded matrix factorization of size two of a polynomial in three variables. We show that we get an inverse to the reduction of singularities considered by C. T. C. Wall. We study this for the full strongly exceptional collections in the triangulated category of graded matrix factorizations constructed by H. Kajiura, K. Saito, and the second author.

0. Introduction

In his classification of singularities [16], C. T. C. Wall noticed that there is a relation between series of singularities given by a reduction. More precisely, he considers a linear reduction L of a complete intersection singularity in $\mathbb{C}^4$ to a hypersurface singularity in $\mathbb{C}^3$ and a reduction Δ of a hypersurface singularity in $\mathbb{C}^3$ of corank 3 to one of corank 2. The aim of this paper is to show that as an inverse of such a reduction one can consider a graded matrix factorization of size two.

We consider weighted homogeneous deformations of an invertible polynomial. We use the approach of [1] to define graded matrix factorizations for such deformations. We classify weighted homogeneous deformations of invertible polynomials of three variables and we define Dolgachev numbers for them. We consider a graded matrix factorization of size two of such a deformation and associate a complete intersection singularity to it. We show that we get the original singularity back by L - or Δ -reduction. Moreover we consider graded matrix factorizations of size two of invertible polynomials in two variables and show that they are related to a reduction between hypersurface singularities in $\mathbb{C}^2$ considered in [10].

In [12], H. Kajiura, K. Saito, and the second author constructed a full strongly exceptional collection in the triangulated category of graded matrix factorizations of a polynomial associated to a regular system of weights whose smallest exponents are equal to -1 . These polynomials define the 14 exceptional unimodal hypersurface

W. Ebeling (✉): Institut für Algebraische Geometrie, Leibniz Universität Hannover, Postfach 6009, D-30060 Hannover, Germany
e-mail: ebeling@math.uni-hannover.de

A. Takahashi: Department of Mathematics, Graduate School of Science, Osaka University, Toyonaka Osaka 560-0043, Japan
e-mail: takahashi@math.sci.osaka-u.ac.jp

Mathematics Subject Classification: 32S05 · 13C14 · 18G80

singularities, the 6 heads of the bimodal series of hypersurface singularities, and 2 further singularities. They are weighted homogeneous deformations of invertible polynomials of three variables. The exceptional collection is described by a certain quiver. The graded matrix factorizations of size two listed in the paper correspond to the outermost vertices of certain arms of the quiver. We show that they correspond to singularities where the corresponding length of the arm (which is one of the Dolgachev numbers) is increased by one. We thus obtain the picture of Fig. 2.

There is Arnold's strange duality between the 14 exceptional unimodal hypersurface singularities and the extension [10] of it to the bimodal series and the complete intersection singularities which reduce to the 14 exceptional unimodal singularities and the 6 heads of the bimodal series. This can be considered as a special kind of mirror symmetry. As a corollary of our main result we get that the reduction is mirror dual to a special adjacency between these singularities.

1. Graded matrix factorizations for invertible polynomials

A polynomial $f(x_1, \dots, x_n) \in \mathbb{C}[x_1, \dots, x_n]$ is called *weighted homogeneous*, if there are positive integers $w_1, \dots, w_n$ and d such that $f(\lambda^{w_1}x_1, \dots, \lambda^{w_n}x_n) = \lambda^d f(x_1, \dots, x_n)$ for $\lambda \in \mathbb{C}^*$. We call $(w_1, \dots, w_n; d)$ a *system of weights*. If $\gcd(w_1, \dots, w_n, d) = 1$, then the system of weights is called *reduced*.

Definition 1. A weighted homogeneous polynomial $f(x_1, \dots, x_n)$ is called *invertible* if the following conditions are satisfied

- (i) it can be written

$$f(x_1, \dots, x_n) = \sum_{i=1}^n a_i \prod_{j=1}^n x_j^{E_{ij}},$$

where $a_i \in \mathbb{C}^*$, E_{ij} are non-negative integers, and the $n \times n$ -matrix $E := (E_{ij})$ is invertible over $\mathbb{Q}$,

- (ii) it has an isolated singularity at the origin.

Let $f(x_1, \dots, x_n) = \sum_{i=1}^n a_i \prod_{j=1}^n x_j^{E_{ij}}$ be an invertible polynomial. Without loss of generality, we assume that $\det E > 0$. The *canonical system of weights* is the system of weights $(w_1, \dots, w_n; d)$ given by the unique solution of the equation

$$E \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix} = \det(E) \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \quad d := \det(E).$$

The canonical system of weights $(w_1, \dots, w_n; d)$ is in general non-reduced, define

$$c_f := \gcd(w_1, \dots, w_n, d).$$

Definition 2. A *weighted homogeneous deformation of an invertible polynomial* is a polynomial of the form

$$f(x_1, \dots, x_n) = \sum_{i=1}^N a_i \prod_{j=1}^n x_j^{E_{ij}},$$

where $N \geq n$, $a_i \in \mathbb{C}^*$, E_{ij} are non-negative integers, $\sum_{i=1}^N a_i \prod_{j=1}^n x_j^{E_{ij}}$ is an invertible polynomial with canonical weight system $(w_1, \dots, w_n; d)$, and $\prod_{j=1}^n x_j^{E_{ij}}$ are monomials of the same degree d for $1 \leq i \leq N$.

Note that the case $N = n$ is permitted, i.e., the weighted homogeneous deformations of an invertible polynomial include the invertible polynomial itself.

Definition 3. Let $f(x_1, \dots, x_n) = \sum_{i=1}^N a_i \prod_{j=1}^n x_j^{E_{ij}}$ be a weighted homogeneous deformation of an invertible polynomial. Consider the free abelian group $\oplus_{i=1}^n \mathbb{Z}\vec{x}_i \oplus \mathbb{Z}\vec{f}$ generated by the symbols $\vec{x}_i$ for the variables x_i for $i = 1, \dots, n$ and the symbol $\vec{f}$ for the polynomial f . The *maximal grading* L_f of the invertible polynomial f is the abelian group defined by the quotient

$$L_f := \bigoplus_{i=1}^n \mathbb{Z}\vec{x}_i \oplus \mathbb{Z}\vec{f} / I_f,$$

where I_f is the subgroup generated by the elements

$$\vec{f} - \sum_{j=1}^n E_{ij} \vec{x}_j, \quad i = 1, \dots, N.$$

Note that L_f is an abelian group of rank 1 which is not necessarily free. We have the degree map $\deg : L_f \rightarrow \mathbb{Z}$ defined by $\vec{x}_i \mapsto w_i/c_f$, $\vec{f} \mapsto d/c_f$, where $(w_1, \dots, w_n; d)$ is the canonical system of weights of f . This map is an isomorphism if and only if the canonical system of weights is reduced, see [8, Proposition 17].

Let $f(x_1, \dots, x_n)$ be a weighted homogeneous deformation of an invertible polynomial. We recall the definition of an L_f -graded matrix factorization of f , see [1] and the references therein.

Let $S = \mathbb{C}[x_1, \dots, x_n]$. This is naturally an L_f -graded algebra:

$$S = \bigoplus_{\vec{l} \in L_f} S_{\vec{l}}.$$

For any L_f -graded S -module M , let $M(\vec{l})$ denote the L_f -graded S -module, where the grading is shifted by $\vec{l} \in L_f$:

$$M(\vec{l}) = \bigoplus_{\vec{l}' \in L_f} M(\vec{l})_{\vec{l}'}, \quad M(\vec{l})_{\vec{l}'} := M_{\vec{l}+\vec{l}'},$$

The grading shift by $\vec{l} \in L_f$ induces a functor $(\vec{l})$ on the category of finitely generated L_f -graded S -modules.

Definition 4. An L_f -graded matrix factorization of f is

$$\overline{F} := \left(F_0 \begin{array}{c} \xrightarrow{f_0} \\ \xleftarrow{f_1} \end{array} F_1 \right),$$

where F_0 and F_1 are L_f -graded free S -modules of finite rank and $f_0 : F_0 \rightarrow F_1$, $f_1 : F_1 \rightarrow F_0$ are morphisms such that $f_1 \circ f_0 = f \cdot \text{id}_{F_0}$ and $f_0(f_1) \circ f_1 = f \cdot \text{id}_{F_1}$. The rank of F_1 is equal to the rank of F_0 and it is called the *size* or the *rank* of the matrix factorization $\overline{F}$.

Let $f(x_1, \dots, x_n)$ be a weighted homogeneous deformation of an invertible polynomial with reduced weight system $(w_1, \dots, w_n; d)$. We define

$$\varepsilon_f := w_1 + \dots + w_n - d.$$

Let $f(x, y, z)$ be a weighted homogeneous deformation of an invertible polynomial in three variables. We shall associate Dolgachev numbers to f . This is done as follows. Let $R_f := \mathbb{C}[x, y, z]/(f)$. Let $r \geq 3$, $A = (\alpha_1, \dots, \alpha_r)$ and let $\lambda = (\lambda_3, \dots, \lambda_r)$ be an $(r-2)$ -tuple of complex numbers with $\lambda_3 = 1, \lambda_i \neq \lambda_j$ for $i \neq j, i, j = 3, \dots, r$. Let $R_{A, \lambda}$ be the ring associated to the weighted projective line corresponding to the tuple of numbers $A = (\alpha_1, \dots, \alpha_r)$ [11], namely the factor algebra

$$R_{A, \lambda} := \mathbb{C}[X_1, \dots, X_r]/I_\lambda$$

where I_λ is the ideal generated by the polynomials

$$X_i^{\alpha_i} - X_2^{\alpha_2} + \lambda_i X_1^{\alpha_1}, \quad i = 3, \dots, r.$$

We shall consider a classification of weighted homogeneous deformations of invertible polynomials according to the classification of invertible polynomials in three variables in [8]. In each type except V, two of the three monomials are replaced by a product of m terms involving parameters $(\lambda_3, \dots, \lambda_{m+2})$. In the case of type V, we set $m = 1$.

Definition 5. The *Dolgachev numbers* of a weighted homogeneous deformation f of an invertible polynomial is the $(m+2)$ -tuple ($m \geq 1$) of numbers $A = (\alpha_1, \alpha_2, \underbrace{\alpha_3, \dots, \alpha_3}_m)$ defined by the embedding $R_f \hookrightarrow R_{A, \lambda}$ given by Table 1.

It follows from the embedding $R_f \hookrightarrow R_{A, \lambda}$ into the ring of a weighted projective line that $L_f \cong \mathbb{Z}$.

2. Graded matrix factorizations of size two

We shall now consider L_f -graded matrix factorizations of size two for the following two cases

- (A) f is a weighted homogeneous deformation of an invertible polynomial in three variables,

Table 1. The image of the generators in $R_{A,\lambda}$

Type	$f(x, y, z)$	x	y	z
I	$-\prod_{i=3}^{m+2} (y \frac{p_2}{m} - \lambda_i x \frac{p_1}{m}) + z^{p_3}$	X_1	X_2	$X_3 \cdots X_{m+2}$
II ₁	$-x \prod_{i=3}^{m+2} (y \frac{p_2}{m} - \lambda_i x \frac{p_1-1}{m}) + z^{p_3}$	$X_1^{p_3}$	X_2	$X_1 X_3 \cdots X_{m+2}$
II ₂	$-\prod_{i=3}^{m+2} (z \frac{p_3}{m} - \lambda_i x \frac{p_1}{m}) + xy^{p_2}$	$X_1^{p_3}$	$X_3 \cdots X_{m+2}$	$X_1 X_2$
III	$-xy \prod_{i=3}^{m+2} (y \frac{p_2-1}{m} - \lambda_i x \frac{p_1-1}{m}) + z^{p_3}$	$X_1^{p_3}$	$X_2^{p_3}$	$X_1 X_2 X_3 \cdots X_{m+2}$
IV	$-x \prod_{i=3}^{m+2} (y \frac{p_2}{m} - \lambda_i x \frac{p_1-1}{m}) + yz^{p_3}$	$X_1^{p_3} X_2$	$X_2^{p_1}$	$X_1 X_3 \cdots X_{m+2}$
V	$x^{q_1} y + y^{q_2} z - z^{q_3} x$	$X_2 X_3^{q_2}$	$X_3 X_1^{q_3}$	$X_1 X_2^{q_3}$
Type	α_1	α_2	α_3	
I	$\frac{p_1}{m}$	$\frac{p_2}{m}$	p_3	
II ₁	$\frac{p_3}{m}(p_1 - 1)$	$\frac{p_2}{m}$	p_3	
II ₂	$\frac{p_3}{m}(p_1 - 1)$	$\frac{p_3}{m}$	p_2	
III	$\frac{p_3}{m}(p_1 - 1)$	$\frac{p_3}{m}(p_2 - 1)$	p_3	
IV	$\frac{p_3}{m}(p_1 - 1)$	$\frac{p_1 p_2 - p_1 + 1}{m}$	p_3	
V	$q_2 q_3 - q_3 + 1$	$q_3 q_1 - q_1 + 1$	$q_1 q_2 - q_2 + 1$	

(B) f is an invertible polynomial in two variables.

(A) Let $S = \mathbb{C}[x, y, z]$ and let $f(x, y, z)$ be a weighted homogeneous deformation of an invertible polynomial of the form

$$f(x, y, z) = p_1(x, y, z)h_1(x, y, z) + p_2(x, y, z)h_2(x, y, z),$$

where $p_i \in S_{\vec{p}_i}$ for some $\vec{p}_i \in L_f, i = 1, 2$, and p_1, p_2 forms a regular sequence in S . As in [5, Section 2] (cf. [1, Definition 2.11]), we consider the Koszul resolution of the L_f -graded S -module $M = S/(p_1, p_2)$

$$0 \longrightarrow S(-\vec{p}_1 - \vec{p}_2) \longrightarrow S(-\vec{p}_1) \oplus S(-\vec{p}_2) \longrightarrow S \longrightarrow S/(p_1, p_2) \longrightarrow 0.$$

This yields the L_f -graded matrix factorization $Q := \left(F_0 \xrightleftharpoons[f_1]{f_0} F_1 \right)$ of f such that

$$F_0 = S(\vec{0}) \oplus S(\vec{f} - \vec{p}_1 - \vec{p}_2), \quad F_1 := S(\vec{f} - \vec{p}_1) \oplus S(\vec{f} - \vec{p}_2),$$

and f_0 and f_1 are given by the matrices

$$q_0 = \begin{pmatrix} h_1(x, y, z) & -p_2(x, y, z) \\ h_2(x, y, z) & p_1(x, y, z) \end{pmatrix} \text{ and } q_1 = \begin{pmatrix} p_1(x, y, z) & p_2(x, y, z) \\ -h_2(x, y, z) & h_1(x, y, z) \end{pmatrix}$$

respectively.

We associate to the matrix factorization a complete intersection singularity in $\mathbb{C}^4$. Denote the coordinates of $\mathbb{C}^4$ by w, x, y, z .

Definition 6. The complete intersection singularity $(X_Q, 0)$ is the singularity defined by

$$\begin{aligned} \mathbf{F}_Q(w, x, y, z) &= (F_{Q,1}(w, x, y, z), F_{Q,2}(w, x, y, z)) \\ &:= (-h_1(x, y, z) + wp_2(x, y, z), h_2(x, y, z) + wp_1(x, y, z)). \end{aligned}$$

We can get the function f back from $\mathbf{F}_Q$ by the linear reduction considered, in the case $p_1(x, y, z) = y$ and $p_2(x, y, z) = z$, by Wall [16, 7.9]. Namely, let $L_w \mathbf{F}_Q$ be the elimination of the variable w from the functions $F_{Q,1}, F_{Q,2}$. This is given by the resultant $R_w(F_{Q,1}, F_{Q,2})$ of the functions with respect to the variable w . We have

$$L_w \mathbf{F}_Q = R_w(F_{Q,1}, F_{Q,2}) = \det q_1 = f.$$

Now suppose that $p_1(x, y, z) = y$, $p_2(x, y, z) = z$, $h_2(x, y, z) = z$ and $h_1(x, y, z) = 2a(x, y)z + b(x, y)$. Then

$$(F_{Q,1}(w, x, y, z), F_{Q,2}(w, x, y, z)) = (-h_1(x, y, z) + wz, z + wy)$$

and the substitution $z = -wy$ in $F_{Q,1}$ gives

$$\begin{aligned} F_Q(w, x, y) &:= F_{Q,1}(w, x, y, -wy) = -w^2y - h_1(x, y, -wy) \\ &= -w^2y + 2wya(x, y) - b(x, y). \end{aligned}$$

Thus $(X_Q, 0)$ is in this case a hypersurface singularity defined by $F_Q = 0$. The discriminant of the polynomial $F_Q(w, x, y)$ in the variable w is

$$\Delta_w F_Q(w, x, y) = y^2 a(x, y)^2 - yb(x, y).$$

This is the reduction $\Delta_w F_Q$ of the polynomial F_Q considered by Wall in [16, 7.3]. The polynomial f is equal to

$$f(x, y, z) = z^2 + 2ya(x, y)z + yb(x, y).$$

Under the substitution $\tilde{z} := z + ya(x, y)$ this becomes

$$f(x, y, \tilde{z}) = \tilde{z}^2 - y^2 a(x, y)^2 + yb(x, y) = \tilde{z}^2 - \Delta_w F_Q(w, x, y).$$

(B) Now let $S = \mathbb{C}[x, y]$ and let $f(x, y)$ be an invertible polynomial of the form $f(x, y) = x^\alpha y^\beta + y^3$, where $\alpha \geq 5$. We consider the Koszul resolution of the L_f -graded S -module $M = S/(x^2, y^2)$

$$0 \longrightarrow S(-2\vec{x} - 2\vec{y}) \longrightarrow S(-2\vec{x}) \oplus S(-2\vec{y}) \longrightarrow S \longrightarrow S/(x^2, y^2) \longrightarrow 0.$$

This yields the L_f -graded matrix factorization $Q := \left(F_0 \begin{smallmatrix} f_0 \\ \xrightarrow{\quad} \\ f_1 \end{smallmatrix} F_1 \right)$ of f such that

$$F_0 = S(\vec{0}) \oplus S(\vec{f} - 2\vec{x} - 2\vec{y}), \quad F_1 := S(\vec{f} - 2\vec{x}) \oplus S(\vec{f} - 2\vec{y}),$$

and f_0 and f_1 are given by the matrices

$$q_0 = \begin{pmatrix} x^{\alpha-2}y^\beta & -y^2 \\ y & x^2 \end{pmatrix} \text{ and } q_1 = \begin{pmatrix} x^2 & y^2 \\ -y & x^{\alpha-2}y^\beta \end{pmatrix}$$

respectively.

Definition 7. We define a singularity $(X_Q, 0)$ by

$$\mathbf{F}_Q(w, x, y) := (F_{Q,1}(w, x, y), F_{Q,2}(w, x, y)) = \left(x^{\alpha-2}y^\beta + \frac{w}{x}y^2, y - \frac{w}{x}x^2\right).$$

If we substitute $y = \frac{w}{x}x^2$ in $F_{Q,1}$, we obtain the function

$$F_Q(w, x) := F_{Q,1}\left(w, x, \frac{w}{x}x^2\right) = w^3x + x^{\alpha+\beta-2}w^\beta.$$

Thus $(X_Q, 0)$ is in fact a hypersurface singularity defined by $F_Q = 0$.

We get the function f back from F_Q by the reduction considered in [10, p. 19], namely F_Q is of the form $F_Q(w, x) = w^3x + x^3A(w, x)$, where $A(w, x) = x^{\alpha+\beta-5}w^\beta$, and $RF_Q(w, x) = w^3 + x^5A\left(\frac{w}{x}, x\right) = f(x, w)$.

3. Graded matrix factorizations of size two for regular systems of weights with $\varepsilon = -1$

We shall now consider the graded matrix factorizations of size two for the regular systems of weights with $\varepsilon = -1$. They were determined in [12]. There are 22 such systems of weights. They define the Fuchsian hypersurface singularities of genus 0 (see, e.g., [7]). Let $W = (a, b, c; h)$ be such a weight system. By definition, it is reduced. The number ε_W is defined by $\varepsilon_W := a + b + c - h$. Let $A_W = (\alpha_1, \dots, \alpha_r)$ be the signature of the system of weights W . Then $(\alpha_1, \dots, \alpha_r)$ corresponds to the signature of the corresponding Fuchsian singularity. For $r = 3$, there are 14 such weight systems which correspond to the 14 exceptional unimodal singularities. Here the polynomial f_W is an invertible polynomial and its canonical system of weights coincides with $(a, b, c; h)$. For $r = 4$, there are 6 corresponding to the heads of the bimodal series. There are 2 more with $r = 5$. The corresponding equations are weighted homogeneous deformations f_W of an invertible polynomial, see Table 2.

This again implies that $L_{f_W} \cong \mathbb{Z}$. Thus all the regular systems of weights with $\varepsilon_W = -1$ can be given by weighted homogeneous deformations f of invertible polynomials with $\varepsilon_f = -1$ and $L_f \cong \mathbb{Z}$ and the L_f -graded matrix factorizations of f correspond to the graded matrix factorizations considered in [12]. The Dolgachev numbers are equal to the signature of the system of weights.

It will turn out that the graded matrix factorizations of size 2 correspond to some of these singularities, but also to the Fuchsian complete intersection singularities of genus 0 (see [7]). There are 8+5+2 of them for certain signatures $(\alpha_1, \dots, \alpha_r)$.

Let W be a regular system of weights with $\varepsilon_W = -1$ and signature $A_W = (\alpha_1, \dots, \alpha_r)$. In [12], a full strongly exceptional collection in the triangulated category of graded matrix factorizations associated to W was constructed. The full strongly exceptional collection is described by the graph shown in Fig. 1.

It turns out that a graded matrix factorization of size 2 corresponds to the end point V_{i, α_i} of some arm.

Theorem 1. Let f be a weighted homogeneous deformation of an invertible polynomial with $\varepsilon_f = -1$ and $L_f \cong \mathbb{Z}$. Let $A = (\alpha_1, \dots, \alpha_r)$ be the Dolgachev

Table 2. Type of f_W

W	f_W	Type	p_1, p_2, p_3	$A = A_W$
6,14,21;42	$-(y^3 - x^7) + z^2$	I, $m = 1$	3,7,2	3,7,2
4,10,15;30	$-x(y^5 - x^2) + z^2$	$\Pi_1, m = 1$	3,5,2	4,5,2
3,8,12;24	$-x(y^4 - x) + z^3$	$\Pi_1, m = 1$	2,4,3	3,4,3
6,8,15;30	$-x(y^3 - x^4) + z^2$	$\Pi_1, m = 1$	5,3,2	8,3,2
4,6,11;22	$-xy(y^2 - x^3) + z^2$	III, $m = 1$	4,3,2	6,4,2
3,5,9;18	$-x(y^3 - x) + yz^3$	IV, $m = 1$	2,3,3	3,5,3
4,5,10;20	$-x(y^2 - x) + z^5$	$\Pi_1, m = 1$	2,2,5	5,2,5
3,4,8;16	$-x(y^2 - x) + yz^4$	IV, $m = 1$	2,2,4	4,3,4
6,8,9;24	$-x(y^2 - x^3) + z^3$	$\Pi_1, m = 1$	4,2,3	9,2,3
4,6,7;18	$-x(y^3 - x^2) + yz^2$	IV, $m = 1$	3,3,2	4,7,2
3,5,6;15	$-xy(y - x^2) + z^3$	III, $m = 1$	3,2,3	6,3,3
4,5,6;16	$-x(y^2 - x^3) + yz^2$	IV, $m = 1$	4,2,2	6,5,2
3,4,5;13	$-z(xz - y^2) + yx^3$	V	3,2,2	3,4,5
3,4,4;12	$-xy(y - x) + z^4$	III, $m = 1$	2,2,4	4,4,4
2,6,9;18	$-x(y^3 - x)(y^3 - \lambda_4 x) + z^2$	$\Pi_1, m = 2$	3,6,2	2,3,2,2
2,4,7;14	$-xy(y - x^2)(y - \lambda_4 x^2) + z^2$	III, $m = 2$	5,3,2	4,2,2,2
2,4,5;12	$-x(y^2 - x)(y^2 - \lambda_4 x) + yz^2$	IV, $m = 2$	3,4,2	2,5,2,2
2,3,6;12	$-(z^3 - x)(z^3 - \lambda_4 x) + xy^2$	$\Pi_2, m = 2$	2, 2, 6	3, 3, 2, 2
	$-(z^2 - x)(z^2 - \lambda_4 x) + xy^3$		2, 3, 4	2, 2, 3, 3
2,3,4;10	$-x(y - x^2)(y - \lambda_4 x^2) + yz^2$	IV, $m = 2$	5,2,2	4,3,2,2
2,3,3;9	$-x(y - x)(y - \lambda_4 x) + yz^3$	IV, $m = 2$	3,2,3	3,2,3,3
2,2,5;10	$-xy(y - x)(y - \lambda_4 x)(y - \lambda_5 x) + z^2$	III, $m = 3$	4,4,2	2,2,2,2,2
2,2,3;8	$-x(y - x)(y - \lambda_4 x)(y - \lambda_5 x) + yz^2$	IV, $m = 3$	4,3,2	2,3,2,2,2

Fig. 1. The graph $T_{\alpha_1, \dots, \alpha_r}^+$

numbers of f . If the L_f -graded matrix factorization V_{i,α_i} is of rank 2, then the corresponding singularity $\mathbf{F}_{V_{i,\alpha_i}}$ has the signature $(\tilde{\alpha}_1, \dots, \tilde{\alpha}_r)$ with $\tilde{\alpha}_i = \alpha_i + 1$ and $\tilde{\alpha}_j = \alpha_j$ for $j \neq i$.

Proof. We consider the matrix factorizations for the cases of Table 1 and examine conditions under which there exist maps of the local ring of the corresponding complete intersection singularity to the ring of a weighted projective line of type $(\tilde{\alpha}_1, \dots, \tilde{\alpha}_r)$. We indicate the matrix factorization, the mapping, the index i where we have an increase of α_i by one, and the condition under which the mapping gives an embedding.

(I) $m \geq 2, m|p_1, m|p_2$. We consider the matrix factorization

$$q_0 = \begin{pmatrix} \prod_{i=4}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1}{m}}) & -z \\ z^{p_3-1} & -(y^{\frac{p_2}{m}} - x^{\frac{p_1}{m}}) \end{pmatrix}.$$

Mapping:

$$\begin{aligned} x &= X_1 X_3^{\frac{p_2}{m}}, \quad y = X_2 X_3^{\frac{p_1}{m}}, \quad z = X_3^{(m-1)\frac{p_1 p_2}{m^2}} X_4 \cdots X_{m+2}, \\ w &= X_4^{p_3-1} \cdots X_{m+2}^{p_3-1} \end{aligned}$$

Increase of α_3 under the condition:

$$(m-1) \frac{p_1 p_2}{m^2} (p_3 - 1) = \frac{p_1 p_2}{m^2} + p_3 + 1. \quad (1)$$

(II) $m \geq 1, m|(p_1 - 1), m|p_2$. We have two essentially different matrix factorizations:

(a)

$$q_0 = \begin{pmatrix} \prod_{i=3}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -z \\ z^{p_3-1} & -x \end{pmatrix}.$$

Mapping:

$$\begin{aligned} x &= X_1^{p_3 + \frac{m+p_2}{p_1-1}}, \quad y = X_1 X_2, \quad z = X_1^{p_2} X_3 \cdots X_{m+2}, \\ w &= X_3^{p_3-1} \cdots X_{m+2}^{p_3-1} \end{aligned}$$

Increase of α_1 under the condition:

$$p_2(p_3 - 1)(p_1 - 1) = p_3(p_1 - 1) + m + p_2. \quad (2)$$

(b)

$$q_0 = \begin{pmatrix} x \prod_{i=4}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -z \\ z^{p_3-1} & -(y^{\frac{p_2}{m}} - x^{\frac{p_1-1}{m}}) \end{pmatrix}.$$

Mapping:

$$x = X_1^{p_3} X_3^{\frac{p_2}{m}}, \quad y = X_2 X_3^{\frac{p_1-1}{m}}, \quad z = X_1 X_3^{\frac{p_2}{m} + (m-1)\frac{(p_1-1)p_2}{m^2}} X_4 \cdots X_{m+2},$$

$$w = X_1^{p_3-1} X_4^{p_3-1} \cdots X_{m+2}^{p_3-1}$$

Increase of α_3 under the condition:

$$\begin{aligned} & \left(\frac{p_2}{m} + (m-1) \frac{(p_1-1)p_2}{m^2} \right) (p_3-1) \\ &= \frac{(p_1-1)p_2}{m^2} + p_3 + 1. \end{aligned} \quad (3)$$

(II₂) $m \geq 2, m|p_1, m|p_3$. We consider the matrix factorization

$$q_0 = \begin{pmatrix} \prod_{i=4}^{m+2} (z^{\frac{p_3}{m}} - \lambda_i x^{\frac{p_1}{m}}) & -y \\ xy^{p_2-1} & -(z^{\frac{p_3}{m}} - x^{\frac{p_1}{m}}) \end{pmatrix}.$$

Mapping:

$$\begin{aligned} x &= X_1^{p_3} X_3^{\frac{p_3}{m}}, \quad y = X_3^{(m-1)\frac{p_1 p_3}{m^2}} X_4 \cdots X_{m+2}, \quad z = X_1 X_2 X_3^{\frac{p_1}{m}}, \\ w &= X_1^{(m-1)\frac{p_3}{m}} X_4^{p_2-1} \cdots X_{m+2}^{p_2-1} \end{aligned}$$

Increase of α_3 under the condition:

$$\frac{p_3}{m} + (m-1) \frac{p_1 p_3}{m^2} (p_2-1) = (m-1) \frac{p_1 p_3}{m^2} + p_2 + 1. \quad (4)$$

One can show that the other possible matrix factorization

$$q_0 = \begin{pmatrix} \prod_{i=4}^{m+2} (z^{\frac{p_3}{m}} - \lambda_i x^{\frac{p_1}{m}}) & -x \\ y^{p_2} & -(z^{\frac{p_3}{m}} - x^{\frac{p_1}{m}}) \end{pmatrix}$$

has no solution.

(III) $m \geq 1, m|(p_1-1), m|(p_2-1)$. We have the essentially different matrix factorizations:

(a)

$$q_0 = \begin{pmatrix} y \prod_{i=3}^{m+2} (y^{\frac{p_2-1}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -z \\ z^{p_3-1} & -x \end{pmatrix}.$$

Mapping:

$$\begin{aligned} x &= X_1^{p_3 + \frac{(m-1)+p_2}{p_1-1}}, \quad y = X_1 X_2^{p_3}, \quad z = X_1^{p_2} X_2 X_3 \cdots X_{m+2}, \\ w &= X_2^{p_3-1} X_3^{p_3-1} \cdots X_{m+2}^{p_3-1} \end{aligned}$$

Increase of α_1 under the condition:

$$p_2(p_3-1)(p_1-1) = p_3(p_1-1) + (m-1) + p_2. \quad (5)$$

(b)

$$q_0 = \begin{pmatrix} xy \prod_{i=4}^{m+2} (y^{\frac{p_2-1}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -z \\ z^{p_3-1} & -(y^{\frac{p_2-1}{m}} - x^{\frac{p_1-1}{m}}) \end{pmatrix}.$$

Mapping:

$$x = X_1^{p_3} X_3^{\frac{p_2-1}{m}}, y = X_2^{p_3} X_3^{\frac{p_1-1}{m}}, z = X_1 X_2 X_3^{\frac{p_1-1}{m} + \frac{p_2-1}{m} + (m-1)\frac{(p_1-1)(p_2-1)}{m^2}}$$

$$X_4 \cdots X_{m+2}, w = X_1^{p_3-1} X_2^{p_3-1} X_4^{p_3-1} \cdots X_{m+2}^{p_3-1}$$

Increase of α_3 under the condition:

$$\left(\frac{p_1-1}{m} + \frac{p_2-1}{m} + (m-1)\frac{(p_1-1)(p_2-1)}{m^2} \right) (p_3-1)$$

$$= \frac{(p_1-1)(p_2-1)}{m^2} + p_3 + 1. \quad (6)$$

(IV) $m \geq 1, m|(p_1-1), m|p_2$. We have three essentially different factorizations.

(a)

$$q_0 = \begin{pmatrix} \prod_{i=3}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -z \\ yz^{p_3-1} & -x \end{pmatrix}.$$

Mapping:

$$x = X_1^{p_3 + \frac{m+p_2}{p_1-1}} X_2, y = X_1 X_2^{p_1}, z = X_1^{p_2} X_3 \cdots X_{m+2},$$

$$w = X_2^{p_1-1} X_3^{p_3-1} \cdots X_{m+2}^{p_3-1}$$

Increase of α_1 under the condition:

$$p_2(p_3-1)(p_1-1) = p_3(p_1-1) + (m+1-p_1) + p_2. \quad (7)$$

(b)

$$q_0 = \begin{pmatrix} \prod_{i=3}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -y \\ z^{p_3} & -x \end{pmatrix}.$$

Mapping:

$$x = X_1^{p_3} X_2^{p_3}, y = X_2^{p_1 + \frac{(p_1-1)(p_3-1)+m}{p_2}}, z = X_1 X_2 X_3 \cdots X_{m+2},$$

$$w = X_3^{p_3} \cdots X_{m+2}^{p_3}$$

Increase of α_2 under the condition:

$$p_2(p_1(p_3-1) - p_3) = (p_1-1)(p_3-1) + m. \quad (8)$$

This equation is equivalent to Eq. (7).

(c)

$$q_0 = \begin{pmatrix} x \prod_{i=4}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -z \\ yz^{p_3-1} & -(y^{\frac{p_2}{m}} - x^{\frac{p_1-1}{m}}) \end{pmatrix}.$$

Mapping:

$$x = X_1^{p_3} X_2 X_3^{\frac{p_2}{m}}, \quad y = X_2^{p_1} X_3^{\frac{p_1-1}{m}}, \quad z = X_1 X_3^{\frac{p_2}{m} + (m-1)\frac{(p_1-1)p_2}{m^2}} X_4 \cdots X_{m+2},$$

$$w = X_1^{p_3-1} X_2^{1+(m-1)\frac{p_1-1}{m}} X_4^{p_3-1} \cdots X_{m+2}^{p_3-1}$$

Increase of α_3 under the condition:

$$\left(\frac{p_2}{m} + (m-1)\frac{(p_1-1)p_2}{m^2} \right) (p_3-1) + \frac{p_1-1}{m} = \frac{(p_1-1)p_2}{m^2} + p_3 + 1. \quad (9)$$

Similarly as in (Π_2) , one can show that the other matrix factorization

$$q_0 = \begin{pmatrix} x \prod_{i=4}^{m+2} (y^{\frac{p_2}{m}} - \lambda_i x^{\frac{p_1-1}{m}}) & -y \\ z^{p_3} & -(y^{\frac{p_2}{m}} - x^{\frac{p_1-1}{m}}) \end{pmatrix}.$$

has no solution.

(d) We also have to consider a special matrix factorization in the case $m = 1$. We can write the equation as

$$f(x, y, z) = -y(xy^{p_2-1} - z^{p_3}) + x^{p_1}.$$

Then we have the matrix factorization

$$q_0 = \begin{pmatrix} xy^{p_2-1} - z^{p_3} & -x \\ x^{p_1-1} & -y \end{pmatrix}$$

Mapping:

$$x = X_2^{p_3} X_3^{p_3}, \quad y = x_2^{p_1 + \frac{2}{p_2-1}}, \quad z = X_1 X_2 X_3, \quad w = X_3^{p_3(p_1-1)}$$

Increase of α_2 under the condition:

$$p_3(p_1-1)(p_2-1) = p_1(p_2-1) + 2. \quad (10)$$

This equation is equivalent to Eq. (8) with $m = 1$.

(V) The equation is

$$f(x, y, z) = -z(xz^{p_3-1} - y^{p_2}) + yx^{p_1}.$$

Then we have two matrix factorizations:

$$(a) \quad q_0 = \begin{pmatrix} xz^{p_3-1} - y^{p_2} & -x \\ yx^{p_1-1} & -z \end{pmatrix}$$

Mapping:

$$x = X_2^{p_2} X_3^{p_3}, \quad y = X_1^{p_3} X_2 X_3, \quad z = X_1 X_2^{p_1 p_2 - p_2 + 1}, \quad w = X_1^{p_3-1} X_3^{p_1 p_2 - p_2 + 1}$$

Increase of α_2 under the condition:

$$p_2(p_1-1)(p_3-1) = (p_1-1)(p_3-1) + 2. \quad (11)$$

$$(b) \quad q_0 = \begin{pmatrix} xz^{p_3-1} - y^{p_2} & -y \\ x^{p_1} & -z \end{pmatrix}$$

Mapping:

$$x = X_1 X_2 X_3^{p_2}, \quad y = X_1^{p_1 p_3 - p_1 + 1} X_3, \quad z = X_1^{p_1} X_2^{p_1}, \quad w = X_2^{p_1} X_3^{p_1 p_2}$$

Increase of α_1 under the condition (11).

In the cases I, II₁(b), II₂, III(b), and IV(c), one can exchange the factor with $\lambda_3 (= 1)$ in the lower right corner of the matrix factorization by any of the factors corresponding to λ_i , $i = 4, \dots, m+2$, and interchange X_3 and X_i . This leads to an increase of α_i instead of α_3 . In Case III(a), one can interchange the variables x and y to get an increase of α_2 .

The solutions to the Eqs. (1)–(11) are precisely the entries of Table 2 and two extra ones. Equation (1) has the solution I, $m = 2$, $(p_1, p_2, p_3) = (4, 4, 3)$, $A = (2, 2, 3, 3)$. This corresponds to the minimally elliptic singularity $V'_{(1)}$ of [6] with weight system $W = (3, 3, 4; 12)$ and $\varepsilon_W = -2$. Equation (9) has the additional solution IV, $m = 1$, $(p_1, p_2, p_3) = (3, 3, 4)$, $A = (8, 7, 4)$. This corresponds to a singularity with the weight system $W = (12, 8, 7; 36)$ with $\varepsilon_W = -9$. $\square$

Remark 1. The proof shows that the statement of Theorem 1 only holds in this very special situation. Namely, it shows that, except for two exceptional cases, the weighted homogeneous deformations f of an invertible polynomial with $\varepsilon_f = -1$ are the only weighted homogeneous deformations of an invertible polynomial with the property that there exists an L_f -graded matrix factorization Q of size 2 such that the corresponding singularity $\mathbf{F}_Q$ has the signature $(\tilde{\alpha}_1, \dots, \tilde{\alpha}_r)$ with $\tilde{\alpha}_i = \alpha_i + 1$ and $\tilde{\alpha}_j = \alpha_j$ for $j \neq i$ where $A = (\alpha_1, \dots, \alpha_r)$ are the Dolgachev numbers of f .

Remark 2. The reduction is a relation between the singularities. Not all possible extensions of the signatures are realised. For example, the following pairs are missing: $(2, 3, 7) \not\leftarrow (2, 3, 8)$, $(2, 4, 5) \not\leftarrow (2, 4, 6)$, $(3, 3, 4) \not\leftarrow (3, 3, 5)$, $(2, 2, 2, 3) \not\leftarrow (2, 2, 2, 4)$. In these cases, the graded matrix factorization V_{r, α_r} for the singularities with signature $(2, 3, 7)$, $(2, 4, 5)$, $(3, 3, 4)$, and $(2, 2, 2, 3)$ is of rank 4. In these cases, the function f_W can be given by $f_W(x, y, z) = f'_W(x, y) + z^2$, where $f'_W(x, y)$ is the invertible polynomial in two variables given by

$$(2, 3, 7): x^7 + y^3 \quad (2, 4, 5): x^5 y + y^3 \quad (3, 3, 4): x^8 + y^3 \quad (2, 2, 2, 3): x^6 y + y^3.$$

We shall consider the L_f -graded matrix factorization of size 2 of f'_W and the corresponding reduction R defined in Sect. 2(B). We enhance the reduction by these cases.

Example 1. We consider the case $W = (4, 10, 15; 30)$, $A_W = (2, 4, 5)$, $f_W = x^5 y + y^3 + z^2$, $f'_W = x^5 y + y^3$. This is the singularity E_{13} in Arnold's notation. We consider the graded matrix factorization of f'_W given by

$$q_0 = \begin{pmatrix} x^3 y & -y^2 \\ y & x^2 \end{pmatrix}.$$

Then $\mathbf{F}_Q(w, x) = w^3 x + w x^4$. This is the singularity Z_{12} with signature $(2, 4, 6)$.

Table 3. Dolgachev numbers, Kodaira types, and grades

Type of Dolgachev numbers	Kodaira type	ℓ	$D = -Z^2$
$2, 3, c; c \geq 7$	$\text{II}(k)$	1	$c - 6$
$2, b, c; b, c \geq 4$	$\text{III}(k_1, k_2)$	2	$b + c - 8$
$a, b, c; a, b, c \geq 3$	$\text{IV}(k_1, k_2, k_3)$	3	$a + b + c - 9$
$a, b, c, d; a, b, c, d \geq 2$	$\text{I}_0^*(k_1, k_2, k_3, k_4)$	4	$a + b + c + d - 8$

The isolated hypersurface singularities defined by regular systems of weights with $\varepsilon = -1$ and $r \leq 4$ and the isolated complete intersection singularities which reduce to them are Kodaira singularities [10] or minimally elliptic singularities [13]. The minimal resolution is of Kodaira type II, III, IV or I_0^* (see [10]) and it carries a fundamental cycle Z (see [3, p. 132]). The self-intersection number of the fundamental cycle $D := -Z^2$ is called the *grade* of the singularity [10]. Let $(X, 0)$ be any of the singularities of Table 2 with Dolgachev numbers $\alpha_1, \dots, \alpha_r$ and $r \leq 4$ or of the isolated complete intersection singularities which reduce to them. According to [10] or [13], there is the correspondence between certain types of Dolgachev numbers, Kodaira types, the number ℓ of parameters k_i of the Kodaira type, and the grades indicated in Table 3 (see also [15, p. 515]).

By [16, Corollary 7.9.2], the Kodaira type and hence the number ℓ of a singularity and its reduction are the same.

The possible reductions between these singularities are visualized in Fig. 2. Here a usual arrow refers to a reduction corresponding to Theorem 1, a dashed arrow to the special reduction R described in Sect. 2(B). The Dolgachev numbers of the regular systems of weights for $\varepsilon = -1$ are denoted by usual letters, those of the remaining singularities are denoted by scriptstyle letters. We also indicate the names of Arnold for the hypersurface singularities [2] and of Wall [16] (see also [7]) for the complete intersection singularities.

On the other hand, each of these singularities has another triple or quadruple of numbers, the *Gabrielov numbers*, see [8] for the definition in the case of the hypersurface singularities and [9] in the case of the complete intersection singularities. They also have types like the Dolgachev numbers, but for them the columns corresponding to ℓ and D are interchanged, see Table 4. Note that the grade D corresponds to the type of Gabrielov numbers, see [13, Theorem 3.13]. More precisely, according to [13, Theorem 3.13], for $1 \leq D \leq 3$, the singularity is a hypersurface singularity and, for $D = 4$, it is a complete intersection singularity. For $D = 1$, the singularity is given by an equation of the form $z^2 + y^3 + g(x, y, z) = 0$, where the polynomial $g(x, y, z)$ has only terms of degree ≥ 4 , see also [13, Table 1]. For $D = 2$, the singularity is given by an equation of the form $z^2 + h(x, y) = 0$, where the polynomial $h(x, y)$ has only terms of degree ≥ 4 , see also [13, Table 2]. Finally, for $D = 3$, the equation is $f(x, y, z) = 0$, where the polynomial $f(x, y, z)$ has only terms of degree ≥ 3 .

The hypersurface singularities with $r = 3$ are the 14 exceptional unimodal singularities. There is Arnold’s strange duality [2] between these singularities which interchanges Dolgachev and Gabrielov numbers.

Fig. 3. The pyramid of s-adjacencies between the 14 exceptional unimodal singularities

Remark 3. Note that the s-adjacency is a special case of adjacency. For example, the general adjacency between the 14 exceptional unimodal singularities corresponds to the natural partial ordering between the Gabrielov numbers, see [4].

A comparison of Fig. 2 restricted to the 14 exceptional unimodal singularities, Fig. 3, and Arnold’s strange duality yields the following corollary of Theorem 1.

Corollary 1. *Denote Arnold’s strange duality by $X \leftrightarrow X^*$. If a singularity Y is a reduction of X , then the dual singularity X^* is s-adjacent to the dual singularity Y^* .*

Acknowledgements This work has been partially supported by DFG. The second named author is also supported by JSPS KAKENHI Grant Numbers JP16H06337 and JP21H04994.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly

from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data Availability Statement Data sharing not applicable to this article as no datasets were generated or analysed during the current study

Funding Open Access funding enabled and organized by Projekt DEAL.

References

- [1] Aramaki, D., Takahashi, A.: Maximally-graded matrix factorizations for an invertible polynomial of chain type. *Adv. Math.* **373**, 107320 (2020)
- [2] Arnold, V.I., Gusein-Zade, S.M., Varchenko, A.N.: *Singularities of Differentiable Maps*, vol. I. Birkhäuser, Boston Basel Berlin (1985)
- [3] Artin, M.: On isolated rational singularities of surfaces. *Am. J. Math.* **88**, 129–136 (1966)
- [4] Brieskorn, E.: Die Hierarchie der 1-modularen Singularitäten. *Manuscripta Math.* **27**(2), 183–219 (1979)
- [5] Buchweitz, R.-O., Greuel, G.-M., Schreyer, F.-O.: Cohen-Macaulay modules on hypersurface singularities. II. *Invent. Math.* **88**(1), 165–182 (1987)
- [6] Ebeling, W.: The Milnor lattices of the elliptic hypersurface singularities. *Proc. London Math. Soc.* (3) **53**(1), 85–111 (1986)
- [7] Ebeling, W.: The Poincaré series of some special quasihomogeneous surface singularities. *Publ. Res. Inst. Math. Sci.* **39**(2), 393–413 (2003)
- [8] Ebeling, W., Takahashi, A.: Strange duality of weighted homogeneous polynomials. *Compositio Math.* **147**, 1413–1433 (2011)
- [9] Ebeling, W., Takahashi, A.: Strange duality between hypersurface and complete intersection singularities. *Arnold Math. J.* **2**, 277–298 (2016)
- [10] Ebeling, W., Wall, C.T.C.: Kodaira singularities and an extension of Arnold’s strange duality. *Compositio Math.* **56**, 3–77 (1985)
- [11] Geigle, W., Lenzing, H.: A class of weighted projective curves arising in representation theory of finite-dimensional algebras. In: *Singularities, representation of algebras, and vector bundles* (Lambrech, 1985), pp. 265–297, *Lecture Notes in Math.*, **1273**, Springer, Berlin, 1987
- [12] Kajiura, H., Saito, K., Takahashi, A.: Triangulated categories of matrix factorizations for regular systems of weights with $\varepsilon = -1$. *Adv. Math.* **220**, 1602–1654 (2009)
- [13] Laufer, H.B.: On minimally elliptic singularities. *Am. J. Math.* **99**(6), 1257–1295 (1977)
- [14] Laufer, H.B.: Ambient deformations for exceptional sets in two-manifolds. *Invent. Math.* **55**(1), 1–36 (1979)
- [15] Saito, K.: Regular system of weights and associated singularities. *Complex Analytic Singularities. Adv. Stud. Pure Math.* **8**, 479–526 (1987)
- [16] Wall, C.T.C.: Notes on the classification of singularities. *Proc. London Math. Soc.* (3) **48**(3), 461–513 (1984)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.