Correction to: Does stent type impact coil embolization outcomes in extended follow-up of small-sized aneurysms (< 10 mm)?

Jeongjun Lee¹ · Young Dae Cho² · Dong Hyun Yoo² · Hyun-Seung Kang³ · Won-Sang Cho³ · Jeong Eun Kim³ · Jusun Moon⁴ · Moon Hee Han^{2,3}

Published online: 7 May 2018

© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: Neuroradiology 2018

<https://doi.org/10.1007/s00234-018-2022-4>

The original version of this article contained a mistake in Figure 4: Figure 4a and d were interchanged. The correct figure is shown below. The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s00234-018-2022-4>

✉ Young Dae Cho
aronnn@naver.com

¹ Department of Neurosurgery, Dongguk University Ilsan Hospital, College of Medicine, Dongguk University, Ilsan, South Korea

² Department of Radiology, Seoul National University Hospital, Seoul National University College of Medicine, 101 Daehak-ro, Jongno-gu, Seoul 03080, South Korea

³ Department of Neurosurgery, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, South Korea

⁴ Department of Neurology, Korea University Anam Hospital, Korea University College of Medicine, Seoul, South Korea

Fig. 4 Representative case of minor recanalization in coiled aneurysm with Neuroform stent. **a** Preembolization DSA of posterior communicating artery aneurysm. **b–d** Six-month follow-up TOF-MRA (arrow indicates the minor recanalized portion; **b** MIP image; **c** source image; **d** DSA)