



## Correction to: Near-occlusion is difficult to diagnose with common carotid ultrasound methods

Elias Johansson<sup>1,2</sup> · Davide Vanoli<sup>3</sup> · Isa Bråten-Johansson<sup>1</sup> · Lucy Law<sup>1</sup> · Richard I Aviv<sup>4</sup> · Allan J Fox<sup>4</sup>

Published online: 21 March 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

### Correction to: *Neuroradiology* (2021)

<https://doi.org/10.1007/s00234-021-02687-x>

Unfortunately, the article was published online prematurely with errors. Initials were presented as numbers and not yet replaced with the actual initials. A translation legend for blinded information was needlessly included.

The original article has been corrected.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s00234-021-02687-x>

---

✉ Elias Johansson  
elias.johansson@umu.se

<sup>1</sup> Department of Clinical Science, Umeå University, Umeå, Sweden

<sup>2</sup> Wallenberg Center for Molecular Medicine (WCMM), Umeå University, Umeå, Sweden

<sup>3</sup> Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden

<sup>4</sup> Department of Medical Imaging, Sunnybrook Health Science Center, University of Toronto, Toronto, Canada