

## Erratum to: Simplified skin excision pattern for skin-sparing/reducing mastectomy and direct-to-implant single-stage immediate (DISSI) breast reconstruction

B. Atiyeh<sup>1</sup> · N. Papazian<sup>1</sup> · H. Becker<sup>2</sup> · S. Dibo<sup>1</sup> · A. Ibrahim<sup>1</sup>

Published online: 3 July 2015

© Springer-Verlag Berlin Heidelberg 2015

### Erratum to: European Journal of Plastic Surgery DOI 10.1007/s00238-015-1107-0

The original version of this article contained a mistake in the order of authors as well as in the caption of Figs. 3, 4, 5, and 6. The correct order of authors is B. Atiyeh, N. Papazian, H. Becker, S. Dibo & A. Ibrahim.

On the other hand, caption of Figs. 3, 4, 5, and 6 should not include Becker 35. Below are the corrected captions:

Fig. 3 **a, b, c** Patient with ptosis and right breast upper quadrant tumor. **d** Preoperative surgical planning with extended skin excision over the upper lateral quadrant. **e** Medially

based dermal barrier flap—DBF. Pectoralis major—PM detached inferiorly. 495 cc Siltex™ Contour Profile®, Cohesive II™ medium height high profile prosthesis inserted. DBF and PM sutured together to prevent muscle retraction and to maintain the implant firmly in position. **f** Immediate result of right IBR and contralateral circumvertical mastopexy. **g, h, i** Result at 3 months

Fig. 4 **a, b, c** Patient presenting for bilateral SRM and IBR. **d, e, f** One month postoperative result following reconstruction with 555 cc Siltex™ Contour Profile®, Cohesive II™ medium height high profile prosthesis. **g, h, i** One year result following nipple-areola reconstruction

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00238-015-1107-0>.

---

✉ A. Ibrahim  
ai12@aub.edu.lb

<sup>1</sup> Plastic and Reconstructive Surgery, American University of Beirut, Medical Center, Beirut, Lebanon

<sup>2</sup> Plastic and Reconstructive Surgery, 670 Glades Road, Suite 220, Boca Raton, FL 33431, USA

Fig. 5 **a, b, c** Patient presenting for bilateral SRM and IBR. **d, e, f** One month postoperative result following reconstruction with 620 cc Siltex™ Contour Profile®, Cohesive II™ medium height high profile prosthesis. **g, h, i** One year postoperative result

Fig. 6 **a, b, c** Patient presenting for bilateral SRM and IBR. **d, e** Preoperative planning for areola preserving mastectomy. **f, g, h** Six months postoperative result of IBR with 555 cc Siltex™ Contour Profile®, Cohesive II™ medium height high profile prosthesis