



Correction to: Current trends in operative treatment of carpometacarpal osteoarthritis: a survey of European hand surgeons

Qi Yin¹ · Merel J.-L. Berkhout^{2,3} · Marco J. P. F. Ritt^{1,3}

Published online: 8 June 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: **European Journal of Plastic Surgery**

<https://doi.org/10.1007/s00238-019-01528-8>

The entries in Tables 1, 2 and 3 were captured as $N = 433439$ instead of $N = 439$. The correct tables are shown below.

Table 1 Survey demographics: country of practice

Country of practice	Response, $N = 439$	% Total responses
Austria	25	5.7
Belgium	26	5.9
Bulgaria	3	0.7
Czech Republic	12	2.7
Estonia	6	1.4
Finland	4	0.9
France	39	8.9
Germany	99	22.6
Greece	22	5.0
Hungary	17	3.9
Ireland	1	0.2
Italy	2	0.5
Latvia	6	1.4
Netherlands	72	16.4
Norway	10	2.3
Poland	8	1.8
Portugal	3	0.7
Romania	1	0.2
Sweden	15	3.4
Switzerland	45	10.3
Turkey	23	5.2

The online version of the original article can be found at <https://doi.org/10.1007/s00238-019-01528-8>

✉ Qi Yin
Q.Yin@vumc.nl

¹ Department of Plastic, Reconstructive and Hand Surgery, Amsterdam UMC, Location VUmc, P.O. Box 7057, 1007 MB Amsterdam, Netherlands

² Department of Plastic, Reconstructive and Hand Surgery, Alrijne Hospital, P.O. Box 4220, 2350 CC Leiderdorp, Netherlands

³ The Hand Clinic, Keienbergweg 8, 1101 GB Amsterdam, Netherlands

Table 2 Survey demographics: years of experience as hand surgeon

Years of experience as hand surgeon	Response, <i>N</i> = 439	% Total responses
≤5	74	16.9
6–15	179	40.8
16–25	122	27.8
>25	64	14.6

Table 3 Preferred surgical procedures for CMC-I OA, arranged by popularity

Preferred surgical procedure	Response	% Responses of total participants, <i>N</i> = 439
Trapeziectomy with LRTI	202	46.0
Prosthetic joint replacement	113	25.7
Metal on polyethylene/metal	86	76.1
Pyrocarbon	21	18.6
Bioresorbable	3	2.7
Other	3	2.7
Trapeziectomy with interpositional arthroplasty	100	22.8
Tendon	89	89.0
Capsule	7	7.0
Other	4	4.0
Trapeziectomy only	93	21.2
Trapeziectomy with ligament reconstruction only	82	18.7
Arthrodesis of MC I and trapezium	33	7.5
Surgical denervation	24	5.5
Trapeziectomy with hematoma distraction only (temporary K-wire fixation)	20	4.6
Eaton-Littler ligament reconstruction	17	3.9
Wilson's osteotomy	14	3.2
Suspensionplasty with tendon, without trapeziectomy	8	1.8
Suspensionplasty with foreign material (e.g., Mini TightRope® or other suture), without trapeziectomy	6	1.4
Arthrodesis of MC I and MC II	0	0
Other	15	3.4

The sum of responses exceeds the total number of participants (*N* = 439), because more than one surgical procedure could be opted as a treatment of choice. In the second column, the responses are expressed as percentage of the total number of participants; these percentages thus do not add up to 100

The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.