



# Temporal and Spatial Distribution of Mercury in Gulls Eggs from the Iberian Peninsula

M. Glória Pereira<sup>1</sup> · Alan Lawlor<sup>1</sup> · Albert Bertolero<sup>2</sup> · Sergi Díez<sup>3</sup> · Richard F. Shore<sup>1</sup> · Silvia Lacorte<sup>3</sup>

Received: 20 August 2018 / Accepted: 27 November 2018 / Published online: 18 December 2018  
© The Author(s) 2018

## Abstract

We examined how coastal mercury contamination varied spatially and temporally across the Iberian Peninsula by measuring mercury concentrations in the eggs of the sentinel biomonitor yellow-legged gull (*Larus michahellis*). Samples were collected from eight colonies that ranged from the Atlantic across the south and northern areas of the Mediterranean. We also measured Hg residues in eggs of the one of the most endangered gull species in the world, the Audouin's gull (*Larus audouinii*) from the Ebro Delta, where colonies of yellow-legged and Audouin's gull co-occur. Fresh eggs were collected in 2009 and 2016 and samples were pooled from each colony for analysis. Mercury concentrations in yellow-legged gulls ranged between 0.4 and 2.8 mg/kg dry weight (dw); although there were no significant differences in concentrations between sampling periods, significant differences were found between colonies. Higher concentrations were associated with northern Mediterranean colonies (Columbretes and Ebro Delta), likely due to proximity to emission sources, circulatory marine currents and diet composition. Mercury concentrations in yellow-legged gull eggs were lower than those reported to result in impaired hatching. Residues in Audouin's gull eggs from the Ebro Delta were significantly higher (4.0–5.6 mg/kg dw) than those in yellow-legged gull from the same location, probably associated with dietary differences. Mercury levels in Audouin's gull were ten times above the benchmark suggested to reduce nest success by 10%. Overall, these results raise concern for adverse health impacts in this protected seabird species and further investigation in Audouin's gull eggs from the Ebro Delta is recommended.

Mercury (Hg) is a toxic, nonessential heavy metal that is released into the environment from natural and anthropogenic sources. Its wide spatial distribution is linked to emissions from various sources, such as fossil fuel combustion, chlor-alkali plants, and agriculture, and is considered a global environmental pollutant (Driscoll et al. 2013).

In Spain, point-source emissions of Hg include eight chlor-alkali plants and Hg mining activity responsible for most contamination at local and regional scales. Even though Hg mining ceased in 2004, Hg is still detected in a wide

range of environmental matrices, e.g., in soils and plants from the mining sites in Almaden (Molina et al. 2006) and from an abandoned cinnabar mining site (Garcia-Sanchez et al. 2009); in food web components collected close to a chlor-alkali plant (Carrasco et al. 2008, 2011a); in downstream (Carrasco et al. 2011b) river water and sediments of Catalonia (Roig et al. 2011) and Asturias (Garcia-Ordiales et al. 2018); and in marine macrophytes and sediments of the Spanish Mediterranean coast (Sanchiz et al. 2000).

Due to its physical and chemical properties, Hg occurs in the environment in different forms and it is of particular concern in aquatic ecosystems. Here, inorganic Hg is bio-transformed into the highly toxic, methylmercury (MeHg), which tends to bioaccumulate and bioconcentrate in food webs. Although MeHg tends to represent less than 1% of the total Hg in fresh and marine waters, c. 96% of the total Hg in top predators is MeHg (Wiener et al. 2003). Hg persistence is also attributed to its slow metabolism by vertebrates (Walker and Livingstone 1992). High levels of Hg exposure have been related to a wide range of adverse effects (Giesy et al. 1994), including reprotoxicity (Furness 1993;

✉ M. Glória Pereira  
mdgds@ceh.ac.uk

<sup>1</sup> Centre for Ecology and Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster LA1 4AP, UK

<sup>2</sup> Associació Ornitològica Picampall de les Terres de l'Ebre, Amposta, Catalonia, Spain

<sup>3</sup> Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research, IDÆA-CSIC, Jordi Girona 18-26, 08034 Barcelona, Catalonia, Spain

Grassman et al. 1998; Boening 2000; Kenow et al. 2003), with MeHg being considered highly neurotoxic for humans and wildlife (Diez 2009).

Organisms that readily respond to contamination events can be used as bioindicators of contamination, providing information on spatial and temporal variation in environmental concentrations (Bishop et al. 1995). Birds and their eggs have been used as such indicators with the use of coastal-nesting birds as sentinels to detect changes in the bioavailability and transfer of Hg through coastal marine systems being well established (Pereira et al. 2009; Dittmann et al. 2012; Champoux and Boily 2017). Hg residues, in particular, have been measured in eggs (contents rather than in eggshells, Peterson et al. 2017), because they tend to have a consistent composition unlike body tissues, which can vary markedly between and within species in mass and lipid content. Furthermore, Hg accumulation in eggs of income breeders species is attributed to the feeding habits of the female in the breeding area (Sanpera et al. 2000), reflecting recent dietary exposure (1–2 weeks before egg laying) (Furness 1993; Becker et al. 1994). Therefore, eggs appear to be good indicators of localised and contemporary environmental exposure. In Spain, Hg concentrations have been reported in the eggs of common terns (*Sterna hirundo*) (Guitart et al. 2003) and of Audouin's gulls (*Larus audouinii*) from the Ebro Delta (Morera et al. 1997; Garcia-Tarrason et al. 2013), in Audouin's gull eggs from Chafarinas Islands (Sanpera et al. 2000), and in yellow-legged gull (*Larus michahellis*) eggs from the Medes Islands (Sanpera et al. 1997).

Studies on Hg residues in gull eggs conducted to date in Portugal and Spain have focused mostly on single locations and do not provide information on how contamination patterns vary spatially. The Iberian Peninsula has both Mediterranean and Atlantic coasts and pollution levels are expected to vary as a result. The Mediterranean is an enclosed basin with limited exchange of deep water with the outer oceans; water circulation is dominated by salinity and temperature differences rather than winds. These geographic and climatological conditions make the Mediterranean a sink for pollutants from river and treated and untreated wastewater discharges, and run-off (Sanchez-Avila et al. 2009). Furthermore, there are areas that contain long-established chemical industries and associated contamination; for instance, large amounts of industrial waste containing high Hg concentrations (up to 436 µg/g) were dumped into a reservoir adjacent to the Ebro river (Carrasco et al. 2008, 2011a) with potential transfer downstream to the delta (Carrasco et al. 2011a, b). Compared with the Mediterranean coast, pollution in the Atlantic coast of the Iberian Peninsula is lower due to dilution by the seawater and oceanic currents (Yamashita et al. 2005), although there are hotspots of contamination associated with human activities (Rubio et al. 2000).


Selenium (Se) often co-occurs with Hg, and there is some evidence that Se may be protective of Hg toxicity in several organisms (Ralston and Raymond 2013; Cusack et al. 2017; Spiller 2018). We also determined Se concentrations in the eggs collected in 2009, because whilst Se is an essential element for developing embryos, excess levels produce embryotoxic and teratogenic effects, embryonic malformations, and increased mortality. Also, simultaneously high concentrations of Se and Hg can have synergistically negative effects on embryonic development in birds (Heinz and Hoffman 1998).

The overarching goal of the present study was to investigate how concentrations of Hg vary across the Iberian Peninsula, both in the Atlantic and Mediterranean coasts. Such information is important to understand which areas, and the wildlife that they support, may be at most risk from contamination. We assessed this variation by measuring Hg in the eggs of the bio-indicator species, yellow-legged gull (*Larus michahellis*). We used this species because of its wide breeding distribution and because it is a coastal feeder (Ramos et al. 2009). We report temporal and spatial variations in total Hg concentrations in eggs taken from nesting colonies around the Iberian Peninsula in 2009 and 2016. Where possible, we compare Hg concentrations in yellow-legged gull eggs with those in the eggs of Audouin's gulls, a highly endangered species. Until the mid-2000s, the Ebro Delta hosted 70% of the world's breeding population (Fernandez-Chacon et al. 2013), but this has declined to only 3% in 2017 (Genovart et al. 2018). Our goal was to determine whether residues in yellow-legged gull eggs can be used as a surrogate measure of exposure in Audouin's gulls. Finally, and given the uncertainty with regard to the protective effect of Se on Hg toxicity, we relate Hg to Se concentrations.

## Materials and Methods

### Sampling Sites and Egg Collection

Fresh, yellow-legged gull eggs were collected from eight colonies, in Spring 2009 and 2016, with the permission of National and Natural Parks. The sampling areas were representative of two geographical areas: the Atlantic Ocean and the Mediterranean Sea. The Atlantic colonies were Berlengas and Cies Islands, with the Mediterranean colonies located at the northern (NM: Medes, Ebro Delta and Columbretes Islands) and southern Spanish coasts (SM: Dragonera, Mar Menor and Chafarinas Islands) (Fig. 1; Table 1). The NM and SM sites are influenced by the Northern current and the Algerian current, respectively (Milot 1987, 1999). The characteristics of the different sites, including distance from mainland and anthropogenic



**Fig. 1** Colony locations from which eggs were sampled. These areas represent the most important gull colonies of the Iberian Peninsula and they all are declared Special Protection Area for birds

pressures, are described in Table 1. Fresh Audouin's gull eggs also were collected from the Ebro Delta in the 2009 and 2016 breeding seasons.

At each colony, 36 eggs were randomly collected from 3 different areas within the colony (hereafter termed sub-colonies) such that 12 eggs were taken per subcolony. We adopted the sampling method described by Vicente et al. (2012), who reported that sample pooling successfully encompassed within-colony variability of residue concentrations. It also followed the recommendations of the United Nations Environment Programme and OSPAR for sampling and processing of biota samples (OSPAR Commission 2007; UN Environment). We collected the first laid egg, because it typically contains the highest within-clutch contaminant concentration, resulting from transfer from female to eggs (Sanpera et al. 1997). The contents of 12 eggs from each subcolony were pooled. Thus, there were three pooled samples per colony for analysis, except where limited colony size meant that only 12 eggs in total were collected (Columbretes in both years, Mar Menor and Berlingas in 2016 and Dragonera in 2009). In those cases, there was only one pooled sample per colony for analysis. No eggs were collected from Dragonera in 2016.

## Hg Analysis

Hg concentrations were measured in approximately 1-g subsamples that were dried to constant weight at 80 °C for 24 h, solubilised at room temperature overnight in 2 ml of (Analar) nitric acid, then heated at 90 °C (20 min) and then 120 °C (1 h). To aid the digestion of the organic matter, 0.5 ml of 30% hydrogen peroxide was added, which was then heated at 120 °C for 15 min. Digests were diluted with double-deionised water to a known volume and 10% acid strength. The resulting digests were analysed for Hg by Inductively Coupled Plasma—Mass Spectrometry (ICP-MS) using a Perkin Elmer DRCII ICP-MS with standard operating conditions. Hg concentrations are expressed as mg/kg dry weight (dw). For quality control, blanks and three certified reference materials (pig kidney, TORT-2 and DOLT-4) were run alongside samples; average recoveries ( $n = 3$ ) in the reference materials were 93% (RSD = 2.1%), 104% (RSD = 10.1%), and 98.6% (RSD = 17.3%), respectively. The sample limit of detection (LoD) and was determined as  $3 \times$  the standard deviation of the blank and was 0.1 mg/kg. Samples were blank corrected, when blank concentrations were above LoD.

**Table 1** Description of the colonies sampled in 2009 and 2016, ordered from northeast to northwest of the Iberian Peninsula, area of the colony, *L. michahellis* population in all colonies and *L. audouin* population in the Ebro Delta, food resources, distance from main human settlements and human impact

Colony	Area (km <sup>2</sup> )	<i>L. michahellis</i> population	Resources	Distance from human settlements (km)	Human impact	References
Medes 3°13'E 42°02'N	1.8	7291	Refuse tips	0.87	Industry, tourism	Bosch et al. (1994)
Ebro Delta 0°40'E 40°35'N	320	9744 14,177 <i>L. audouin</i>	Refuse tips, waste, crabfish Benthic-mesopelagic fish	7.5	Rice fields, chlor-alkali plant	Ramos et al. (2009) and Oro et al. (2009)
Columbretes 0°41'E 39°54'N	0.19	500	Pelagic fish	53.78	Tourism, agriculture	Ramos et al. (2009)
Dragonera 2°18'E 39°35'N	3	4500	Refuse tips	0.94	Textile, leather, tourism	Bermejo et al. (2009)
Mar Menor 0°42'W 37°43'N	0.165	1272	Refuse tips, brackish, freshwater and pelagic preys	1.57	Agriculture, greenhouses	Ramos et al. (2009)
Chafarinas 2°25'W 35°10'N	0.525	5700	Epipelagic fish	6.67	Agriculture	Arcos et al. (2002) and Pedrocchi et al. (2002)
Berlengas 9°30'E 39°24'N	0.788	23,000	Refuse tips	10.32	Industry, agriculture, boating	Moreno et al. (2010)
Cies 8°54'W 42°13'N	4.4	15,654	Most pelagic preys, but also benthic preys and refuse tips	2.54	Fishing, boating, chlor-alkali	Moreno et al. (2010)

At the same time that we analysed Hg in the 2009 eggs, a suite of metals also was analysed (data not presented here), including selenium (Se). The average recoveries ( $n = 3$ ) for Se in the reference materials (pig kidney, TORT-2 and DOLT-4) were 99% (RSD = 5.1%), 131% (RSD = 7.4%), and 111% (RSD = 3.3%) and the LOD was 0.05 mg/kg. The molar ratio of Hg to Se was calculated by dividing the calculated number of moles of Hg with those of selenium.

### Statistical Analysis

Hg concentrations in yellow-legged gulls from the various locations and for both sampling dates were log<sub>10</sub> transformed for analysis, and so geometric means and  $\pm$  geometric 95% confidence intervals (CI) are presented. The significance of differences between years and sites were analysed using a general linear model that included sampling year and site as random variables; differences between sites were assessed using Tukey post hoc tests. Data for Hg concentrations for yellow-legged gulls from Dragonera were not included in the analysis, because no eggs were sampled in 2016. The underlying assumptions of the model of homogeneity of variance (Barlett's test) and normality of residuals (Kolmogorov–Smirnov test) were met. A pairwise *t* test was


used to compare Hg concentrations between Audouin's and yellow-legged gull eggs from the Ebro Delta.

## Results and Discussion

### Temporal Variation in Hg in Yellow-Legged Gulls

The results of the Hg analyses in eggs collected from the various sites in 2009 and 2016 are presented in Fig. 2a. There was no significant difference in concentrations between the 2 years across the sites ( $F_{(1,26)} = 0.02$ ,  $P = 0.876$ ). This suggests that contamination remained relatively unchanged over the 7-year period. When colonies were grouped as Atlantic, NM, or SM, there were, again, no obvious differences in temporal trends, with the % change between 2009 and 2016 ranging between  $-0.1$  and 44%,  $-22$  and 7.5%, and  $-0.1$  and  $-12.9\%$ , respectively (Fig. 2a). The largest % changes were noticed in colonies with only one pooled sample in one or both sampling years; thus, in these cases data may be considered less robust. Changes in geometric mean concentration for sites with multiple pooled samples in either sampling year were only from  $-3.6$  to 7.5%. At the Ebro Delta, the removal of upstream contaminated sediments began in 2013 (Garcia-Tarrason et al. 2013). However, no marked

**Fig. 2** Geometric mean ( $\pm 95\%$  CI) concentration of total Hg in yellow-legged gull eggs collected from various colonies on the Iberian Peninsula. **a** Hg concentrations (mg/kg dw) in eggs collected in 2009 and 2016. **b** Data from 2009 and 2016 combined. Letters on top of bars indicates significant differences



changes in Hg concentrations in yellow-legged gull eggs (Fig. 2a) seem to be associated with dredging operations, perhaps reflecting the generalist diet of this species.

Overall, the lack of temporal changes in concentrations across the Iberian Peninsula contrasts with studies of Hg contamination in bird eggs elsewhere. Hg concentrations decreased in herring gull (*Larus argentatus*) eggs in the Great Lakes between 1970 and 2000, but temporal trends have since varied between locations, with increases in some cases and further declines in others (Blukacz-Richards et al. 2017). Braune et al. (2016) observed that residues in glaucous gull (*Larus hyperboreus*) eggs in the Canadian Arctic decreased between 1993 and 2003 but subsequently started to increase, although similar temporal trends were not always observed in other species measured

in the same study. Mercury concentrations also have been shown to have increased since the year 2000 in common loons (*Gavia immer*; Meyer et al. 2011) and in bald eagles (*Haliaeetus leucocephalus*; Pittman et al. 2011) in North America.

### Spatial Variation in Hg Concentrations in Yellow-Legged Gull Eggs

Hg concentrations in yellow-legged gull eggs ranged between 0.4 and 1.77 mg/kg dw, except in Columbretes, which contained up to 2.8 mg/kg dw. There were significant differences in concentrations between sites ( $F_{6,26} = 1877$ ,  $P < 0001$ ; Fig. 2b). They were highest in two of the three NM sites (i.e., Columbretes and Ebro Delta) and were



significantly higher than in the two SM sites (Mar Menor and Chafarina; Fig. 2b). Concentrations in the single 2009 pooled sample from Dragonera (not included in the statistical analysis) was 0.9 mg/kg dw, which is 1.5 and 3 times lower than the 2009 concentrations in the Ebro Delta and Columbretes Islands, respectively. Concentrations in Medes were significantly lower than those at the other two northern sites and were similar to those found in the more southern colonies (Fig. 2b) and also to those recorded in 1992 ( $0.52 \pm 0.29$   $\mu\text{g/g dw}$ ) at the same location (Sanpera et al. 1997). Although there is some evidence that Atlantic fauna accumulate less Hg compared with biota from the Mediterranean (Renzoni et al. 1998), the mean Hg residues in eggs from the two Atlantic colonies generally fell between the values recorded in eggs from the NM and SM colonies (Fig. 2b). The residues obtained in samples from Berlengas were similar to those ( $1.4$   $\mu\text{g/g dw}$ ) reported previously for yellow-legged gull eggs from the same area (Monteiro et al. 1999).

The spatial variation of Hg in yellow-legged gull eggs can be explained by the distribution of anthropogenic and natural sources of Hg. The NM Ebro Delta colony is close to industrialised and highly populated areas, and concentrations most likely reflect Hg inputs into the Ebro River from intensive agricultural and industrial activity within its watershed, including inputs from nearby chemical plants (Morera et al. 1997; Sanchiz et al. 2000; Sanpera et al. 2000). However, the highest mean Hg concentration was found in eggs from Columbretes Islands (Fig. 2b). Compared with the Ebro Delta, the Columbretes Islands are more isolated from major point sources of Hg, and so anthropogenic inputs alone cannot account for the relatively high concentrations at this colony, up to twofold higher than in the Ebro Delta. The Catalan-Balearic Sea in the NM carries cold northern waters from the Gulf of Lion southward along the continental slope in the Balearic Sea, and this current bifurcates at the northern end of the Eivissa Channel (Millot 1999), potentially transporting atmospheric Hg from the North towards Columbretes.

We expect that dietary differences between colonies may also contribute to the intercolony variation of Hg and in future work we plan to perform stable isotopes analyses to ascertain whether diet contributes to the observed differences in mercury concentrations. Yellow-legged gulls in the Iberian Peninsula feed in both marine and freshwater environments, prey on smaller seabirds, and also take the eggs of terrestrial and coastal birds (Ramos et al. 2009). Distance to the mainland is another factor influencing feeding ecology, with scavenging of waste from rubbish tips a more important part of the diet in birds close to the mainland (Ramos et al. 2013). Among the yellow-legged gull colonies studied, Columbretes Islands is the furthest site from the mainland ( $\approx 54$  km; Table 1), and therefore, gulls have a


more fish-based rather than waste-based diet (Ramos et al. 2009, 2013), which may account for the higher concentrations measured. Similarly, scavenging on waste tips also may explain why Hg levels in eggs from the Medes Islands were lower than those in eggs from the other NM colonies (Ebro Delta and Columbretes). Yellow-legged gulls from Medes are highly dependent on feeding at refuse tips (Bosch et al. 1994), which may reduce their exposure to Hg compared with feeding on more natural marine or freshwater-based diet (Leonzio et al. 1986).

There was little difference in Hg contamination between the two Atlantic colonies of Cies and Berlengas. The main anthropogenic source of Hg in Cies is the chlor-alkali plant situated on the south coast of the Ria of Pontevedra (Galicia) (Otero and Fernandez-Sanjurjo 2000). Whilst it may be expected that this may contribute to elevated contamination, Cies is an open Atlantic site and currents may dilute contaminant loadings into the food chain. Furthermore, refuse from rubbish tips is an important source in the diet of gulls from both Cies and Berlengas (Moreno et al. 2010). This commonality in diet sources may account for the overall similarity in the Hg levels between the colonies. It was also noticeable that the concentrations in eggs from Cies and Berlengas were similar to those from the (SM) colony at Mar Menor and higher than those from the Chafarinas (Fig. 2b). Given the potential of Atlantic oceanic currents to dilute inputs into the marine food chain, residues in the eggs of Atlantic gulls were expected to be lower than in the SM. However, a study of perfluorinated compounds in yellow-legged gulls eggs from the same colonies found no differences between these two water bodies (Vicente et al. 2012), and, as it has been suggested, temporary influx of Atlantic water into the SM may reduce contamination levels in the southern colonies.

### Comparison Between Hg Concentrations in Audouin's and Yellow-Legged Gull Eggs

In the three Ebro Delta subcolonies, concentrations were significantly higher for Audouin's than yellow-legged gulls eggs ( $3.41 \pm 0.52$  mg/kg dw (mean  $\pm$  SD);  $t_{(5)} = -14.64$ ;  $P < 0.001$ ; Fig. 3). Concentrations were of the same order of magnitude as those reported in other studies in Audouin's eggs from the Ebro Delta during 1992 ( $5.05 \pm 1.5$  mg/kg dw; Morera et al. 1997) and 1994 ( $5.0 \pm 1.54$  mg/kg dw; Sanpera et al. 2000), suggesting Hg accumulation has remained fairly unchanged for the past c. 25 years.

The Hg concentrations in Audouin's gull eggs from the Ebro delta are amongst the highest reported for seabird eggs anywhere (Ackerman et al. 2013). Audouin's gulls are specialised pelagic marine predators that feed mainly on clupeiform fish (Witt et al. 1981). During the breeding season, they feed preferentially on trawler discards (Oro and Ruiz



**Fig. 3** Mercury concentration in the eggs of yellow-legged gulls and Audouin's gull from three subcolonies of the Ebro Delta collected in 2009 (clear bars) and 2016 (dark bars)

1997), which contain both demersal and mesopelagic fish. A continuous release of Hg, from the chlor-alkali plant located 60 km upstream in the dissolve and particulate phase (Turull et al. 2017) would increase the Hg body burden of particular demersal fish species. Hg concentrations in demersal fish from the Ebro Delta can be more than double those in epipelagic fish (Arcos et al. 2002), and consequently Hg concentrations can sharply increase by a diet containing demersal fish (Arcos et al. 2002). Thus, it seems likely that the comparatively high Hg levels in Audouin's gull eggs around the Ebro delta are a result of higher contamination inputs in this area and to the birds feeding habits. Furthermore, it has been suggested (Henny et al. 2002) that some waterbird species are able to perform mercury demethylation when exposed to high concentrations. This does not appear to be the case with Audouin gulls, although we cannot be certain of this since we did not measure levels in adults.

### Toxicological Implications of Hg in Gull Eggs


Seabirds tend to accumulate high levels of Hg because aquatic conditions facilitate conversion of inorganic Hg into bioavailable MeHg, which bioaccumulates in fish and their predators (Monteiro and Furness 2001; Wiener et al. 2003). In this study, we analysed total Hg, because it has been shown to be a good proxy for the most toxic form of Hg (MeHg) in bird eggs. Ackerman et al. (2013) have reported that, across 22 different species of birds, 96% of the total Hg in eggs is MeHg.

Because the sensitivity of birds to Hg toxicity can differ widely among species (Ohlendorf and Heinz 2011), it is

difficult to compare egg Hg concentrations against known toxicity thresholds. Therefore, we have provided several toxicity benchmarks suggested to impair bird reproduction. Laboratory studies revealed that Hg can have deleterious effects in birds including decreased egg weight, embryo malformations, lower hatchability, decreased chick growth, and reduced survival of the young (Thompson 1996; Burger and Gochfeld 1997). For example, a 10% reduction in nest success corresponded with 0.11 mg/kg ww (0.55 mg/kg dw; Jackson et al. 2011) of mercury concentration in the egg. In fact, an egg Hg concentration of 0.6 mg/kg ww, equivalent to approximately 3 mg/kg dw (assuming eggs contain 80% water; Thompson 1996), has been suggested as a conservative generic value below which embryotoxic effects in most avian species are unlikely (Heinz and Hoffman 1998; Shore et al. 2011). Consistent with this, Evers et al. (2003) used Hg concentrations in eggs to estimate reproductive risk in loons (*Gavia immer*) and identified a “moderate risk category” of 2.82–6.10 mg/kg dw that reflected “elevated levels of environmental Hg that may be indicative of significant reproductive impairment in some individuals in some avian species, including loons”. However, the toxicological significance of Hg to seabirds in particular is difficult to interpret (Thompson 1996). An egg Hg concentration > 3.5 mg/kg ww (c. 17.5 mg/kg dw) has been suggested as an adverse effect level for common terns (Shore et al. 2011), but Vermeer et al. (1973) found that hatching success of herring gull eggs was unaffected by Hg concentrations in the first-laid egg between 2.3 and 15.8 mg/kg ww (c. 10–80 mg/kg dw). In the current study, none of the yellow-legged gull pooled samples from any colony exceeded 3 mg/kg dw. Thus, it seems unlikely that Hg contamination alone currently poses a significant reproductive risk to this species around the Iberian Peninsula. However, Hg concentrations in the Audouin's gull eggs ranged from 4.0 to 5.9 mg/kg dw. These concentrations are higher than the proposed generic adverse effect concentrations within the range of the “moderate risk” proposed for loons, generally greater than the adverse effect concentration proposed for common terns, but lower than the concentrations associated with no effect in herring gulls, as reported by Vermeer et al. (1973).

Furthermore, in a laboratory study where eggs were injected with methylmercury, it was shown that different bird species, including different species of gulls, have a wide range of sensitivities to this metal (Heinz et al. 2009). For example, based on dose–response curves and the median lethal concentration LC(50), Heinz et al. (2009) found that that the laughing gull (*Larus atricilla*) had relatively low sensitivity (LC(50)s  $\geq 1$   $\mu\text{g/g}$  Hg), whereas the herring gull (*Larus argentatus*) had medium sensitivity (LC(50)s > 0.25  $\mu\text{g/g}$  Hg but < 1  $\mu\text{g/g}$  Hg).

**Fig. 4** Relationship between mean Hg and Se concentrations in eggs from species collecting in 2009 by sampling sites. The stippled line represents the 1:1 molar ratio between Se and total Hg and the dotted line represents the relationship between Hg and Se for yellow-legged gulls (using all data points)



## Relationship Between Hg and Se

Selenium concentrations  $> 5.0$  mg/kg dw in eggs are in excess of the normal range, and 6.4 mg/kg dw approaches levels that induce reproductive impairment (Ohlendorf and Heinz 2011). Se concentrations in the present study ranged between 1.9 and 4.1 mg/kg dw, in the yellow-legged gulls and  $\sim 4$  in the Audouin and were below concentrations likely to be associated with adverse effects. However, taken into account all the measurements, there was a significant and positive correlation between yellow-legged egg concentrations of Hg and Se ( $r^2 = 0.796$ ,  $P < 0.001$ ; Fig. 4). Positive correlations also were found for eggs of the common loon (Scheuhammer et al. 2001), Forster's tern (*Sterna forsteri*), and black skimmer (*Rynchops niger*; King et al. 1991) but not previously for eggs of herring gull, Caspian tern (*Hydroprogne caspia*), or least tern (*Sternula antillarum*; Burger and Gochfeld 1995).

Molar ratios of Se to Hg differed substantially among species, with mean values for Audouin of 2.0 and for yellow-legged between 3.7 and 13.0. The mean molar ratios for both species were all greater than 1:1 (Fig. 4), with the lowest ratio value corresponding to Audouin gull in the Ebro Delta; for the yellow-legged gull the values decreased in the following order: Columbretes  $>$  Ebro Delta  $>$  Dragonera  $>$  Mar Menor  $>$  Cies  $>$  Berlengas  $>$  Medes  $>$  Chafarinas. Eggs with higher levels of Hg generally had lower molar ratios (Fig. 4), due to high Se concentrations. In fact, in sampling sites close to agricultural activities (such as the Ebro Delta), Se levels are high due to its considerable use in agriculture, as an additive to insecticides, fertilizers, and foliar sprays (Kabata-Pendias 2011).

## Conclusions

Mercury concentrations in yellow-legged gull eggs from the Iberian Peninsula appeared to have remained stable between 2009 and 2016 but tended to be higher in the NM (Ebro Delta and Columbretes) than in the SM or the Atlantic colonies. This spatial variation is likely to be the result of a combination of local pollution, oceanic currents, and birds feeding habits. Independently, concentrations of Hg and Se in the yellow-legged were below those associated with embryotoxicity, but there was a positive correlation between Hg and Se concentrations and these two elements can act synergistically to exert embryotoxicity.

Hg concentrations in eggs of Audouin's gulls from the Ebro Delta were three to four times higher than those of yellow-legged gulls from the same area. This may be the result of this species taking more fish than yellow-legged gulls do, including demersal fish discards from fishing boats. The Hg concentrations in Audouin's gull eggs are of a magnitude associated with embryotoxicity in other species. Mercury levels in Audouin's gull are more than ten times above the benchmark suggested to reduce nest success by 10%. Further investigation of the scale and effects of Hg and Se contamination in Audouin's gull eggs from the Ebro Delta is merited.

**Acknowledgements** From Spain, we acknowledge the Islas Atlánticas National Park, Refugio de Caza de las Islas Chafarinas, Illes Medes Natural Park, Delta de l'Ebre Natural Park, Columbretes Islands Natural Park, Montgrí, les Illes Medes i el Baix Ter Natural Park, that authorized our work. From Portugal, we acknowledge Maria Felisbina Quadrado and Sofia Batista from the Portuguese Environment Agency (APA), Department of Water Resources for the support; Lurdes Morais from Portuguese Protected Area of Berlengas-Portuguese Institute for the Conservation of Nature and Forests (ICNF) for the sampling permission. The Servei de Protecció i Gestió de la Fauna (Catalan Government) and Direcció General de Medi Natural i d'Avaluació Ambiental (Generalitat Valenciana) are acknowledged for sampling permission. National and Natural Park's staff are acknowledged for their help in



the gull eggs collection and transport. We are especially grateful to Javier Zapata (Chafarinas), José Antonio Fernández-Bouza and Vicente Piorro (Atlantic Islands), Juan Jiménez (Generalitat Valenciana), Miguel Ángel Monsalve Dolz (Equipo de Seguimiento de Fauna—VAERSA, Generalitat Valenciana), Francesc Vidal and Antoni Curcó (PN Delta de l'Ebre) for their enthusiasm, implication, logistic support, and valuable contributions to the project. Dr. Jacob González-Solis, Dr. Dani Oro, Javier Zapata, Cristina García and Marc Bosch are acknowledged for aiding in the sampling campaigns carried out in 2009. This study was financed by the Spanish Autonomous Organism of National Parks [Projects 2012/768 and 2009/038]. R. Ballesteros is acknowledged for his help in preparing the egg material for Hg analysis. From UK, this work was supported by the Natural Environment Research Council as part of the CEH programme delivering National Capability.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Ackerman JT, Herzog MP, Schwarzbach SE (2013) Methylmercury is the predominant form of mercury in bird eggs: a synthesis. *Environ Sci Technol* 47:2052–2060. <https://doi.org/10.1021/es304385y>
- Arcos JM, Ruiz X, Bearhop S, Furness RW (2002) Mercury levels in seabirds and their fish prey at the Ebro Delta (NW Mediterranean): the role of trawler discards as a source of contamination. *Mar Ecol Prog Ser* 232:281–290. <https://doi.org/10.3354/meps232281>
- Becker PH, Henning D, Furness RW (1994) Differences in mercury contamination and elimination during feather development in gull and tern broods. *Arch Environ Contam Toxicol* 27:162–167
- Bermejo A, Molina B, Cantos FJ, J M (eds) (2009) Gaviotas reidora, sombría y patiamarilla en España. Población en 2007–2009 y método de censo. SEO/BirdLife, Madrid
- Bishop CA, Koster MD, Check AA, Hussell DJT, Jock K (1995) Chlorinated hydrocarbons and mercury in the sediments, red-winged blackbirds (*Agelaius phoeniceus*) and tree swallows (*Tachycineta bicolor*) from wetlands in the Great lakes-St Lawrence river basin. *Environ Toxicol Chem* 14:491–501
- Blukacz-Richards EA, Visha A, Graham ML, McGoldrick DL, de Solla SR, Moore DJ, Arhonditsis GB (2017) Mercury levels in herring gulls and fish: 42 years of spatio-temporal trends in the Great Lakes. *Chemosphere* 172:476–487. <https://doi.org/10.1016/j.chemosphere>
- Boening DW (2000) Ecological effects, transport, and fate of mercury: a general review. *Chemosphere* 40:1335–1351. [https://doi.org/10.1016/s0045-6535\(99\)00283-0](https://doi.org/10.1016/s0045-6535(99)00283-0)
- Bosch M, Oro D, Ruiz X (1994) Dependence of yellow-legged gulls (*Larus cachinnans*) on food from human activity in two western Mediterranean colonies. *Avocetta* 18:135–139
- Braune BM, Gaston AJ, Mallory ML (2016) Temporal trends of mercury in eggs of five sympatrically breeding seabird species in the Canadian Arctic. *Environ Pollut* 214:124–131. <https://doi.org/10.1016/j.envpol.2016.04.006>
- Burger J, Gochfeld M (1995) Heavy-metal and selenium concentrations in eggs of herring-gulls (*larus-argentatus*)—temporal differences from 1989 to 1994. *Arch Environ Contam Toxicol* 29:192–197
- Burger J, Gochfeld M (1997) Risk, mercury levels, and birds: relating adverse laboratory effects to field biomonitoring. *Environ Res* 75:160–172
- Carrasco L, Diez S, Soto DX, Catalan J, Bayona JM (2008) Assessment of mercury and methylmercury pollution with zebra mussel (*Dreissena polymorpha*) in the Ebro River (NE Spain) impacted by industrial hazardous dumps. *Sci Total Environ* 407:178–184. <https://doi.org/10.1016/j.scitotenv.2008.07.031>
- Carrasco L, Barata C, Garcia-Berthou E, Tobias A, Bayona JM, Diez S (2011a) Patterns of mercury and methylmercury bioaccumulation in fish species downstream of a long-term mercury-contaminated site in the lower Ebro River (NE Spain). *Chemosphere* 84:1642–1649. <https://doi.org/10.1016/j.chemosphere.2011.05.022>
- Carrasco L, Benjam L, Benito J, Bayona JM, Diez S (2011b) Methylmercury levels and bioaccumulation in the aquatic food web of a highly mercury-contaminated reservoir. *Environ Int* 37:1213–1218. <https://doi.org/10.1016/j.envint.2011.05.004>
- Champoux L, Boily M (2017) Temporal trends of mercury and organohalogen contaminants in great blue heron eggs from the St. Lawrence River, Quebec, Canada, 1991–2011, and relationships with tracers of feeding ecology. *Sci Total Environ* 609:1270–1285. <https://doi.org/10.1016/j.scitotenv.2017.07.223>
- Cusack LK, Eagles-Smith C, Harding AK, Kile M, Stone D (2017) Selenium: mercury molar ratios in freshwater fish in the Columbia River Basin: potential applications for specific fish consumption advisories. *Biol Trace Elem Res* 178:136–146. <https://doi.org/10.1007/s12011-016-0907-9>
- Diez S (2009) Human health effects of methylmercury exposure. In: Whitacre DM (ed) Reviews of environmental contamination and toxicology, vol 198. Springer, New York, pp 111–132
- Dittmann T et al (2012) Large-scale spatial pollution patterns around the North Sea indicated by coastal bird eggs within an EcoQO programme. *Environ Sci Pollut Res* 19:4060–4072. <https://doi.org/10.1007/s11356-012-1070-2>
- Driscoll CT, Mason RP, Chan HM, Jacob DJ, Pirrone N (2013) Mercury as a global pollutant: sources, pathways, and effects. *Environ Sci Technol* 47:4967–4983. <https://doi.org/10.1021/es305071v>
- Evers DC, Taylor KM, Major A, Taylor RJ, Poppenga RH, Scheuhammer AM (2003) Common loon eggs as indicators of methylmercury availability in North America. *Ecotoxicology* 12:69–81. <https://doi.org/10.1023/a:1022593030009>
- Fernandez-Chacon A et al (2013) When to stay, when to disperse and where to go: survival and dispersal patterns in a spatially structured seabird population. *Ecography* 36:1117–1126. <https://doi.org/10.1111/j.1600-0587.2013.00246.x>
- Furness RW (1993) Birds as monitors of pollutants. In: Furness RW, Greenwood JJD (eds) Birds as monitors of environmental change. Chapman & Hall, London, pp 86–143
- Garcia-Ordiales E et al (2018) Occurrence and speciation of arsenic and mercury in estuarine sediments affected by mining activities (Asturias, northern Spain). *Chemosphere* 198:281–289. <https://doi.org/10.1016/j.chemosphere.2018.01.146>
- Garcia-Sanchez A, Murcigo A, Alvarez-Ayuso E, Santa-Regina I, Rodríguez-González MA (2009) Mercury in soils and plants in an abandoned cinnabar mining area (SW Spain). *J Hazard Mater* 168:1319–1324. <https://doi.org/10.1016/j.jhazmat.2009.03.009>
- Garcia-Tarrason M, Pacho S, Jover L, Sanpera C (2013) Anthropogenic input of heavy metals in two Audouin's gull breeding colonies. *Mar Pollut Bull* 74:285–290. <https://doi.org/10.1016/j.marpolbul.2013.06.043>
- Genovart M, Oro D, Tenan S (2018) Immature survival, fecundity and density-dependence, drive global population dynamics in a long-lived bird. *Ecology* 99:2823–2832. <https://doi.org/10.1002/ecy.2515>
- Giesy JP, Ludwig JP, Tillitt DE (1994) Deformities in birds of the Great Lakes region: assessing causality. *Environ Sci Technol* 28:128–135

- Grassman KA, Scanlon PF, Fox GA (1998) Reproductive and physiological effects of environmental contaminants in fish-eating birds of the Great Lakes: a review of historical trends. *Environ Monit Assess* 53:117–145
- Guitart R, Mateo R, Sanpera C, Hernandez-Matias A, Ruiz X (2003) Mercury and selenium levels in eggs of common terns (*Sterna hirundo*) from two breeding colonies in the Ebro Delta, Spain. *Bull Environ Contam Toxicol* 70:71–77. <https://doi.org/10.1007/s00128-002-0157-8>
- Heinz GH, Hoffman DJ (1998) Methylmercury chloride and selenomethionine interactions on health and reproduction in mallards. *Environ Toxicol Chem* 17:139–145. [https://doi.org/10.1897/1551-5028\(1998\)017%3c0139:mcasio%3e2.3.co;2](https://doi.org/10.1897/1551-5028(1998)017%3c0139:mcasio%3e2.3.co;2)
- Heinz GH et al (2009) Species differences in the sensitivity of avian embryos to methylmercury. *Arch Environ Contam Toxicol* 56:129–138
- Henny CJ, Hill EF, Hoffman DJ, Spalding MG, Grove RA (2002) Nineteenth century mercury: hazard to wading birds and cormorants of the Carson River, Nevada. *Ecotoxicology* 11:213–231. <https://doi.org/10.1023/a:1016327602656>
- Jackson AK et al (2011) Mercury exposure affects the reproductive success of a free-living terrestrial songbird, the carolina wren (*Thryothorus ludovicianus*). *Auk* 128:759–769. <https://doi.org/10.1525/auk.2011.11106>
- Kabata-Pendias A (2011) Trace elements in soils and plants, 4th edn. CRC Press, Boca Raton, p 505
- Kenow KP, Gutreuter S, Hines RK, Meyer MW, Fournier F, Karasov WH (2003) Effects of methyl mercury exposure on the growth of juvenile common loons. *Ecotoxicology* 12:171–182. <https://doi.org/10.1023/a:1022598525891>
- King KA, Custer TW, Quinn JS (1991) Effects of mercury, selenium, and organochlorine contaminants on reproduction of forster terns and black skimmers nesting in a contaminated texas bay. *Arch Environ Contam Toxicol* 20:32–40
- Leonzio C, Fossi C, Focardi S (1986) Lead, mercury, cadmium and selenium in 2 species of gull feeding on inland dumps, and in marine areas. *Sci Total Environ* 57:121–127. [https://doi.org/10.1016/0048-9697\(86\)90017-3](https://doi.org/10.1016/0048-9697(86)90017-3)
- Meyer MW, Rasmussen PW, Watras CJ, Fovold BM, Kenow KP (2011) Bi-phasic trends in mercury concentrations in blood of Wisconsin common loons during 1992–2010. *Ecotoxicology* 20:1659–1668. <https://doi.org/10.1007/s10646-011-0759-1>
- Millot C (1987) Circulation in the western mediterranean-sea. *Oceanol Acta* 10:143–149
- Millot C (1999) Circulation in the western Mediterranean Sea. *J Mar Syst* 20:423–442. [https://doi.org/10.1016/s0924-7963\(98\)00078-5](https://doi.org/10.1016/s0924-7963(98)00078-5)
- Molina JA, Oyarzun R, Esbri JM, Higuera P (2006) Mercury accumulation in soils and plants in the Almaden mining district, Spain: one of the most contaminated sites on Earth. *Environ Geochem Health* 28:487–498. <https://doi.org/10.1007/s10653-006-9058-9>
- Monteiro LR, Furness RW (2001) Kinetics, dose-response, and excretion of methylmercury in free-living adult Cory's shearwaters. *Environ Sci Technol* 35:739–746
- Monteiro LR, Granadeiro JP, Furness RW, Oliveria P (1999) Contemporary patterns of mercury contamination in the Portuguese Atlantic inferred from mercury concentrations in seabird tissues. *Mar Environ Res* 47:137–156. [https://doi.org/10.1016/s0141-1136\(98\)00110-x](https://doi.org/10.1016/s0141-1136(98)00110-x)
- Moreno R, Jover L, Munilla I, Velando A, Sanpera C (2010) A three-isotope approach to disentangling the diet of a generalist consumer: the yellow-legged gull in northwest Spain. *Mar Biol* 157:545–553. <https://doi.org/10.1007/s00227-009-1340-9>
- Morera M, Sanpera C, Crespo S, Jover L, Ruiz X (1997) Inter- and intraclutch variability in heavy metals and selenium levels in Audouin's gull eggs from the Ebro Delta, Spain. *Arch Environ Contam Toxicol* 33:71–75
- Ohlendorf HM, Heinz GH (2011) Selenium in birds. In: Beyer WN, Meador JP (eds) *Environmental contaminants in biota—interpreting tissue concentrations*, 2nd edn. CRC Press, Boca Raton, pp 669–701
- Oro D, Ruiz X (1997) Exploitation of trawler discards by breeding seabirds in the north-western Mediterranean: differences between the Ebro Delta and the Balearic Islands areas. *ICES J Mar Sci* 54:695–707. <https://doi.org/10.1006/jmsc.1997.0246>
- Oro D, Perez-Rodriguez A, Martinez-Vilalta A, Bertolero A, Vidal F, Genovart M (2009) Interference competition in a threatened seabird community: a paradox for a successful conservation. *Biol Conserv* 142:1830–1835. <https://doi.org/10.1016/j.bioco.2009.03.023>
- OSPAR Commission (2007) Background document on the EcoQO on mercury and organohalogens in seabird eggs. <https://www.ospar.org/documents?v=7089>. Accessed June 2018
- Otero XL, Fernandez-Sanjurjo MJ (2000) Mercury in faeces and feathers of yellow-legged gulls (*Larus cachinnans*) and in soils from their breeding sites (Cies Islands-NW Spain) in the vicinity of a chlor-alkali plant. *Fresenius Environ Bull* 9:56–63
- Pedrocchi V, Oro D, Gonzalez-Solis J, Ruiz X, Jover L (2002) Differences in diet between the two largest breeding colonies of Audouin's gulls: the effects of fishery activities. *Scientia Marina* 66:313–320
- Pereira MG, Walker LA, Best J, Shore RF (2009) Long-term trends in mercury and PCB congener concentrations in gannet (*Morus bassanus*) eggs in Britain. *Environ Pollut* 157:155–163. <https://doi.org/10.1016/j.envpol.2008.07.019>
- Peterson SH, Ackerman JT, Eagles-Smith CA, Hartman CA, Herzog MP (2017) A critical evaluation of the utility of eggshells for estimating mercury concentrations in avian eggs. *Environ Toxicol Chem* 36:2417–2427. <https://doi.org/10.1002/etc.3777>
- Pittman HT, Bowerman WW, Grim LH, Grubb TG, Bridges WC (2011) Using nestling feathers to assess spatial and temporal concentrations of mercury in bald eagles at Voyageurs National Park, Minnesota, USA. *Ecotoxicology* 20:1626–1635. <https://doi.org/10.1007/s10646-011-0736-8>
- Ralston NVC, Raymond LJ (2013) Selenium status and intake influences mercury exposure risk assessments. In: Banuelos GS, Lin Z-Q, Yin X (eds) *Selenium in the environment and human health*. CRC, London, pp 203–205
- Ramos R, Ramirez F, Sanpera C, Jover L, Ruiz X (2009) Diet of yellow-legged gull (*Larus michahellis*) chicks along the Spanish Western Mediterranean coast: the relevance of refuse dumps. *J Ornithol* 150:265–272. <https://doi.org/10.1007/s10336-008-0346-2>
- Ramos R, Ramirez F, Jover L (2013) Trophodynamics of inorganic pollutants in a wide-range feeder: the relevance of dietary inputs and biomagnification in the yellow-legged gull (*Larus michahellis*). *Environ Pollut* 172:235–242. <https://doi.org/10.1016/j.envpol.2012.09.014>
- Renzoni A, Zino F, Franchi E (1998) Mercury levels along the food chain and risk for exposed populations. *Environ Res* 77:68–72. <https://doi.org/10.1006/enrs.1998.3832>
- Roig N, Nadal M, Sierra J, Ginebreda A, Schuhmacher M, Domingo JL (2011) Novel approach for assessing heavy metal pollution and ecotoxicological status of rivers by means of passive sampling methods. *Environ Int* 37:671–677. <https://doi.org/10.1016/j.envint.2011.01.007>
- Rubio B, Nombela MA, Vilas F (2000) Geochemistry of major and trace elements in sediments of the Ria de Vigo (NW Spain): an assessment of metal pollution. *Mar Pollut Bull* 40:968–980. [https://doi.org/10.1016/s0025-326x\(00\)00039-4](https://doi.org/10.1016/s0025-326x(00)00039-4)
- Sanchez-Avila J, Bonet J, Velasco G, Lacorte S (2009) Determination and occurrence of phthalates, alkylphenols, bisphenol A, PBDEs, PCBs and PAHs in an industrial sewage grid discharging

- to a Municipal Wastewater Treatment Plant. *Sci Total Environ* 407:4157–4167. <https://doi.org/10.1016/j.scitotenv.2009.03.016>
- Sanchiz C, Garcia-Carrascosa AM, Pastor A (2000) Heavy metal contents in soft-bottom marine macrophytes and sediments along the Mediterranean coast of Spain. *Marine Ecology-Pubblicazioni Della Stazione Zoologica Di Napoli I* 21:1–16. <https://doi.org/10.1046/j.1439-0485.2000.00642.x>
- Sanpera C, Morera M, Crespo S, Ruiz X, Jover L (1997) Trace elements in clutches of yellow-legged Gulls, *Larus cachinnans*, from the Medes Islands, Spain. *Bull Environ Contam Toxicol* 59:757–762
- Sanpera C, Morera M, Ruiz X, Jover L (2000) Variability of mercury and selenium levels in clutches of audouin's gulls (*Larus audouinii*) breeding at the Chafarinas Islands, southwest Mediterranean. *Arch Environ Contam Toxicol* 39:119–123
- Scheuhammer AM, Perrault JA, Bond DE (2001) Mercury, methylmercury, and selenium concentrations in eggs of common loons (*Gavia immer*) from Canada. *Environ Monit Assess* 72:79–94. <https://doi.org/10.1023/a:1011911805216>
- Shore RP, Pereira MG, Walker LA, Thompson DR (2011) Mercury in nonmarine birds and mammals. In: Beyer WM, Meador JP (eds) *Environmental contaminants in biota: interpreting tissue concentrations*, 2nd edn. CRC Press, Taylor & Francis Group, London, pp 609–627
- Spiller HA (2018) Rethinking mercury: the role of selenium in the pathophysiology of mercury toxicity. *Clin Toxicol* 56:313–326. <https://doi.org/10.1080/15563650.2017.1400555>
- Thompson DR (1996) Mercury in birds and terrestrial mammals. In: Beyer WN, Heinz GH, Redmon-Norwood AW (eds) *Environmental contaminants in wildlife: interpreting tissue concentrations*. CRC Lewis Publishers, Boca Raton, pp 341–356
- Turull M, Grmanova G, Dago A, Arino C, Diez S, Diaz-Cruz JM, Esteban M (2017) Phytochelatin synthesis in response to Hg uptake in aquatic plants near a chlor-alkali factory. *Chemosphere* 176:74–80. <https://doi.org/10.1016/j.chemosphere.2017.02.092>
- UN Environment. Global monitoring plan on persistent organic pollutants protocol for the sampling and pre-treatment of national samples within the UNEP/GEF projects to support the global monitoring plan of POPs 2016–2019. <http://wedocs.unep.org/bitstream/handle/20.500.11822/21023/SOP-National%20Sample%202017%20En.pdf?sequence=1&isAllowed=y>. Accessed June 2018
- Vermeer K, Armstrong FA, Hatch DRM (1973) Mercury in aquatic birds at clay-lake, western-ontario. *J Wildl Manag* 37:58–61. <https://doi.org/10.2307/3799738>
- Vicente J, Bertolero A, Meyer J, Viana P, Lacorte S (2012) Distribution of perfluorinated compounds in yellow-legged gull eggs (*Larus michahellis*) from the Iberian Peninsula. *Sci Total Environ* 416:468–475. <https://doi.org/10.1016/j.scitotenv.2011.11.085>
- Walker C, Livingstone DR (1992) *Persistent pollutants in marine ecosystems*. Pergamon Press, Oxford
- Wiener JG, Krabbenhoft DP, Heinz GH, Scheuhammer AM (2003) *Ecotoxicology of mercury*. In: Hoffman DJ, Rattner BA, Burton GA Jr, Cairns J Jr (eds) *The handbook of ecotoxicology*. Lewis Publishers, New York, pp 409–463
- Witt H-HC et al (1981) Comparative feeding ecology of Audouin's gull *Larus audouinii* and the herring gull *L. argentatus* in the Mediterranean. *Ibis* 1:519–526
- Yamashita N, Kannan K, Taniyasu S, Horii Y, Petrick G, Gamo T (2005) A global survey of perfluorinated acids in oceans. *Mar Pollut Bull* 51:658–668. <https://doi.org/10.1016/j.marpolbul.2005.04.026>