



Invariant Measures for the Stochastic One-Dimensional Compressible Navier–Stokes Equations

Michele Coti Zelati¹ · Nathan Glatt-Holtz² · Konstantina Trivisa³

Published online: 16 July 2019
© The Author(s) 2019

Abstract

We investigate the long-time behavior of solutions to a stochastically forced one-dimensional Navier–Stokes system, describing the motion of a compressible viscous fluid, in the case of linear pressure law. We prove existence of an invariant measure for the Markov process generated by strong solutions. We overcome the difficulties of working with non-Feller Markov semigroups on non-complete metric spaces by generalizing the classical Krylov–Bogoliubov method, and by providing suitable polynomial and exponential moment bounds on the solution, together with pathwise estimates.

Keywords Navier–Stokes system · Compressible fluid · Stochastic perturbation · Invariant measure

Mathematics Subject Classification 35R60 · 37L40 · 35Q35 · 60H30

1 Introduction

We consider the question of existence of an invariant measure for the stochastically forced Navier–Stokes equations of compressible fluid flows in one space dimension. Formulated on the unit interval $(0, 1)$, the equations read

✉ Michele Coti Zelati
m.coti-zelati@imperial.ac.uk

Nathan Glatt-Holtz
negh@tulane.edu

Konstantina Trivisa
trivisa@math.umd.edu

¹ Department of Mathematics, Imperial College London, London SW7 2AZ, UK

² Department of Mathematics, Tulane University, New Orleans, LA 70118, USA

³ Department of Mathematics, University of Maryland, College Park, MD 20742, USA

$$\rho_t + (\rho u)_x = 0 \quad (1.1)$$

$$d(\rho u) + \left(\rho u^2 + A^2 \rho \right)_x dt = u_{xx} dt + \rho \sigma dW, \quad (1.2)$$

where $\rho = \rho(t, x)$ is the fluid density and $u = u(t, x)$ is the fluid velocity. System (1.1)–(1.2) is appropriately written in dimensionless form, and is supplemented by the initial conditions

$$\rho(0, x) = \rho_0(x), \quad u(0, x) = u_0(x), \quad (1.3)$$

where ρ_0, u_0 are assigned functions defined on the interval $[0, 1]$, with the (strictly positive) initial density having normalized mass

$$\int_0^1 \rho_0(x) dx = 1.$$

In view of mass conservation, we obviously have

$$\int_0^1 \rho(t, x) dx = 1, \quad \forall t \geq 0. \quad (1.4)$$

Moreover, we assume homogeneous Dirichlet boundary conditions for the velocity u , namely

$$u(t, 0) = u(t, 1) = 0, \quad \forall t \geq 0. \quad (1.5)$$

Above, the driving noise is given by a collection of independent white noise processes, suitably colored in space (see (2.1) below), while $A > 0$ is a dimensionless parameter, inversely proportional to the Mach number. When considering strong (in the deterministic sense) solutions to (1.1)–(1.5), the natural phase space for our system is

$$\mathcal{X} = \left\{ (\rho, u) \in H^1(0, 1) \times H_0^1(0, 1) : \int_0^1 \rho(x) dx = 1, \rho > 0 \right\}. \quad (1.6)$$

The main result of this article is the following.

Theorem 1.1 *For every fixed $A > 0$, the Markov semigroup $\{\mathcal{P}_t\}_{t \geq 0}$ associated to (1.1)–(1.5) possesses an invariant probability measure $\mu_A \in \mathfrak{P}(\mathcal{X}_{L^2})$. Furthermore,*

$$\int_{\mathcal{X}} \left[A^2 \|(\log \rho)_x\|_{L^2}^2 + \|u_x\|_{L^2}^2 \right] d\mu_A(\rho, u) \leq \|\sigma\|_{L^\infty}^2. \quad (1.7)$$

Here, \mathcal{X}_{L^2} denotes the set \mathcal{X} in (1.6), endowed with the $L^2 \times L^2$ metric, and $\|\sigma\|_{L^\infty}$ is given by (2.2) below.

The existence of statistically stationary states to randomly driven systems in fluid dynamics is of basic importance from both mathematical and experimental viewpoints. On the one hand, the existence of an invariant measure provides information on the generic long-time behavior of the system. On the other hand, under ergodicity assumptions, it provides a link between experimental observations (for example, in turbulence theory) and theoretical predictions.

In the context of fluid dynamics, a satisfactory theory of invariant measures has been developed for two-dimensional, incompressible flows [6,7,14,22,26,27,36,37,41,42,52]. In three dimensions, we mention the works [11,12,23], and especially [21], in which solutions which are strictly stationary stochastic processes are constructed, but the concept of invariant measure as a steady state is not well defined due to the absence of the Markov property. In this sense, the situation for multi-dimensional compressible flows is similar. To the best of our knowledge, the only relevant articles are [24], in which an invariant measure is constructed for an approximation of the Lagrangian version of (1.1)–(1.2) and for a different pressure law, and the more recent [4], in which statistically stationary solutions (but no invariant measure) are constructed. There are, however, results for both the dissipative and non-dissipative one-dimensional Burgers equation [10,13,25,44,51].

In this article, we prove existence of an invariant measure for the compressible Navier–Stokes system. The main advantage in working in one space dimension is that the equations are globally well-posed and can be solved pathwise, leaning on various works in the deterministic setting [28,29,31–34,43,46]. For the global solvability in the stochastic case, we mention [48,49], valid in one space dimension, and [2,5,45,47,50] for the multi-dimensional case. For the deterministic set-up, we refer to the classical references [16,38], and to [18,19] for long-time behavior results.

In turn, (1.1)–(1.5) generates a proper Markov semigroup, and the concept of an invariant measure can be defined in a standard manner. However, there are several obstructions on the path towards the proof of Theorem 1.1. The main ingredients of our approach and contributions to the existing theory on invariant measures can be summarized as follows:

- *An L^2 -based continuity result via the derivation of polynomial and exponential moment bounds* As it is clear from its statement, the topology on \mathcal{X} plays an essential role. The metric space \mathcal{X}_{L^2} is not complete (hence not Polish), due to both the open condition $\rho > 0$ and the fact that the Sobolev space H^1 is not a closed subspace of L^2 . This choice is dictated by the available continuous dependence estimates (cf. Theorem 2.7), which make use of the so-called relative entropy functional and essentially provide an L^2 -based continuity result. Such a result crucially depends on polynomial and exponential moment bounds of the solution to (1.1)–(1.5), which are carried out in detail in Sect. 2.1, and on careful pathwise estimates (see Sect. 2.2).
- *Introduction of a (larger) class of functions on \mathcal{X}_{L^2} which is invariant under the Markov semigroup \mathcal{P}_t* Closely related to the issue mentioned above, a second difficulty lies in the fact that the Markov semigroup associated to (1.1)–(1.5) is not known to be Feller (cf. Sect. 3.1), namely \mathcal{P}_t may not map $C_b(\mathcal{X}_{L^2})$ (the real-

valued, continuous bounded functions on \mathcal{X}_{L^2}) into itself. This is again attributable to the mismatch between the set \mathcal{X} and the L^2 topology, and causes problems when trying to apply the classical Krylov–Bogoliubov procedure to prove existence and, in particular, invariance of a probability measure constructed as a subsequential limit of time-averaged measures. We circumvent this issue by defining a class of functions on \mathcal{X}_{L^2} that is slightly larger than $C_b(\mathcal{X}_{L^2})$, which is invariant under the Markov semigroup \mathcal{P}_t , and still well-behaved in the above limiting procedure.

- *Derivation of lower and upper bounds of the density with the correct time-averaged growth* Lastly, the lack of instantaneous smoothing in the density equation (1.1) constitutes an obstacle towards compactness estimates for time-averaged measures. However, an energy structure already exploited in [32,43] allows to obtain a dissipation term involving $\|(\log \rho)_x\|_{L^2}$, thanks to the linear pressure law adopted here. Consequently, upper and lower bounds on the density (which, in 1D, can be proven with more general pressure laws) have the correct time-averaged growth and provide suitable tightness estimates for time-averaged measures (cf. Sect. 3.2).
- The present article is, according to our knowledge, the first that presents rigorous results on the existence of invariant measures for compressible flows.

Remark 1.2 (The low Mach number limit) One may be tempted to study the behavior of the measures μ_A as $A \rightarrow \infty$, in the spirit of compressible-incompressible limits [3,15,17,39,40]. From (1.7), it is clear that, as $A \rightarrow \infty$, the density component of the measures concentrates on sets such that $\log \rho = 0$, namely, $\rho = 1$. However, (1.1) implies that $u_x = 0$, and due to the homogeneous boundary conditions (1.5), we deduce that also $u = 0$. This is inconsistent with the second equation, as long as $\sigma \neq 0$. By replacing σ with $A^{-\eta}\sigma$, for any $\eta > 0$, it is easily seen from (1.7) that $\mu_A \rightarrow \delta_{(\rho=1, u=0)}$ as $A \rightarrow \infty$, essentially describing rigid body motion.

Extensions and Further Developments

Theorem 1.1 is in fact true for pressure laws of the type

$$p(\rho) \sim \rho, \quad \text{as } \rho \rightarrow 0,$$

and any polynomial growth as $\rho \rightarrow \infty$. The case of pure power law $p(\rho) = \rho^\gamma$, for $\gamma > 1$, remains open. Another important question is the uniqueness (and hence ergodicity) and the attracting properties of the measure μ_A . It is worth to point out that the problem is highly degenerate, since the noise only acts on one component of the phase space. We conjecture that uniqueness holds, since the unforced system evolves (at a very slow rate) towards the unique steady state $(\rho, u) = (1, 0)$.

Outline of the Article

Section 2 is dedicated to the properties of solutions to the compressible Navier–Stokes equations. Energy estimates and moment bounds are proven in Sect. 2.1, pathwise estimates in Sect. 2.2, while uniqueness and continuous dependence on data are discussed in Sects. 2.3 and 2.4, respectively. In Sect. 3 we prove Theorem 1.1, setting up the

Markovian framework and discussing existence of invariant measure for non-Feller Markov processes on non-complete metric spaces.

Notation and Conventions

Throughout the paper, c will denote a *generic* positive constant *independent* of A , whose value may change even in the same line of a certain equation. In the same way, C_1, C_2, \dots denote specific large deterministic constants. We will denote by \mathcal{X}_{L^2} the set \mathcal{X} endowed with $L^2 \times L^2$ metric. Notice that \mathcal{X}_{L^2} is clearly a metric space. When we want to indicate the set \mathcal{X} endowed with the $H^1 \times H^1$ metric, we write \mathcal{X}_{H^1} . By $\mathcal{B}(\mathcal{X}_{L^2})$, we denote the family of Borel subsets of \mathcal{X}_{L^2} . With the symbol $M_b(\mathcal{X}_{L^2})$ (resp. $C_b(\mathcal{X}_{L^2})$) we refer to the set of all real valued bounded measurable (resp. bounded continuous) functions on \mathcal{X}_{L^2} . Finally, $\mathfrak{P}(\mathcal{X}_{L^2})$ is the set of all probability measures on \mathcal{X}_{L^2} .

2 Existence and Uniqueness of Solutions

In view of the particular form of the stochastic forcing in (1.2), the proof of existence and uniqueness of pathwise solutions borrows many ideas from the deterministic case. Fix a stochastic basis

$$S = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W).$$

We write the noise term in (1.2) as

$$\sigma dW = \sum_{k=1}^{\infty} \sigma_k(x) dW^k(t). \tag{2.1}$$

The sequence $W(t) = \{W^k(t)\}_{k \in \mathbb{N}}$ consists of independent copies of the standard one-dimensional Wiener process (Brownian motion). As such, for each k , $dW^k(t)$ is formally a white noise which, in particular, is stationary in time. Throughout the article, we will assume

$$\|\sigma_{xx}\|_{L^2}^2 := \int_0^1 \sum_{\ell=1}^{\infty} |(\sigma_{\ell})_{xx}(x)|^2 dx < \infty, \quad \sigma_{\ell} \in H^2 \cap H_0^1.$$

In particular, this implies

$$\|\sigma\|_{L^{\infty}}^2 := \sup_{x \in (0,1)} \sum_{\ell=1}^{\infty} |\sigma_{\ell}(x)|^2 < \infty. \tag{2.2}$$

By considering the change of variable $\tilde{u} = u - \sigma W$, a pathwise approach for the existence of solutions (even in the multidimensional case) has been carried out in [20] (see also [47–49] and the more recent [5] for local existence of strong solutions). In

the deterministic setting and in one space dimension, the local existence of strong solutions has been proved in [46], while their global existence can be found in [43], in the case of pressure law of the type ρ^γ , for $\gamma > 1$. That this result can be extended to our case is a consequence of the estimates provided in the work [32] on (deterministic) weak solutions and their regularization properties. We summarize these observations in the theorem below.

Theorem 2.1 *Fix a stochastic basis $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W)$. For any $(\rho_0, u_0) \in \mathcal{X}$, there exists a unique $(\rho(t; \rho_0), u(t; u_0))$ satisfying (1.1)–(1.5), in the time integrated sense and with the regularity*

$$\rho \in L_{loc}^\infty(0, \infty; H^1), \quad u \in L_{loc}^\infty(0, \infty; H_0^1) \cap L_{loc}^2(0, \infty; H^2),$$

almost surely. Moreover, the map $t \mapsto (\rho(t; \rho_0), u(t; u_0))$ is weakly continuous, and for every $t \geq 0$,

$$\rho(t) \in L^\infty \quad \text{and} \quad \frac{1}{\rho(t)} \in L^\infty,$$

almost surely.

The bounds in the Theorem 2.1 (see also Lemma 2.2) for the density show that neither vacuum states nor concentration states can occur, no matter how large the initial datum is. This is one of several important differences between the Navier–Stokes equations and the inviscid Euler equations, for which vacuum states may in fact occur for large initial data and for certain equations of state (cf. [8,9]). It is also relevant in this regard that solutions of the Navier–Stokes equations show certain instabilities when vacuum states are allowed (cf. Hoff and Serre [30]).

As far as the regularity of solutions is concerned, we need quantitative estimates and moment bounds that are new to the best of our knowledge. These estimates are contained in the Propositions 2.3 and 2.4 below, and, in particular, agree with the regularity expressed in the theorem above.

2.1 Energy Estimates

For a pair $(\rho, u) \in \mathcal{X}$, we define the entropy function in the classical way as

$$\mathcal{H}(\rho, u) = \int_0^1 \left(\frac{1}{2} \rho u^2 + A^2 \rho \log \rho \right) dx.$$

In the one-dimensional setting, it turns out the strong solutions are also well-behaved with respect to the modified energy functional

$$\mathcal{E}(\rho, u) = \mathcal{H}(\rho, u) + \frac{1}{2} \int_0^1 \left(\frac{\rho_x u}{\rho} + \frac{1}{2} \frac{\rho_x^2}{\rho^3} \right) dx.$$

Several quantities are controlled by \mathcal{E} . In particular, uniform estimates on \mathcal{E} entail lower bounds on ρ .

Lemma 2.2 *Assume that $(\rho, u) \in \mathcal{X}$ are such that $\mathcal{E}(\rho, u) < \infty$. Then*

$$e^{-[8\mathcal{E}(\rho,u)]^{1/2}} \leq \rho(x) \leq e^{[8\mathcal{E}(\rho,u)]^{1/2}}, \quad \forall x \in [0, 1], \tag{2.3}$$

and

$$\|\rho_x\|_{L^2}^2 \leq 8\mathcal{E}(\rho, u)e^{3[8\mathcal{E}(\rho,u)]^{1/2}}. \tag{2.4}$$

Moreover,

$$\|u\|_{L^2}^2 \leq 2\mathcal{H}(\rho, u)e^{[8\mathcal{E}(\rho,u)]^{1/2}}. \tag{2.5}$$

Proof As a preliminary observation we note that (pointwise)

$$\frac{\rho_x u}{\rho} \leq \rho u^2 + \frac{1}{4} \frac{\rho_x^2}{\rho^3}. \tag{2.6}$$

Moreover, since $\rho \in H^1$ and

$$\int_0^1 \rho \, dx = 1, \tag{2.7}$$

we have

$$\int_0^1 \rho \log \rho \, dx = \int_0^1 [\rho \log \rho - \rho + 1] \, dx \geq 0.$$

Therefore

$$\mathcal{E}(\rho, u) = \int_0^1 \left(\frac{1}{2} \rho u^2 + A^2 \rho \log \rho + \frac{1}{2} \frac{\rho_x u}{\rho} + \frac{1}{4} \frac{\rho_x^2}{\rho^3} \right) dx \geq \frac{1}{8} \int_0^1 \frac{\rho_x^2}{\rho^3} dx. \tag{2.8}$$

By continuity of ρ and (2.7), we can choose $x_0 \in (0, 1)$ such that $\rho(x_0) = 1$ and write

$$|\log \rho(x)| = \left| \int_{x_0}^x (\log \rho)_y dy \right| \leq \int_0^1 \frac{|\rho_y|}{\rho} dy.$$

Hence,

$$|\log \rho(x)| \leq \int_0^1 \frac{|\rho_x| \rho^{1/2}}{\rho^{3/2}} dx \leq \left[\int_0^1 \frac{\rho_x^2}{\rho^3} dx \right]^{1/2}.$$

Consequently,

$$\|\log \rho\|_{L^\infty} \leq \left[\int_0^1 \frac{\rho_x^2}{\rho^3} dx \right]^{1/2}.$$

From (2.8), the first bound (2.3) follows immediately. Regarding (2.4), we combine the upper bound in (2.3) with (2.8) to deduce that

$$\int_0^1 \rho_x^2 dx \leq \|\rho\|_{L^\infty}^3 \int_0^1 \frac{\rho_x^2}{\rho^3} dx \leq 8\mathcal{E}(\rho, u)e^{3[8\mathcal{E}(\rho, u)]^{1/2}}$$

Finally, (2.5) follows from the lower bound in (2.3). □

It is fairly clear from the above estimates that if $(\rho, u) \in \mathcal{X}$, then $\mathcal{E}(\rho, u) < \infty$. Viceversa, if (ρ, u) are smooth functions such that $\mathcal{E}(\rho, u) < \infty$, then $(\rho, u) \in \mathcal{X}$. The above lemma provides a quantification of this dichotomy.

It is crucial for us to prove various estimates on strong solutions to the compressible Navier–Stokes equations, keeping particular attention to the dependence on the parameter $A > 0$. We collect these estimates in the next two propositions, whose proofs are carried out in the subsequent sections.

Proposition 2.3 *Consider a solution (ρ, u) to (1.1)–(1.5) with initial data $(\rho_0, u_0) \in \mathcal{X}$ in (1.3) such that*

$$\mathcal{E}(\rho_0, u_0) < \infty.$$

Then there hold the entropy inequality

$$\mathbb{E}\mathcal{H}(\rho, u)(t) + \mathbb{E} \int_0^t \|u_x\|_{L^2}^2 ds \leq \mathcal{H}(\rho_0, u_0) + \frac{1}{2}\|\sigma\|_{L^\infty}^2 t, \tag{2.9}$$

and the energy inequality

$$\begin{aligned} \mathbb{E}\mathcal{E}(\rho, u)(t) + \frac{1}{2}\mathbb{E} \int_0^t \|u_x\|_{L^2}^2 ds + \frac{A^2}{2}\mathbb{E} \int_0^t \|(\log \rho)_x\|_{L^2}^2 ds \\ \leq \mathcal{E}(\rho_0, u_0) + \frac{1}{2}\|\sigma\|_{L^\infty}^2 t, \end{aligned} \tag{2.10}$$

for all $t \geq 0$. Moreover, the exponential martingale estimate

$$\begin{aligned} \mathbb{P} \left[\sup_{t \geq 0} \left(\mathcal{E}(\rho, u)(t) + \frac{1}{4} \int_0^t \left[\|u_x\|_{L^2}^2 + A^2 \|(\log \rho)_x\|_{L^2}^2 \right] ds \right. \right. \\ \left. \left. - \frac{1}{2}\|\sigma\|_{L^\infty}^2 t \right) - \mathcal{E}(\rho_0, u_0) \geq R \right] \leq e^{-\gamma_0 R}, \end{aligned} \tag{2.11}$$

holds for every $R \geq 0$ and for

$$\gamma_0 := \frac{\min \{1, 4A^2\}}{4\|\sigma\|_{L^\infty}^2}.$$

As a consequence, for every $m \geq 1$, there exists a constant $c_m > 0$ such that

$$\begin{aligned} \mathbb{E} \sup_{t \in [0, T]} \left(\mathcal{E}(\rho, u)(t) + \frac{1}{4} \int_0^t \left[\|u_x\|_{L^2}^2 + A^2 \|(\log \rho)_x\|_{L^2}^2 \right] ds \right)^m \\ \leq c_m \left(\mathcal{E}(\rho_0, u_0)^m + \|\sigma\|_{L^\infty}^{2m} T^m + \gamma_0^{-m} \right), \end{aligned} \tag{2.12}$$

for every $T \geq 1$, and the exponential moment bound holds

$$\begin{aligned} \mathbb{E} \exp \left(\frac{\gamma_0}{2} \sup_{t \in [0, T]} \left(\mathcal{E}(\rho, u)(t) + \frac{1}{4} \int_0^t \left[\|u_x\|_{L^2}^2 + A^2 \|(\log \rho)_x\|_{L^2}^2 \right] ds \right) \right) \\ \leq \exp \frac{\gamma_0}{2} \left(\mathcal{E}(\rho_0, u_0) + \frac{1}{2} \|\sigma\|_{L^\infty}^2 T \right), \end{aligned} \tag{2.13}$$

q for all $T \geq 1$.

The next result is instead a collection of similar, yet pathwise, estimates.

Proposition 2.4 Consider a solution (ρ, u) to (1.1)–(1.5) with initial data $(\rho_0, u_0) \in \mathcal{X}$ such that $\mathcal{E}(\rho_0, u_0) < \infty$. For every $T \geq 0$ and almost surely, there holds

$$\begin{aligned} \sup_{t \in [0, T]} \left[\mathcal{E}(\rho(t), u(t)) + \int_0^t \left[\|u_x\|_{L^2}^2 + A^2 \|(\log \rho)_x\|_{L^2}^2 \right] dt \right] \\ \leq C_1(\sigma W, A, \mathcal{E}(\rho_0, u_0), T). \end{aligned} \tag{2.14}$$

In particular,

$$\begin{aligned} \sup_{t \in [0, T]} \left[\|\rho(t)\|_{L^\infty} + \|\rho^{-1}(t)\|_{L^\infty} \right] \\ \leq C_2(\sigma W, A, \mathcal{E}(\rho_0, u_0), T). \end{aligned} \tag{2.15}$$

Moreover,

$$\sup_{t \in [0, T]} \left[\|u_x(t)\|_{L^2}^2 + \int_0^t \left[\int_0^1 \frac{|u_{xx}|^2}{\rho} dx \right] dt \right] \leq C_3(\sigma W, A, \mathcal{E}(\rho_0, u_0), \|(u_0)_x\|_{L^2}, T). \tag{2.16}$$

In the above the constants C_i can be explicitly computed.

The proof of (2.15) is simply a consequence of (2.3). We begin with Proposition 2.3.

2.1.1 Entropy Estimates

According to the Itô lemma

$$\begin{aligned} d(\rho u^2) &= d\left(\frac{1}{\rho}(\rho u)^2\right) = -\rho_t u^2 dt + \frac{1}{\rho}(2\rho u \cdot d(\rho u) + d(\rho u)d(\rho u)) \\ &= (\rho u)_x u^2 dt + 2u \left[u_{xx} - (\rho u^2 + A^2 \rho)_x \right] dt \\ &\quad + 2 \sum_{\ell=1}^{\infty} \rho u \sigma_{\ell} dW^{\ell} + \sum_{\ell=1}^{\infty} \rho |\sigma_{\ell}|^2 dt. \end{aligned}$$

Hence, integrating by parts,

$$\begin{aligned} d\frac{1}{2} \int_0^1 \rho u^2 dx + \|u_x\|_{L^2}^2 dt &= -A^2 \int_0^1 \rho_x u dx dt \\ &\quad + \sum_{\ell=1}^{\infty} \left(\int_0^1 \rho u \sigma_{\ell} dx \right) dW^{\ell} + \frac{1}{2} \sum_{\ell=1}^{\infty} \int_0^1 \rho |\sigma_{\ell}|^2 dx dt. \end{aligned}$$

Moreover, by integration by parts and using only (1.1), we have

$$\frac{d}{dt} \int_0^1 \rho \log \rho dx = - \int_0^1 (\rho u)_x [\log \rho + 1] dx = \int_0^1 \rho_x u dx.$$

Combining the above computations we find

$$d\mathcal{H}(\rho, u) + \|u_x\|_{L^2}^2 dt = \sum_{\ell=1}^{\infty} \left(\int_0^1 \rho u \sigma_{\ell} dx \right) dW^{\ell} + \frac{1}{2} \sum_{\ell=1}^{\infty} \int_0^1 \rho |\sigma_{\ell}|^2 dx dt, \tag{2.17}$$

so that, for every $t \geq 0$, we find the entropy balance

$$\mathbb{E}\mathcal{H}(\rho, u)(t) + \mathbb{E} \int_0^t \|u_x\|_{L^2}^2 ds = \mathcal{H}(\rho_0, u_0) + \frac{1}{2} \mathbb{E} \sum_{\ell=1}^{\infty} \int_0^t \int_0^1 \rho |\sigma_{\ell}|^2 dx ds.$$

Notice that by (1.4) and (2.2), we have

$$\sum_{\ell=1}^{\infty} \int_0^t \int_0^1 \rho |\sigma_{\ell}|^2 dx ds \leq \|\sigma\|_{L^{\infty}}^2 t,$$

so that

$$\mathbb{E}\mathcal{H}(\rho, u)(t) + \mathbb{E} \int_0^t \|u_x\|_{L^2}^2 ds \leq \mathcal{H}(\rho_0, u_0) + \frac{1}{2} \|\sigma\|_{L^{\infty}}^2 t.$$

This is precisely (2.9).

2.1.2 Energy Estimates

Let $\ell = \rho^{-1/2}$ so that

$$\begin{aligned} \ell_x &= -\frac{1}{2} \frac{\rho_x}{\rho^{3/2}}, & \ell_x^2 &= \frac{1}{4} \frac{\rho_x^2}{\rho^3}, \\ (\ell_x)_t &= -\frac{1}{2} \frac{(\rho_x)_t}{\rho^{3/2}} + \frac{3}{4} \frac{\rho_x \rho_t}{\rho^{5/2}} = \frac{1}{2} \frac{(\rho u)_{xx}}{\rho^{3/2}} - \frac{3}{4} \frac{\rho_x (\rho u)_x}{\rho^{5/2}}, \\ \ell_{xx} &= -\frac{1}{2} \frac{\rho_{xx}}{\rho^{3/2}} + \frac{3}{4} \frac{\rho_x^2}{\rho^{5/2}}. \end{aligned}$$

As such

$$\begin{aligned} (\ell_x^2)_t + (\ell_x^2 u)_x &= -\frac{\rho_x}{\rho^{3/2}} ((\ell_x)_t + \ell_{xx} u) + \ell_x^2 u_x \\ &= -\frac{\rho_x}{\rho^{3/2}} \left(\frac{1}{2} \frac{2\rho_x u_x + \rho u_{xx}}{\rho^{3/2}} - \frac{3}{4} \frac{\rho_x \rho u_x}{\rho^{5/2}} \right) + \frac{1}{4} \frac{\rho_x^2 u_x}{\rho^3} \\ &= -\frac{1}{2} \frac{\rho_x u_{xx}}{\rho^2}. \end{aligned}$$

Notice that this computation only makes use of (1.1). On the other hand

$$\begin{aligned} d\left(\frac{\rho_x u}{\rho}\right) &= \rho u \left(\frac{\rho_x}{\rho^2}\right)_t dt + \frac{\rho_x}{\rho^2} d(\rho u) \\ &= \rho u \left(\frac{(\rho_x)_t}{\rho^2} - 2\frac{\rho_x \rho_t}{\rho^3}\right) dt + \frac{\rho_x}{\rho^2} \left(u_{xx} - (\rho u^2 + A^2 \rho)_x\right) dt \\ &\quad + \sum_{\ell=1}^{\infty} \frac{\rho_x}{\rho} \sigma_{\ell} dW^{\ell} \\ &= \frac{\rho_x u_{xx}}{\rho^2} dt - \left(\frac{u(\rho u)_{xx}}{\rho} - 2\frac{u \rho_x (\rho u)_x}{\rho^2} + \frac{\rho_x (\rho u^2 + A^2 \rho)_x}{\rho^2}\right) dt \\ &\quad + \sum_{\ell=1}^{\infty} \frac{\rho_x}{\rho} \sigma_{\ell} dW^{\ell} \\ &= \frac{\rho_x u_{xx}}{\rho^2} dt - \left((uu_x)_x - u_x^2 + \frac{2\rho_x uu_x}{\rho} - \frac{\rho_x^2 u^2}{\rho^2} + \frac{u^2 \rho_{xx}}{\rho} + A^2 \frac{\rho_x^2}{\rho^2}\right) dt \\ &\quad + \sum_{\ell=1}^{\infty} \frac{\rho_x}{\rho} \sigma_{\ell} dW^{\ell} \\ &= \frac{\rho_x u_{xx}}{\rho^2} dt - \left((uu_x)_x - u_x^2 + \left(\frac{\rho_x u^2}{\rho}\right)_x + A^2 \frac{\rho_x^2}{\rho^2}\right) dt \\ &\quad + \sum_{\ell=1}^{\infty} \frac{\rho_x}{\rho} \sigma_{\ell} dW^{\ell}. \end{aligned}$$

Combining the above two equations and weighting appropriately we infer

$$d \left(\frac{\rho_x u}{\rho} + 2\ell_x^2 \right) + \left(\frac{\rho_x u^2}{\rho} + uu_x + 2\ell_x^2 u \right)_x dt + A^2 \frac{\rho_x^2}{\rho^2} dt = u_x^2 dt + \sum_{\ell=1}^{\infty} \frac{\rho_x}{\rho} \sigma_{\ell} dW^{\ell},$$

and hence

$$d \int_0^1 \left(\frac{\rho_x u}{\rho} + \frac{1}{2} \frac{\rho_x^2}{\rho^3} \right) dx + A^2 \int_0^1 \frac{\rho_x^2}{\rho^2} dx dt = \|u_x\|_{L^2}^2 dt + \sum_{\ell=1}^{\infty} \left(\int_0^1 \frac{\rho_x}{\rho} \sigma_{\ell} dx \right) dW^{\ell}. \tag{2.18}$$

We sum together (2.17) and half of (2.18) to finally obtain

$$\begin{aligned} d\mathcal{E}(\rho, u) + \frac{1}{2} \|u_x\|_{L^2}^2 dt + \frac{A^2}{2} \int_0^1 \frac{\rho_x^2}{\rho^2} dx dt \\ = \sum_{\ell=1}^{\infty} \left(\int_0^1 \left(\rho u + \frac{1}{2} \frac{\rho_x}{\rho} \right) \sigma_{\ell} dx \right) dW^{\ell} \\ + \frac{1}{2} \sum_{\ell=1}^{\infty} \int_0^1 \rho |\sigma_{\ell}|^2 dx dt. \end{aligned} \tag{2.19}$$

Hence

$$\begin{aligned} \mathbb{E}\mathcal{E}(\rho, u)(t) + \frac{1}{2} \mathbb{E} \int_0^t \|u_x\|_{L^2}^2 ds + \frac{A^2}{2} \mathbb{E} \int_0^t \int_0^1 \frac{\rho_x^2}{\rho^2} dx ds \\ = \mathcal{E}(\rho_0, u_0) + \frac{1}{2} \mathbb{E} \sum_{\ell=1}^{\infty} \int_0^t \int_0^1 \rho |\sigma_{\ell}|^2 dx ds, \end{aligned}$$

so that

$$\mathbb{E}\mathcal{E}(\rho, u)(t) + \frac{1}{2} \mathbb{E} \int_0^t \|u_x\|_{L^2}^2 ds + \frac{A^2}{2} \mathbb{E} \int_0^t \int_0^1 \frac{\rho_x^2}{\rho^2} dx ds \leq \mathcal{E}(\rho_0, u_0) + \frac{1}{2} \|\sigma\|_{L^{\infty}}^2 t,$$

and therefore (2.10) holds.

2.1.3 Exponential Estimates

It remains to prove (2.11), which follows from (2.19) and an application of the exponential martingale estimate

$$\mathbb{P} \left[\sup_{t \geq 0} \left(Z(t) - \frac{\gamma}{2} (Z)(t) \right) \geq R \right] \leq e^{-\gamma R}, \quad \forall R, \gamma > 0, \tag{2.20}$$

valid for any continuous martingale $\{Z(t)\}_{t \geq 0}$ with quadratic variation $\langle Z \rangle(t)$. Indeed, consider the time integrated version of (2.19), namely

$$\begin{aligned} \mathcal{E}(\rho, u)(t) &+ \frac{1}{2} \int_0^t \|u_x\|_{L^2}^2 ds + \frac{A^2}{2} \int_0^t \|(\log \rho)_x\|_{L^2}^2 ds \\ &= \mathcal{E}(\rho_0, u_0) + Z(t) + \frac{1}{2} \sum_{\ell=1}^{\infty} \int_0^t \int_0^1 \rho |\sigma_\ell|^2 dx ds. \end{aligned} \tag{2.21}$$

where $Z(t)$ is the martingale

$$Z(t) = \sum_{\ell=1}^{\infty} \int_0^t \left(\int_0^1 \left(\rho u + \frac{1}{2} \frac{\rho_x}{\rho} \right) \sigma_\ell dx \right) dW^\ell$$

with quadratic variation

$$\langle Z \rangle(t) = \sum_{\ell=1}^{\infty} \int_0^t \left[\int_0^1 \left(\rho u + \frac{1}{2} \frac{\rho_x}{\rho} \right) \sigma_\ell dx \right]^2 ds.$$

In view of the Poincaré-like inequality [32]

$$\int_0^1 \rho u^2 dx \leq \|u_x\|_{L^2}^2, \tag{2.22}$$

the mass constraint (1.4) and (2.2), we have

$$\sum_{\ell=1}^{\infty} \int_0^t \left[\int_0^1 \rho u \sigma_\ell dx \right]^2 ds \leq \|\sigma\|_{L^\infty}^2 \int_0^t \|u_x\|_{L^2}^2 ds$$

and

$$\frac{1}{4} \sum_{\ell=1}^{\infty} \int_0^t \left[\int_0^1 \frac{\rho_x}{\rho} \sigma_\ell dx \right]^2 ds \leq \frac{1}{4} \|\sigma\|_{L^\infty}^2 \int_0^t \|(\log \rho)_x\|_{L^2}^2 ds.$$

As a consequence, the quadratic variation of $Z(t)$ can be estimated as

$$\langle Z \rangle(t) \leq 2\|\sigma\|_{L^\infty}^2 \int_0^t \left[\|u_x\|_{L^2}^2 + \frac{1}{4} \|(\log \rho)_x\|_{L^2}^2 \right] ds. \tag{2.23}$$

Now, from (2.21) we infer that the functional

$$\Psi(t) = \mathcal{E}(\rho, u)(t) + \frac{1}{4} \int_0^t \|u_x\|_{L^2}^2 ds + \frac{A^2}{4} \int_0^t \|(\log \rho)_x\|_{L^2}^2 ds$$

satisfies with probability one the inequality

$$\begin{aligned} \Psi(t) - \frac{1}{2} \|\sigma\|_{L^\infty}^2 t &\leq \mathcal{E}(\rho_0, u_0) + \left[Z(t) - \frac{\gamma_0}{2} \langle Z \rangle(t) \right] \\ &\quad + \frac{\gamma_0}{2} \langle Z \rangle(t) - \frac{1}{4} \int_0^t \|u_x\|_{L^2}^2 ds - \frac{A^2}{4} \int_0^t \|(\log \rho)_x\|_{L^2}^2 ds \end{aligned} \tag{2.24}$$

where we conveniently fix the constant $\gamma_0 > 0$ as

$$\gamma_0 := \frac{\min\{1, 4A^2\}}{4\|\sigma\|_{L^\infty}^2}.$$

In this way, from (2.23) it follows that

$$\frac{\gamma_0}{2} \langle Z \rangle(t) - \frac{1}{4} \int_0^t \|u_x\|_{L^2}^2 ds - \frac{A^2}{4} \int_0^t \|(\log \rho)_x\|_{L^2}^2 ds \leq 0$$

and thus (2.24) implies

$$\Psi(t) - \frac{1}{2} \|\sigma\|_{L^\infty}^2 t \leq \mathcal{E}(\rho_0, u_0) + \left[Z(t) - \frac{\gamma_0}{2} \langle Z \rangle(t) \right]. \tag{2.25}$$

In turn, from (2.20) and the above (2.25) we deduce that

$$\begin{aligned} &\mathbb{P} \left[\sup_{t \geq 0} \left(\Psi(t) - \frac{1}{2} \|\sigma\|_{L^\infty}^2 t \right) - \mathcal{E}(\rho_0, u_0) \geq R \right] \\ &\leq \mathbb{P} \left[\sup_{t \geq 0} \left(Z(t) - \frac{\gamma_0}{2} \langle Z \rangle(t) \right) \geq R \right] \leq e^{-\gamma_0 R}, \end{aligned} \tag{2.26}$$

for every $R \geq 0$. This is exactly (2.11).

2.1.4 Polynomial and Exponential Moments

We now focus on (2.12), which is, in fact, an easy consequence of (2.11). Indeed, (2.26) implies that

$$\mathbb{P} \left[\sup_{t \in [0, T]} \Psi(t) - \mathcal{E}(\rho_0, u_0) - \frac{1}{2} \|\sigma\|_{L^\infty}^2 T \geq R \right] \leq e^{-\gamma_0 R}, \tag{2.27}$$

where Ψ is the functional defined in (2.24). Using that, for a non-negative random variable Z , any $m \geq 1$ and any constant $c > 0$, there holds

$$\mathbb{E}[Z^m] \leq 2^{m-1} \mathbb{E}[(Z - c)^m \mathbb{1}_{Z > c}] + 2^{m-1} c^m$$

$$= 2^{m-1} \int_0^\infty \mathbb{P}[Z - c > \lambda^{1/m}] d\lambda + 2^{m-1} c^m,$$

we infer from (2.27) that for some constant $c_m > 0$, independent of γ_0, T and $\mathcal{E}(\rho_0, u_0)$ there holds

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} \Psi(t)^m \right] &\leq 2^{m-1} \int_0^\infty e^{-\gamma_0 \lambda^{1/m}} d\lambda + 2^{m-1} \left(\mathcal{E}(\rho_0, u_0) + \frac{1}{2} \|\sigma\|_{L^\infty}^2 T \right)^m \\ &\leq c_m \left(\gamma_0^{-m} + \mathcal{E}(\rho_0, u_0)^m + \|\sigma\|_{L^\infty}^2 T^m \right), \end{aligned}$$

which is precisely (2.12). For (2.13), the idea is similar. Indeed, for a non-negative random variable Z and any $\delta, c > 0$, we also have that that

$$\mathbb{E}[e^{\delta Z}] \leq e^{\delta c} \mathbb{E} \left[e^{\delta(Z-c)} \mathbb{1}_{Z>c} \right] = e^{\delta c} \int_1^\infty \mathbb{P} \left[Z - c > \frac{\ln \lambda}{\delta} \right] d\lambda.$$

Hence, for any $\delta < \gamma_0$, we deduce from (2.27) that

$$\begin{aligned} \mathbb{E} \left[\exp \left(\delta \sup_{t \in [0, T]} \Psi(t) \right) \right] &\leq \exp \left[\delta \left(\mathcal{E}(\rho_0, u_0) + \frac{1}{2} \|\sigma\|_{L^\infty}^2 T \right) \right] \int_1^\infty \frac{1}{\lambda^{\gamma_0/\delta}} d\lambda \\ &= \frac{\delta}{\gamma_0 - \delta} \exp \left[\delta \left(\mathcal{E}(\rho_0, u_0) + \frac{1}{2} \|\sigma\|_{L^\infty}^2 T \right) \right], \end{aligned}$$

and (2.13) simply follows by choosing $\delta = \gamma_0/2$. We concluded the proof of Proposition 2.3.

2.2 Pathwise Estimates

By setting $\tilde{u} = u - w$, with $w = \sigma W$ we obtain from (1.1) to (1.2) that the pair (ρ, \tilde{u}) satisfies the system

$$\begin{aligned} \rho_t + (\rho \tilde{u})_x &= f(\rho, w) \\ (\rho \tilde{u})_t + (\rho \tilde{u}^2 + A^2 \rho)_x &= \tilde{u}_{xx} + \tilde{u} f(\rho, w) + \rho g(\rho, \tilde{u}, w), \end{aligned} \tag{2.28}$$

where

$$f(\rho, w) = -(\rho w)_x, \quad g(\rho, \tilde{u}, w) = \frac{w_{xx}}{\rho} - (\tilde{u} w)_x - w w_x, \tag{2.29}$$

with initial datum $(\rho_0, \tilde{u}_0) = (\rho_0, u_0)$. Notice that from (2.28) we can also write the velocity equation alone as

$$\tilde{u}_t + \tilde{u} \tilde{u}_x + A^2 \frac{\rho_x}{\rho} = \frac{\tilde{u}_{xx}}{\rho} + g(\rho, \tilde{u}, w), \tag{2.30}$$

which will also be useful in the sequel. Without explicit reference, we will use (2.2) multiple times to bound the various norms of w appearing in the estimates.

2.2.1 Pathwise Energy Estimates

We begin by proving (2.14). A lengthy calculation analogous to that in Sect. 2.1.2 shows that

$$\frac{d}{dt} \mathcal{E}(\rho, \tilde{u}) + \frac{1}{2} \|\tilde{u}_x\|_{L^2}^2 + \frac{A^2}{2} \|(\log \rho)_x\|_{L^2}^2 = \mathfrak{F}(\rho, \tilde{u}, f, g),$$

where

$$\begin{aligned} \mathfrak{F}(\rho, \tilde{u}, f, g) &= \int_0^1 \left(\frac{1}{2} f \tilde{u}^2 + A^2 f (\log \rho + 1) + \frac{1}{2} \frac{\rho_x}{\rho^2} f \tilde{u} \right. \\ &\quad \left. + \frac{1}{2} \frac{f_x \tilde{u}}{\rho} - \frac{\rho_x \tilde{u} f}{\rho^2} + \frac{1}{2} \frac{\rho_x f_x}{\rho^3} - \frac{3}{4} \frac{(\rho_x)^2 f}{\rho^4} \right) dx \\ &\quad + \int_0^1 \left(\rho \tilde{u} g + \frac{1}{2} \frac{\rho_x}{\rho} g \right) dx. \end{aligned}$$

Recalling (2.29), we may rewrite the above term as

$$\begin{aligned} \mathfrak{F}(\rho, \tilde{u}, f, g) &= -A^2 \int_0^1 (\rho w)_x \log \rho \, dx - \frac{1}{2} \int_0^1 \left[\frac{\rho_x (\rho w)_{xx}}{\rho^3} - \frac{3}{2} \frac{(\rho_x)^2 (\rho w)_x}{\rho^4} \right. \\ &\quad \left. - \frac{\rho_x}{\rho} \left(\frac{w_{xx}}{\rho} - w w_x \right) \right] dx \\ &\quad - \frac{1}{2} \int_0^1 \left[\left(\tilde{u}^2 - \frac{\rho_x \tilde{u}}{\rho^2} \right) (\rho w)_x + \frac{(\rho w)_{xx} \tilde{u}}{\rho} \right. \\ &\quad \left. - 2\rho \tilde{u} \left(\frac{w_{xx}}{\rho} - (\tilde{u} w)_x - w w_x \right) + \frac{\rho_x}{\rho} (\tilde{u} w)_x \right] dx. \end{aligned}$$

We now integrate by parts multiple times. Clearly,

$$-A^2 \int_0^1 (\rho w)_x \log \rho \, dx = A^2 \int_0^1 w \rho_x \, dx = -A^2 \int_0^1 w_x \rho \, dx.$$

Moreover, expanding all the derivatives and integrating by parts, the second term above reduces to

$$\begin{aligned} &\int_0^1 \left[\frac{\rho_x (\rho w)_{xx}}{\rho^3} - \frac{3}{2} \frac{(\rho_x)^2 (\rho w)_x}{\rho^4} - \frac{\rho_x}{\rho} \left(\frac{w_{xx}}{\rho} - w w_x \right) \right] dx \\ &= \int_0^1 \left[\frac{1}{2} \frac{[(\rho_x)^2]_x w}{\rho^3} + \frac{1}{2} \frac{(\rho_x)^2 w_x}{\rho^3} - \frac{3}{2} \frac{(\rho_x)^3 w}{\rho^4} + \frac{\rho_x}{\rho} w w_x \right] dx \end{aligned}$$

$$= \int_0^1 \frac{\rho_x}{\rho} w w_x dx.$$

Concerning the third term, similar arguments lead to

$$\begin{aligned} & \int_0^1 \left[\left(\tilde{u}^2 - \frac{\rho_x \tilde{u}}{\rho^2} \right) (\rho w)_x + \frac{(\rho w)_{xx} \tilde{u}}{\rho} \right. \\ & \quad \left. - 2\rho \tilde{u} \left(\frac{w_{xx}}{\rho} - (\tilde{u} w)_x - w w_x \right) + \frac{\rho_x}{\rho} (\tilde{u} w)_x \right] dx \\ &= \int_0^1 \left[\left(\tilde{u}^2 - \left(\frac{\tilde{u}}{\rho} \right)_x - \frac{\rho_x \tilde{u}}{\rho^2} \right) (\rho w)_x \right. \\ & \quad \left. + 2\tilde{u}_x w_x + 2\rho \tilde{u} (\tilde{u} w)_x + 2\rho \tilde{u} w w_x + \frac{\rho_x}{\rho} (\tilde{u} w)_x \right] dx \\ &= \int_0^1 \left[\tilde{u}_x w_x + 3\rho \tilde{u}^2 w_x + \tilde{u}^2 \rho_x w + 2\rho \tilde{u} \tilde{u}_x w + 2\rho \tilde{u} w w_x + \frac{\rho_x \tilde{u}}{\rho} w_x \right] dx \\ &= \int_0^1 \left[\tilde{u}_x w_x + 2\rho \tilde{u}^2 w_x + 2\rho \tilde{u} w w_x + \frac{\rho_x \tilde{u}}{\rho} w_x \right] dx. \end{aligned}$$

Therefore, collecting all of the above we find that

$$\begin{aligned} \mathfrak{F}(\rho, \tilde{u}, f, g) &= -A^2 \int_0^1 w_x \rho dx - \frac{1}{2} \int_0^1 \frac{\rho_x}{\rho} w w_x dx \\ & \quad - \frac{1}{2} \int_0^1 \left[\tilde{u}_x w_x + 2\rho \tilde{u}^2 w_x + 2\rho \tilde{u} w w_x + \frac{\rho_x \tilde{u}}{\rho} w_x \right] dx \end{aligned}$$

and using (1.4), (2.22), (2.6) and standard inequalities we deduce that

$$\begin{aligned} & |\mathfrak{F}(\rho, \tilde{u}, f, g)| \\ & \leq A^2 \|w_x\|_{L^\infty} + \|w_x\|_{L^\infty}^3 + \frac{1}{2} \|(\log \rho)_x\|_{L^2} \|w_x\|_{L^2}^2 \\ & \quad + \frac{1}{2} \|\tilde{u}_x\|_{L^2} \|w_x\|_{L^2} + c \|w_x\|_{L^\infty} \mathcal{E}(\rho, \tilde{u}) \end{aligned}$$

implying that

$$\begin{aligned} & \frac{d}{dt} \mathcal{E}(\rho, \tilde{u}) + \frac{1}{4} \|\tilde{u}_x\|_{L^2}^2 + \frac{A^2}{4} \|(\log \rho)_x\|_{L^2}^2 \leq c \|w_x\|_{L^\infty} \mathcal{E}(\rho, \tilde{u}) + \|w_x\|_{L^\infty}^3 \\ & \quad + A^2 \|w_x\|_{L^\infty} + c \|w_x\|_{L^2}^2 \left(1 + \frac{1}{A^2} \|w_x\|_{L^2}^2 \right). \end{aligned}$$

Hence, (2.14) follows from the standard Gronwall lemma, together with the fact that

$$\mathcal{E}(\rho, u) = \int_0^1 \left(\frac{1}{2} \rho (\tilde{u} + w)^2 + A^2 \rho \log \rho \right) dx + \frac{1}{2} \int_0^1 \left(\frac{\rho_x (\tilde{u} + w)}{\rho} + \frac{1}{2} \frac{\rho_x^2}{\rho^3} \right) dx$$

$$\leq c\mathcal{E}(\rho, \tilde{u}) + \int_0^1 \left(\rho w^2 + \frac{\rho_x w}{\rho} \right) dx \leq c\mathcal{E}(\rho, \tilde{u}) + c\|w_x\|_{L^2}^2.$$

2.2.2 H^1 Estimates on the Velocity

To prove (2.16), we multiply (2.30) by u_{xx} and integrate by parts, to obtain the identity

$$\begin{aligned} \frac{d}{dt} \|\tilde{u}_x\|_{L^2}^2 + 2 \int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx &= 2 \int_0^1 \tilde{u} \tilde{u}_x \tilde{u}_{xx} dx \\ &+ 2A^2 \int_0^1 \frac{\rho_x}{\rho} \tilde{u}_{xx} dx - 2 \int_0^1 g(\rho, \tilde{u}, w) \tilde{u}_{xx} dx. \end{aligned} \quad (2.31)$$

Now,

$$\begin{aligned} &2 \left| \int_0^1 \tilde{u} \tilde{u}_x \tilde{u}_{xx} dx \right| \\ &\leq 2 \left(\int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx \right)^{1/2} \|\rho^{1/2} \tilde{u}\|_{L^\infty} \|\tilde{u}_x\|_{L^2} \\ &\leq 2 \left(\int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx \right)^{1/2} \|\rho\|_{L^\infty}^{1/2} \|\tilde{u}_x\|_{L^2}^2 \\ &\leq \frac{1}{3} \int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx + c \|\rho\|_{L^\infty} \|\tilde{u}_x\|_{L^2}^4, \end{aligned}$$

and

$$\begin{aligned} 2A^2 \left| \int_0^1 \tilde{u}_{xx} \frac{\rho_x}{\rho} dx \right| &\leq \frac{1}{3} \int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx + cA^4 \int_0^1 \frac{\rho_x^2}{\rho} dx \\ &\leq \frac{1}{3} \int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx + cA^4 \|\rho\|_{L^\infty} \|(\log \rho)_x\|_{L^2}^2. \end{aligned}$$

Concerning the last term, we find

$$\begin{aligned} &2 \left| \int_0^1 g(\rho, \tilde{u}, w) \tilde{u}_{xx} dx \right| \\ &= 2 \left| \int_0^1 \rho^{1/2} \left(\frac{w_{xx}}{\rho} - (\tilde{u}w)_x - w w_x \right) \frac{\tilde{u}_{xx}}{\rho^{1/2}} dx \right| \\ &\leq c \left[\|\rho^{-1}\|_{L^\infty} \|w_{xx}\|_{L^2}^2 + (1 + \|\rho\|_{L^\infty}) \|w_x\|_{L^\infty}^2 \|\tilde{u}_x\|_{L^2}^2 + \|w_x\|_{L^\infty}^4 \right] \\ &\quad + \frac{1}{3} \int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx. \end{aligned}$$

Thus, (2.31) and the above estimates entail

$$\begin{aligned} & \frac{d}{dt} \|\tilde{u}_x\|_{L^2}^2 + \int_0^1 \frac{|\tilde{u}_{xx}|^2}{\rho} dx \\ & \leq c \left[\|\rho\|_{L^\infty} \|\tilde{u}_x\|_{L^2}^2 + (1 + \|\rho\|_{L^\infty}) \|w_x\|_{L^\infty}^2 \right] \|\tilde{u}_x\|_{L^2}^2 \\ & \quad + c \left[\|\rho^{-1}\|_{L^\infty} \|w_{xx}\|_{L^2}^2 + \|w_x\|_{L^\infty}^4 + A^4 \|\rho\|_{L^\infty} \|(\log \rho)_x\|_{L^2}^2 \right]. \end{aligned}$$

In view of (2.14), (2.15) and Lemma 2.2, we can apply the Gronwall lemma to the above inequality and deduce (2.16).

2.3 Uniqueness of Strong Solutions

Let (ρ, u) and (r, v) be two solutions to (1.1)–(1.2). The relative entropy between (ρ, u) and (r, v) is defined as the functional

$$\mathcal{H}_r(\rho, u|r, v) = \int_0^1 \left(\frac{1}{2} \rho (u - v)^2 + A^2 \rho \log \frac{\rho}{r} \right) dx.$$

It is well-known that the above functional is not a distance, and it is not even symmetric. Nonetheless $\mathcal{H}_r(\rho, u|r, v) = 0$ if and only if $(\rho, u) = (r, v)$. More precisely, the following well-known facts hold.

Lemma 2.5 *Let $(\rho, u) \in \mathcal{X}$ and $(r, v) \in \mathcal{X}$. Then*

$$\frac{\min\{1, A^2\}}{2} \min \left\{ \|\rho^{-1}\|_{L^\infty}, \|r^{-1}\|_{L^\infty} \right\} \left[\|u - v\|_{L^2}^2 + \|\rho - r\|_{L^2}^2 \right] \leq \mathcal{H}_r(\rho, u|r, v) \tag{2.32}$$

and

$$\mathcal{H}_r(\rho, u|r, v) \leq \frac{1}{2} \|\rho\|_{L^\infty} \|u - v\|_{L^2}^2 + \frac{A^2}{2} \max \left\{ \|\rho^{-1}\|_{L^\infty}, \|r^{-1}\|_{L^\infty} \right\} \|\rho - r\|_{L^2}^2. \tag{2.33}$$

Proof The proof is based on the fact that the function $\xi \mapsto F(\xi) := \xi \log \xi - \xi + 1$ is convex. In particular,

$$F'(\xi) = \log \xi, \quad F''(\xi) = \frac{1}{\xi}, \quad \forall \xi > 0.$$

Moreover, due to (1.4), the relative entropy can be rewritten as

$$\mathcal{H}_r(\rho, u|r, v) = \int_0^1 \left(\frac{1}{2} \rho (u - v)^2 + A^2 [F(\rho) - F(r) - (\rho - r)F'(r)] \right) dx.$$

Therefore, using Taylor’s theorem, on the one hand we have the lower bound

$$\frac{1}{2} \min \left\{ \frac{1}{\xi_1}, \frac{1}{\xi_2} \right\} (\xi_1 - \xi_2)^2 \leq F(\xi_1) - F(\xi_2) - (\xi_1 - \xi_2)F'(\xi_2)$$

while on the other hand the upper bound reads

$$F(\xi_1) - F(\xi_2) - (\xi_1 - \xi_2)F'(\xi_2) \leq \frac{1}{2} \max \left\{ \frac{1}{\xi_1}, \frac{1}{\xi_2} \right\} (\xi_1 - \xi_2)^2.$$

Using the above bounds in the expression for \mathcal{H}_r yields precisely (2.32)–(2.33), and the proof is over. □

We then have the following uniqueness result.

Theorem 2.6 *Let (ρ, u) and (r, v) be two solutions of (1.1)–(1.5), corresponding to the same initial condition $(\rho_0, u_0) \in \mathcal{X}$. Then*

$$\mathbb{P}[(\rho(t), u(t)) = (r(t), v(t)), \forall t \geq 0] = 1.$$

Proof We compute the time derivative of $\mathcal{H}_r(\rho, u|r, v)$. Firstly, notice that (1.2) can be rewritten as

$$du + \left(uu_x + A^2 \frac{\rho_x}{\rho} \right) dt = \frac{u_{xx}}{\rho} dt + \sigma dW,$$

so that the difference of the velocities satisfies

$$(u - v)_t + uu_x - vv_x + A^2 \left[\log \left(\frac{\rho}{r} \right) \right]_x = \frac{u_{xx}}{\rho} - \frac{v_{xx}}{r}, \tag{2.34}$$

Therefore, making use of (1.1) and (2.34), we have that

$$\begin{aligned} & \frac{d}{dt} \frac{1}{2} \int_0^1 \rho(u - v)^2 dx \\ &= \frac{1}{2} \int_0^1 \rho_t(u - v)^2 dx + \int_0^1 \rho(u - v)(u - v)_t dx \\ &= -\frac{1}{2} \int_0^1 (\rho u)_x(u - v)^2 dx + \int_0^1 \rho(u - v) \left(\frac{u_{xx}}{\rho} - \frac{v_{xx}}{r} \right) dx \\ &\quad - \int_0^1 \rho(u - v)(uu_x - vv_x) dx - A^2 \int_0^1 \rho(u - v) \left[\log \left(\frac{\rho}{r} \right) \right]_x dx \\ &= \int_0^1 \rho u(u - v)(u - v)_x dx - \|(u - v)_x\|_{L^2}^2 + \int_0^1 \frac{\rho - r}{r} (v - u)v_{xx} dx \\ &\quad - \int_0^1 \rho(u - v)(uu_x - vv_x) dx - A^2 \int_0^1 \rho(u - v) \left[\log \left(\frac{\rho}{r} \right) \right]_x dx \end{aligned}$$

$$\begin{aligned}
 &= -\|(u - v)_x\|_{L^2}^2 + \int_0^1 \frac{\rho - r}{r} (v - u)v_{xx} dx - \int_0^1 v_x \rho (u - v)^2 dx \\
 &\quad - A^2 \int_0^1 \rho (u - v) \left[\log \left(\frac{\rho}{r} \right) \right]_x dx.
 \end{aligned} \tag{2.35}$$

On the other hand, due to (1.1) once more, we compute

$$\begin{aligned}
 \frac{d}{dt} \int_0^1 \rho \log \left(\frac{\rho}{r} \right) dx &= \int_0^1 \rho_t \log \left(\frac{\rho}{r} \right) dx + \int_0^1 \left(\rho_t - \frac{\rho}{r} r_t \right) dx \\
 &= - \int_0^1 (\rho u)_x \log \left(\frac{\rho}{r} \right) dx - \int_0^1 (\rho u)_x dx + \int_0^1 \frac{\rho}{r} (rv)_x dx \\
 &= \int_0^1 \rho u \left[\log \left(\frac{\rho}{r} \right) \right]_x dx + \int_0^1 \rho v_x dx + \int_0^1 \frac{\rho}{r} r_x v dx \\
 &= \int_0^1 \rho u \left[\log \left(\frac{\rho}{r} \right) \right]_x dx - \int_0^1 \rho_x v dx + \int_0^1 \frac{\rho}{r} r_x v dx \\
 &= \int_0^1 \rho (u - v) \left[\log \left(\frac{\rho}{r} \right) \right]_x dx.
 \end{aligned} \tag{2.36}$$

Therefore, combining (2.35) and (2.36), we arrive at

$$\frac{d}{dt} \mathcal{H}_r(\rho, u|r, v) + \|(u - v)_x\|_{L^2}^2 = \int_0^1 \frac{\rho - r}{r} (v - u)v_{xx} dx - \int_0^1 v_x \rho (u - v)^2 dx. \tag{2.37}$$

We now estimate each term in the right-hand side above. We preliminary notice that for any smooth function g such that $g(x_0) = 0$ for some $x_0 \in [0, 1]$, there holds the inequality

$$\|g\|_{L^\infty} \leq \left(\int_0^1 \frac{|g_x|^2}{\rho} dx \right)^{1/2}, \tag{2.38}$$

where ρ can be replaced by any positive function with mass at most 1 (and, in particular, by r). Indeed,

$$\begin{aligned}
 |g(x)| &= \left| \int_{x_0}^x g_y dy \right| = \left| \int_{x_0}^x \rho^{1/2} \frac{g_y}{\rho^{1/2}} dy \right| \\
 &\leq \left(\int_{x_0}^x \rho dy \right)^{1/2} \left(\int_{x_0}^x \frac{|g_y|^2}{\rho} dy \right)^{1/2} \leq \left(\int_0^1 \frac{|g_x|^2}{\rho} dx \right)^{1/2}.
 \end{aligned}$$

Concerning the first term in the right-hand side of (2.37), using standard inequalities and (2.32) we obtain

$$\int_0^1 \frac{\rho - r}{r} (v - u)v_{xx} dx$$

$$\begin{aligned}
&\leq \|r^{-1}\|_{L^\infty}^{1/2} \|\rho - r\|_{L^2} \|v - u\|_{L^\infty} \left(\int_0^1 \frac{|v_{xx}|^2}{r} dx \right)^{1/2} \\
&\leq \|r^{-1}\|_{L^\infty} \left(\int_0^1 \frac{|v_{xx}|^2}{r} dx \right) \|\rho - r\|_{L^2}^2 + \frac{1}{2} \|(u - v)_x\|_{L^2}^2 \\
&\leq \frac{2}{\min\{1, A^2\} \min\{\|\rho^{-1}\|_{L^\infty}, \|r^{-1}\|_{L^\infty}\}} \left(\int_0^1 \frac{|v_{xx}|^2}{r} dx \right) \mathcal{H}_r(\rho, u|r, v) \\
&\quad + \frac{1}{2} \|(u - v)_x\|_{L^2}^2 \\
&\leq \frac{2}{\min\{1, A^2\}} \max\left\{1, \frac{\|r^{-1}\|_{L^\infty}}{\|\rho^{-1}\|_{L^\infty}}\right\} \left(\int_0^1 \frac{|v_{xx}|^2}{r} dx \right) \mathcal{H}_r(\rho, u|r, v) \\
&\quad + \frac{1}{2} \|(u - v)_x\|_{L^2}^2,
\end{aligned}$$

where the last line follows from (2.32). For the second term, we use (2.22) and (2.38) to deduce that

$$\begin{aligned}
\int_0^1 v_x \rho (u - v)^2 dx &\leq \|v_x\|_{L^\infty} \left(\int_0^1 \rho (u - v)^2 dx \right)^{1/2} \|(u - v)_x\|_{L^2} \\
&\leq \left(\int_0^1 \frac{|v_{xx}|^2}{r} dx \right) \mathcal{H}_r(\rho, u|r, v) + \frac{1}{2} \|(u - v)_x\|_{L^2}^2.
\end{aligned}$$

Hence, from (2.37) it follows that

$$\begin{aligned}
&\frac{d}{dt} \mathcal{H}_r(\rho, u|r, v) \\
&\leq \left(1 + \frac{2}{\min\{1, A^2\}} \max\left\{1, \frac{\|r^{-1}\|_{L^\infty}}{\|\rho^{-1}\|_{L^\infty}}\right\} \right) \left(\int_0^1 \frac{|v_{xx}|^2}{r} dx \right) \mathcal{H}_r(\rho, u|r, v).
\end{aligned} \tag{2.39}$$

Thanks to Proposition 2.4, the term multiplying $\mathcal{H}_r(\rho, u|r, v)$ in the right-hand side above is locally integrable in time. We now apply the standard Gronwall lemma to (2.39) and use the fact that

$$\mathcal{H}_r(\rho, u|r, v)|_{t=0} = 0$$

by assumption to conclude that the two solutions are indistinguishable. \square

2.4 Continuous Dependence of Solutions

Regarding the continuous dependence of solutions, the picture is more complicated, and can be proven in the following situation.

Theorem 2.7 *Let $\{(\rho^n, u^n)\}_{n \in \mathbb{N}}$ be a sequence of solutions to (1.1)–(1.5) with initial data $\{(\rho_0^n, u_0^n)\}_{n \in \mathbb{N}}$ such that*

$$\sup_{n \in \mathbb{N}} \mathcal{E}(\rho_0^n, u_0^n) < \infty, \tag{2.40}$$

and let (ρ, u) be a solution to (1.1)–(1.5) with initial datum $(\rho_0, u_0) \in \mathcal{X}$. If

$$(\rho_0^n, u_0^n) \rightarrow (\rho_0, u_0) \text{ in } L^2 \times L^2, \tag{2.41}$$

then

$$(\rho^n, u^n) \rightarrow (\rho, u) \text{ a.s. in } L^2 \times L^2,$$

for all $t \geq 0$.

Proof We use the approach in Theorem 2.6. Arguing in the same way as we did to derive (2.39), we arrive at

$$\begin{aligned} & \frac{d}{dt} \mathcal{H}_r(\rho^n, u^n | \rho, u) \\ & \leq \left(1 + \frac{2}{\min\{1, A^2\}} \max \left\{ 1, \frac{\|\rho^{-1}\|_{L^\infty}}{\|(\rho^n)^{-1}\|_{L^\infty}} \right\} \right) \left(\int_0^1 \frac{|u_{xx}|^2}{\rho} dx \right) \mathcal{H}_r(\rho^n, u^n | \rho, u). \end{aligned} \tag{2.42}$$

Due to the lack of symmetry of the relative entropy, it is important that we consider the above quantity, and not the other possible $\mathcal{H}_r(\rho, u | \rho^n, u^n)$. Let

$$R = \max \left\{ \sup_{n \in \mathbb{N}} \mathcal{E}(\rho_0^n, u_0^n), \mathcal{E}(\rho_0, u_0) \right\} < \infty.$$

In view of Proposition 2.4, we have n -independent almost sure bounds of the form

$$\sup_{t \in [0, T]} \int_0^t \left(\int_0^1 \frac{|u_{xx}|^2}{\rho} dx \right) ds \leq C(T, \sigma W, R, \|(u_0)_x\|_{L^2}),$$

and

$$\sup_{t \in [0, T]} \frac{\|\rho^{-1}\|_{L^\infty}^2}{\|\rho_n^{-1}\|_{L^\infty}^2} \leq C(T, \sigma W, R).$$

The point is that the latter bound does not depend on $\|(u_0^n)_x\|_{L^2}$, which is not, in general, under control uniformly in $n \in \mathbb{N}$. Thanks to Lemma 2.5, the fact that $\mathcal{E}(\rho_0^n, u_0^n) \leq R$ and (2.41), we have that

$$\lim_{n \rightarrow \infty} \mathcal{H}_r(\rho_0^n, u_0^n | \rho_0, u_0) = 0.$$

Thus, the standard Gronwall lemma applied to (2.42) implies

$$\lim_{n \rightarrow \infty} \mathcal{H}_r(\rho^n(t), u^n(t) | \rho(t), u(t)) = 0, \quad \text{a.s.}$$

for all $t \geq 0$. Hence, a further application of Lemma 2.5 leads to the conclusion of the proof. \square

3 Invariant Measures

In this section, we prove of the main results of this article, namely the existence of invariant measures to (1.1)–(1.2). As mentioned earlier, we will work in the phase space

$$\mathcal{X} = \left\{ (\rho, u) \in H^1(0, 1) \times H_0^1(0, 1) : \int_0^1 \rho(x) dx = 1, \rho > 0 \right\}.$$

However, in view of Theorem 2.7, the correct topology does not seem to be the natural one induced by the product norm in \mathcal{X} , since Lemma 2.5 suggests that continuous dependence for our system holds in a weaker sense. We make this precise here below.

3.1 The Markovian Framework

We denote by $\rho(t; \rho_0)$, $u(t; u_0)$ the unique solution to (1.1)–(1.2), with (deterministic) initial data $(\rho_0, u_0) \in \mathcal{X}$. For a set $B \in \mathcal{B}(\mathcal{X}_{L^2})$, we define the transition functions

$$\mathcal{P}_t(\rho_0, u_0, B) = \mathbb{P}((\rho(t; \rho_0), u(t; u_0)) \in B),$$

for any $t \geq 0$. For $t \geq 0$, define the Markov semigroup

$$\mathcal{P}_t \phi(\rho_0, u_0) = \mathbb{E} \phi(\rho(t; \rho_0), u(t; u_0)), \quad \phi \in M_b(\mathcal{X}_{L^2}).$$

The usual semigroup properties

$$\mathcal{P}_0 = \text{identity on } M_b(\mathcal{X}_{L^2})$$

and

$$\mathcal{P}_{t+s} = \mathcal{P}_t \circ \mathcal{P}_s, \quad \forall t, s \geq 0,$$

follow from the existence and uniqueness results for our system. The goal is to prove the existence of a probability measure $\mu \in \mathfrak{P}(\mathcal{X}_{L^2})$ that is *invariant* under \mathcal{P}_t , namely such that

$$\int_{\mathcal{X}} \mathcal{P}_t \phi d\mu = \int_{\mathcal{X}} \phi d\mu, \quad \forall \phi \in C_b(\mathcal{X}_{L^2}).$$

Now, it is clear from the estimates in Propositions 2.3 and 2.4 that

$$\mathcal{P}_t : M_b(\mathcal{X}_{L^2}) \rightarrow M_b(\mathcal{X}_{L^2}),$$

namely, \mathcal{P}_t is well defined on measurable bounded functions. The Feller property, namely that \mathcal{P}_t maps $C_b(\mathcal{X}_{L^2})$ into $C_b(\mathcal{X}_{L^2})$ is more delicate, and we are not able to prove it at the moment. In particular, it does not follow from Theorem 2.7, due to the additional requirement (2.40). However, Theorem 2.7 suggests the following definition.

Definition 3.1 Let $\phi \in M_b(\mathcal{X}_{L^2})$. We say that $\phi : \mathcal{X} \rightarrow \mathbb{R}$ belongs to the class \mathcal{G} if

$$\lim_{n \rightarrow \infty} \phi(\rho^n, u^n) = \phi(\rho, u),$$

whenever

$$\{(\rho^n, u^n)\}_{n \in \mathbb{N}} \subset \mathcal{X}_{L^2}, \quad \lim_{n \rightarrow \infty} (\rho^n, u^n) = (\rho, u) \text{ in } \mathcal{X}_{L^2}, \quad \text{and} \quad \sup_{n \in \mathbb{N}} \mathcal{E}(\rho^n, u^n) < \infty. \tag{3.1}$$

From the above definition and Lemma 2.2, it is clear that

$$C_b(\mathcal{X}_{L^2}) \subset \mathcal{G} \subset C_b(\mathcal{X}_{H^1}).$$

It turns out that \mathcal{G} is also invariant under \mathcal{P}_t .

Lemma 3.2 *The class \mathcal{G} is invariant under the Markov semigroup \mathcal{P}_t , namely*

$$\mathcal{P}_t : \mathcal{G} \rightarrow \mathcal{G}.$$

Proof Let $\{(\rho_0^n, u_0^n)\}_{n \in \mathbb{N}} \subset \mathcal{X}_{L^2}$ be a sequence, complying with (3.1), namely

$$\lim_{n \rightarrow \infty} (\rho_0^n, u_0^n) = (\rho_0, u_0) \text{ in } \mathcal{X}_{L^2}, \quad \text{and} \quad M := \sup_{n \in \mathbb{N}} \mathcal{E}(\rho_0^n, u_0^n) < \infty. \tag{3.2}$$

In light of Theorem 2.7,

$$\lim_{n \rightarrow \infty} (\rho(t; \rho_0^n), u(t; u_0^n)) = (\rho(t; \rho_0), u(t; u_0)) \text{ a.s. in } \mathcal{X}_{L^2}, \quad \forall t \geq 0. \tag{3.3}$$

Moreover, in view of (2.10) and (3.2), there exists a constant $\bar{M} := \bar{M}(t, M, \sigma)$ independent of $n \in \mathbb{N}$ such that

$$\mathbb{E} \mathcal{E}(\rho(t; \rho_0^n), u(t; u_0^n)) \leq \bar{M},$$

which, by Chebyshev’s inequality, it implies that

$$\mathbb{P} [\mathcal{E}(\rho(t; \rho_0^n), u(t; u_0^n)) > R] \leq \frac{\bar{M}}{R}, \tag{3.4}$$

for any $R > 0$. Fix $\phi \in \mathcal{G}$, and let

$$M_\phi := \sup_{(\rho, u) \in \mathcal{X}_{L^2}} |\phi(\rho, u)| < \infty. \tag{3.5}$$

To show that $\mathcal{P}_t \phi \in \mathcal{G}$, we are required to prove that for any fixed $t > 0$ there holds

$$\lim_{n \rightarrow \infty} \mathcal{P}_t \phi(\rho_0^n, u_0^n) = \mathcal{P}_t \phi(\rho_0, u_0).$$

Writing $\mathcal{E}_n = \mathcal{E}(\rho(t; \rho_0^n), u(t; u_0^n))$ for short, we have by definition that

$$\begin{aligned} & \mathcal{P}_t \phi(\rho_0^n, u_0^n) - \mathcal{P}_t \phi(\rho_0, u_0) \\ &= \mathbb{E} [\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))] \\ &= \mathbb{E} [(\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n \leq R}] \\ & \quad + \mathbb{E} [(\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n > R}]. \end{aligned}$$

In light of (3.4)–(3.5),

$$|\mathbb{E} [(\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n > R}]| \leq \frac{2\bar{M}M_\phi}{R}. \tag{3.6}$$

Since $\phi \in \mathcal{G}$, we also deduce by (3.2) and (3.3) that

$$\lim_{n \rightarrow \infty} (\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n \leq R} = 0, \quad \text{a.s.,}$$

and therefore by the bounded convergence theorem that

$$\lim_{n \rightarrow \infty} \mathbb{E} [(\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n \leq R}] = 0. \tag{3.7}$$

Let us now arbitrarily fix $\varepsilon > 0$, and pick $R_\varepsilon = 4\bar{M}M_\phi/\varepsilon$. Invoking (3.6), we deduce that

$$\mathbb{E} [(\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n > R}] \leq \frac{\varepsilon}{2}.$$

Moreover, from (3.7) there exists $n_\varepsilon \in \mathbb{N}$ such that

$$|\mathbb{E} [(\phi(\rho(t; \rho_0^n), u(t; u_0^n)) - \phi(\rho(t; \rho_0), u(t; u_0))) \mathbb{1}_{\mathcal{E}_n \leq R}]| < \frac{\varepsilon}{2}, \quad \forall n \geq n_\varepsilon.$$

Thus,

$$|\mathcal{P}_t \phi(\rho_0^n, u_0^n) - \mathcal{P}_t \phi(\rho_0, u_0)| < \varepsilon, \quad \forall t \geq 0, n \geq n_\varepsilon,$$

and the proof is over. □

3.2 Tightness of Time-Averaged Measures

We prove the existence of an invariant measure for \mathcal{P}_T via the classical Krylov–Bogoliubov procedure. However, since we are working in a non-complete metric space, and with a Markov semigroup that is not Feller, some details do not follow directly from the well-known theory. In what follows, the parameter $A > 0$ appearing in (1.2) is fixed, but we suppress the dependence on it of all the quantities (except for bounds), in order to keep the notation as simple as possible. As initial conditions to our problem, we fix $\rho_0 = 1$ and $u_0 = 0$. Notice that the corresponding energy vanishes, namely $\mathcal{E}(1, 0) = 0$. For $T > 0$, define the time-averaged measure on \mathcal{X} by

$$\mu_T(B) = \frac{1}{T} \int_0^T \mathbb{P}(\rho(t; 1), u(t; 0) \in B) dt, \tag{3.8}$$

where $B \in \mathcal{B}(\mathcal{X}_{L^2})$. The first step is to prove tightness of the family $\{\mu_T\}_{T>0}$. We will make use of the following lemma.

Lemma 3.3 *Assume that $\rho \in H^1$ is such that $\rho > 0$ and*

$$\int_0^1 \rho \, dx = 1, \quad M^2 := \int_0^1 \frac{\rho_x^2}{\rho^2} \, dx = \int_0^1 [(\log \rho)_x]^2 \, dx < \infty.$$

Then, for $x \in [0, 1]$, we have the pointwise bounds

$$e^{-M} \leq \rho(x) \leq e^M, \tag{3.9}$$

and, moreover

$$\int_0^1 \rho_x^2 \, dx \leq M^2 e^{2M}.$$

Proof Firstly, we notice since ρ is continuous and has mean 1, there exists $x_0 \in (0, 1)$ such that $\rho(x_0) = 1$. Therefore, for every $x \in [0, 1]$ we have

$$|\log \rho(x)| = \left| \int_{x_0}^x (\log \rho)_y \, dy \right| \leq \left[\int_0^1 [(\log \rho)_y]^2 \, dy \right]^{1/2} = M, \quad \forall x \in [0, 1].$$

As a consequence,

$$e^{-M} \leq \rho(x) \leq e^M, \quad \forall x \in [0, 1],$$

proving (3.9). Moreover,

$$\int_0^1 \rho_x^2 \, dx \leq \|\rho\|_{L^\infty}^2 \int_0^1 \frac{\rho_x^2}{\rho^2} \, dx.$$

The claim then follows from (3.9). □

As a consequence, tightness of $\{\mu_T\}_{T>0}$ follows in a straightforward manner.

Proposition 3.4 *The family of probability measures $\{\mu_T\}_{T>0} \subset \mathfrak{P}(\mathcal{X}_{L^2})$ is tight. Hence, there exists a subsequence $T_j \rightarrow \infty$ and a measure $\mu \in \mathfrak{P}(\mathcal{X}_{L^2})$ such that*

$$\lim_{j \rightarrow \infty} \int_{\mathcal{X}} \phi \, d\mu_{T_j} = \int_{\mathcal{X}} \phi \, d\mu, \quad \forall \phi \in C_b(\mathcal{X}_{L^2}). \tag{3.10}$$

Proof For any fixed $R, S \geq 1$, define the sets

$$K_R = \left\{ (\rho, u) \in \mathcal{X} : \int_0^1 u_x^2 \, dx + \int_0^1 \frac{\rho_x^2}{\rho^2} \, dx \leq R^2 \right\}$$

and

$$C_S = \left\{ (\rho, u) \in \mathcal{X} : \int_0^1 u_x^2 \, dx + \int_0^1 \rho_x^2 \, dx + \|\rho\|_{L^\infty} + \|\rho^{-1}\|_{L^\infty} \leq S \right\}. \tag{3.11}$$

Note that since C_S is bounded in $H^1 \times H_0^1$ and is closed, it is compact in \mathcal{X}_{L^2} . By Lemma 3.3, we have that if $(\rho, u) \in K_R$, then

$$\int_0^1 \rho_x^2 \, dx \leq R^2 e^{2R}, \quad \int_0^1 u_x^2 \, dx \leq R^2, \quad \|\rho\|_{L^\infty} \leq e^R, \quad \|\rho^{-1}\|_{L^\infty} \leq e^R,$$

In particular, if $R \geq 1$, we obtain

$$\int_0^1 \rho_x^2 \, dx + \int_0^1 u_x^2 \, dx + \|\rho\|_{L^\infty} + \|\rho^{-1}\|_{L^\infty} \leq S_R := 4R^2 e^{2R}, \tag{3.12}$$

which therefore translates into the set inclusion

$$K_R \subset C_{S_R}. \tag{3.13}$$

In view of (3.13) and Chebyshev’s inequality we have

$$\begin{aligned} \mu_T(C_{S_R}) &\geq \mu_T(K_R) = 1 - \mu_T(\mathcal{X} \setminus K_R) \\ &= 1 - \frac{1}{T} \int_0^T \mathbb{P} \left[\int_0^1 \frac{\rho_x^2}{\rho^2} \, dx + \int_0^1 u_x^2 \, dx > R^2 \right] \\ &\geq 1 - \frac{1}{R^2} \frac{1}{T} \int_0^T \mathbb{E} \left[\int_0^1 \frac{\rho_x^2}{\rho^2} \, dx + \int_0^1 u_x^2 \, dx \right]. \end{aligned}$$

In light of the energy inequality (2.10) and the fact that $\mathcal{E}(1, 0) = 0$, it follows that

$$\mu_T(C_{S_R}) \geq 1 - \frac{\|\sigma\|_{L^\infty}^2}{\min\{1, A^2\}} \frac{1}{R^2}. \tag{3.14}$$

Hence, the family $\{\mu_T\}_{T>0}$ is tight, and therefore there exists a subsequential limit $\mu \in \mathfrak{P}(\mathcal{X}_{L^2})$. Note that this uses the direction of Prokhorov’s theorem that does not require completeness (see [1, Theorem 6.1]). Now, since C_{S_R} is a closed set, the Portmanteau theorem implies

$$\mu(\mathcal{X}) \geq \mu(C_{S_R}) \geq \limsup_{T \rightarrow \infty} \mu_T(C_{S_R}) \geq 1 - \frac{\|\sigma\|_{L^\infty}^2}{\min\{1, A^2\}} \frac{1}{R^2}. \tag{3.15}$$

Notice again that this does not require the metric space \mathcal{X}_{L^2} to be complete (see [1, Theorem 2.1]). Thus

$$\mu(\mathcal{X}) \geq \lim_{R \rightarrow \infty} \mu(C_{S_R}) = 1.$$

The proof is therefore concluded. □

3.3 Invariance of the Limit Measure

We aim to prove the following proposition.

Proposition 3.5 *For every fixed $A > 0$, the Markov semigroup $\{\mathcal{P}_t\}_{t \geq 0}$ associated to (1.1)–(1.5) possesses an invariant probability measure $\mu_A \in \mathfrak{P}(\mathcal{X}_{L^2})$. Furthermore,*

$$\int_{\mathcal{X}} \left[A^2 \|(\log \rho)_x\|_{L^2}^2 + \|u_x\|_{L^2}^2 \right] \mu_A(\rho, u) \leq \|\sigma\|_{L^\infty}^2. \tag{3.16}$$

The main issue here is that \mathcal{P}_t is not known to be Feller, and therefore we cannot directly apply (3.10) to $\mathcal{P}_t \phi$. As a first step, we extend property (3.10) to the class \mathcal{G} .

Lemma 3.6 *Let $\{\mu_{T_j}\}_{j \in \mathbb{N}} \subset \mathfrak{P}(\mathcal{X}_{L^2})$ be the convergent sequence in Proposition 3.4. Then*

$$\lim_{j \rightarrow \infty} \int_{\mathcal{X}} \phi \, d\mu_{T_j} = \int_{\mathcal{X}} \phi \, d\mu, \quad \forall \phi \in \mathcal{G}.$$

Proof Let $\phi \in \mathcal{G}$, arbitrarily fix $R \geq 1$, and consider the compact sets C_{S_R} from (3.11), where S_R is given by (3.12). Since $\phi \in \mathcal{G}$, ϕ is bounded on \mathcal{X}_{L^2} , so that

$$M_\phi := \sup_{(\rho, u) \in \mathcal{X}_{L^2}} |\phi(\rho, u)| < \infty.$$

Moreover, the restriction of ϕ to C_{S_R} is continuous (in the $L^2 \times L^2$ metric) on C_{S_R} , for every $R \geq 1$. Since C_{S_R} is a closed set, Tietze extension theorem (see e.g. [35]) guarantees the existence of a function $\tilde{\phi}_R \in C_b(\mathcal{X}_{L^2})$ such that

$$\tilde{\phi}_R = \phi \quad \text{on } C_{S_R}, \quad \sup_{(\rho, u) \in \mathcal{X}_{L^2}} |\tilde{\phi}_R(\rho, u)| \leq M_\phi, \quad \forall R \geq 1. \tag{3.17}$$

Now,

$$\begin{aligned} \left| \int_{\mathcal{X}} \phi \, d\mu_{T_j} - \int_{\mathcal{X}} \phi \, d\mu \right| &\leq \left| \int_{\mathcal{X}} (\phi - \tilde{\phi}_R) \, d\mu_{T_j} \right| \\ &+ \left| \int_{\mathcal{X}} \tilde{\phi}_R \, d\mu_{T_j} - \int_{\mathcal{X}} \tilde{\phi}_R \, d\mu \right| + \left| \int_{\mathcal{X}} (\tilde{\phi}_R - \phi) \, d\mu \right| \end{aligned}$$

Fix $\varepsilon > 0$. By (3.14) and the above (3.17), there exists $R_\varepsilon \geq 1$ such that

$$\begin{aligned} \left| \int_{\mathcal{X}} (\phi - \tilde{\phi}_{R_\varepsilon}) \, d\mu_{T_j} \right| &= \left| \int_{\mathcal{X} \setminus C_{SR_\varepsilon}} (\phi - \tilde{\phi}_{R_\varepsilon}) \, d\mu_{T_j} \right| \leq 2M_\phi \mu_{T_j}(\mathcal{X} \setminus C_{SR_\varepsilon}) \\ &\leq \frac{2M_\phi \|\sigma\|_{L^\infty}^2}{\min\{1, A^2\}} \frac{1}{R_\varepsilon^2} < \frac{\varepsilon}{3}. \end{aligned}$$

uniformly for $j \in \mathbb{N}$, and, similarly, in light of (3.15),

$$\left| \int_{\mathcal{X}} (\tilde{\phi}_{R_\varepsilon} - \phi) \, d\mu \right| < \frac{\varepsilon}{3}.$$

Since $\tilde{\phi}_{R_\varepsilon} \in C_b(\mathcal{X}_{L^2})$, property (3.10) implies the existence of $j_\varepsilon \in \mathbb{N}$ such that

$$\left| \int_{\mathcal{X}} \tilde{\phi}_{R_\varepsilon} \, d\mu_{T_j} - \int_{\mathcal{X}} \tilde{\phi}_{R_\varepsilon} \, d\mu \right| < \frac{\varepsilon}{3}, \quad \forall j \geq j_\varepsilon.$$

Hence, for every $\varepsilon > 0$, there exists $j_\varepsilon \in \mathbb{N}$ such that

$$\left| \int_{\mathcal{X}} \phi \, d\mu_{T_j} - \int_{\mathcal{X}} \phi \, d\mu \right| < \varepsilon, \quad \forall j \geq j_\varepsilon,$$

which proves the claim. □

We are now in the position to prove invariance of the limiting measure and the bound (3.16).

Proof of Proposition 3.5 As before, $A > 0$ is fixed and we suppress the various dependences on it. Let $\phi \in C_b(\mathcal{X}_{L^2})$ be fixed. Thanks to Lemma 3.2,

$$\mathcal{P}_t \phi \in \mathcal{G}, \quad \forall t > 0.$$

In view of Lemma 3.6, we then have that

$$\lim_{j \rightarrow \infty} \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu_{T_j} = \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu, \quad \forall t \geq 0.$$

Hence, by (3.8), we conclude that

$$\begin{aligned} & \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu \\ &= \lim_{j \rightarrow \infty} \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu_{T_j} = \lim_{j \rightarrow \infty} \frac{1}{T_j} \int_0^{T_j} \mathcal{P}_{t+s} \phi(1, 0) \, ds = \lim_{j \rightarrow \infty} \frac{1}{T_j} \int_t^{T_j+t} \mathcal{P}_s \phi(1, 0) \, ds \\ &= \lim_{j \rightarrow \infty} \left(\frac{1}{T_j} \int_0^{T_j} \mathcal{P}_s \phi(1, 0) \, ds + \frac{1}{T_j} \int_{T_j}^{T_j+t} \mathcal{P}_s \phi(1, 0) \, ds - \frac{1}{T_j} \int_0^t \mathcal{P}_s \phi(1, 0) \, ds \right) \\ &= \lim_{j \rightarrow \infty} \int_{\mathcal{X}} \phi \, d\mu_{T_j} = \int_{\mathcal{X}} \phi \, d\mu, \end{aligned}$$

so that invariance follows from the arbitrariness of ϕ . Lastly, (3.16) is deduced directly from (2.10). Indeed, is $(\rho^S, u^S) \in \mathcal{X}$ is a statistically stationary solution distributed as μ , the we can use (2.10) to derive the bound

$$\mathbb{E} \|u_x^S\|_{L^2}^2 + A^2 \mathbb{E} \|(\log \rho^S)_x\|_{L^2}^2 \leq \|\sigma\|_{L^\infty}^2,$$

which is precisely (3.16) after the usual change of variables. The proof is over. □

Acknowledgements The authors would like to thank Jacob Bedrossian, Sandra Cerrai, Dave Levermore, Jonathan Mattingly, Sam Punshon-Smith and Mohammed Ziane for helpful discussions. We also thank the program “New Challenges in PDE: Deterministic Dynamics and Randomness in High Infinite Dimensional Systems”, held at the Mathematical Science Research Institute (MSRI) in the Fall 2015, where some initial steps in the work were formulated. MCZ was in part supported by the National Science Foundation under the Award DMS-1713886. K.T. gratefully acknowledges the support in part by the National Science Foundation under the Award DMS-1614964. NGH was in part supported by the National Science Foundation under the Award DMS-1313272 and Simons Foundation 515990.

Compliance with Ethical Standards

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Appendix A: A General Result on Existence of Invariant Measures

The results in Sect. 3 on the existence of invariant measures can be stated in greater generality as follows. Let (\mathcal{X}, d_s) be a metric space, and assume we are given another metric d_w on \mathcal{X} such that

- (1) Any d_s -bounded subset of \mathcal{X} is d_w -precompact.
- (2) If $d_s(x_n, y_n) \rightarrow 0$ as $n \rightarrow \infty$ for some $x_n, y_n \in \mathcal{X}$, then $d_w(x_n, y_n) \rightarrow 0$ as $n \rightarrow \infty$.

We denote by \mathcal{X}_w the space \mathcal{X} endowed with the metric d_w . Given $x \in \mathcal{X}$, let $\{X_t^x\}_{t \geq 0}$ be a Markov process on \mathcal{X} with $X_0^x = x$, and define the time-averaged measures

$$\mu_T^x(B) = \frac{1}{T} \int_0^T \mathbb{P}(X_t^x \in B) dt, \quad T > 0, \tag{A.1}$$

where $B \in \mathcal{B}(\mathcal{X}_w)$, the Borel σ -algebra of \mathcal{X}_w . For $t \geq 0$, define the Markov semigroup

$$\mathcal{P}_t \phi(x) = \mathbb{E} \phi(X_t^x), \quad \phi \in M_b(\mathcal{X}).$$

We assume that the Markov semigroup is well defined on measurable bounded functions, namely the property that $\mathcal{P}_t : M_b(\mathcal{X}_w) \rightarrow M_b(\mathcal{X}_w)$ is well-defined.

Definition A.1 Let $\phi \in M_b(\mathcal{X}_w)$. We say that $\phi : \mathcal{X} \rightarrow \mathbb{R}$ belongs to the class \mathcal{G} if $\phi \in C(K)$ for any d_s -bounded set $K \subset \mathcal{X}$.

The following is a slight variant of the Krylov–Bogoliubov procedure to prove the existence of an invariant measure for \mathcal{P}_t .

Theorem A.2 Assume that for some initial condition $x \in \mathcal{X}$ the following two properties hold.

- (Tightness) For every $\varepsilon \in (0, 1]$, there exists a d_s -bounded and d_w -closed set $K_\varepsilon \subset \mathcal{X}$ such that the time-averaged measures $\{\mu_T^x\}_{T > 0}$ satisfy

$$\inf_{T > 0} \mu_T^x(K_\varepsilon) \geq 1 - \varepsilon. \tag{A.2}$$

- (Almost Feller) For all $t > 0$,

$$\mathcal{P}_t C_b(\mathcal{X}_w) \subset \mathcal{G}. \tag{A.3}$$

Then there exists an invariant measure μ for \mathcal{P}_t .

Proof From the tightness property (A.2) and the fact that d_s -bounded sets are d_w -precompact, we infer the existence of a sequence $T_j \rightarrow \infty$ and a measure $\mu \in \mathfrak{P}(\mathcal{X}_w)$ such that

$$\lim_{j \rightarrow \infty} \int_{\mathcal{X}} \phi d\mu_{T_j}^x = \int_{\mathcal{X}} \phi d\mu, \quad \forall \phi \in C_b(\mathcal{X}_w). \tag{A.4}$$

Let $\varepsilon > 0$, and let K_ε be the corresponding sets as above. Now, since K_ε is a d_w -closed set, the Portmanteau theorem and (A.2) imply

$$\mu(K_\varepsilon) \geq \limsup_{T \rightarrow \infty} \mu_T(K_\varepsilon) \geq 1 - \varepsilon. \tag{A.5}$$

We now claim that, in fact,

$$\lim_{j \rightarrow \infty} \int_{\mathcal{X}} \phi d\mu_{T_j}^x = \int_{\mathcal{X}} \phi d\mu, \quad \forall \phi \in \mathcal{G}. \tag{A.6}$$

To prove the claim, let $\phi \in \mathcal{G}$ be arbitrarily fixed, and set

$$M_\phi := \sup_{y \in \mathcal{X}} |\phi(y)| < \infty.$$

It is important to notice here that M_ϕ does not depend on ε . We can then apply Tietze extension theorem (see e.g. [35]) and deduce the existence of a function $\tilde{\phi}_\varepsilon \in C_b(\mathcal{X}_w)$ such that

$$\tilde{\phi}_\varepsilon = \phi \quad \text{on } K_\varepsilon, \quad \sup_{y \in \mathcal{X}} |\tilde{\phi}_\varepsilon(y)| \leq M_\phi, \quad \forall \varepsilon \in (0, 1]. \tag{A.7}$$

Now,

$$\begin{aligned} \left| \int_{\mathcal{X}} \phi \, d\mu_{T_j}^x - \int_{\mathcal{X}} \phi \, d\mu \right| &\leq \left| \int_{\mathcal{X}} (\phi - \tilde{\phi}_\varepsilon) \, d\mu_{T_j}^x \right| \\ &+ \left| \int_{\mathcal{X}} \tilde{\phi}_\varepsilon \, d\mu_{T_j}^x - \int_{\mathcal{X}} \tilde{\phi}_\varepsilon \, d\mu \right| + \left| \int_{\mathcal{X}} (\tilde{\phi}_\varepsilon - \phi) \, d\mu \right|. \end{aligned}$$

By (A.2) and the above (A.7), we have

$$\left| \int_{\mathcal{X}} (\phi - \tilde{\phi}_\varepsilon) \, d\mu_{T_j}^x \right| = \left| \int_{\mathcal{X} \setminus K_\varepsilon} (\phi - \tilde{\phi}_\varepsilon) \, d\mu_{T_j}^x \right| \leq 2M_\phi \mu_{T_j}^x(\mathcal{X} \setminus K_\varepsilon) < 2M_\phi \varepsilon.$$

uniformly for $j \in \mathbb{N}$, and, similarly, in light of (A.5),

$$\left| \int_{\mathcal{X}} (\tilde{\phi}_\varepsilon - \phi) \, d\mu \right| < 2M_\phi \varepsilon.$$

Since $\tilde{\phi}_\varepsilon \in C_b(\mathcal{X}_{L^2})$, property (A.4) implies the existence of $j_\varepsilon \in \mathbb{N}$ such that

$$\left| \int_{\mathcal{X}} \tilde{\phi}_\varepsilon \, d\mu_{T_j}^x - \int_{\mathcal{X}} \tilde{\phi}_\varepsilon \, d\mu \right| < 2M_\phi \varepsilon, \quad \forall j \geq j_\varepsilon.$$

Hence, for every $\varepsilon > 0$, there exists $j_\varepsilon \in \mathbb{N}$ such that

$$\left| \int_{\mathcal{X}} \phi \, d\mu_{T_j}^x - \int_{\mathcal{X}} \phi \, d\mu \right| < 6M_\phi \varepsilon, \quad \forall j \geq j_\varepsilon,$$

which proves (A.6). We are now left to prove that μ is invariant for \mathcal{P}_t . Let $\phi \in C_b(\mathcal{X}_w)$ be fixed. Thanks to assumption (A.3),

$$\mathcal{P}_t \phi \in \mathcal{G}, \quad \forall t > 0.$$

In view of (A.6), we then have that

$$\lim_{j \rightarrow \infty} \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu_{T_j}^x = \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu, \quad \forall t \geq 0.$$

Hence, by (A.1), we conclude that

$$\begin{aligned}
 & \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu \\
 &= \lim_{j \rightarrow \infty} \int_{\mathcal{X}} \mathcal{P}_t \phi \, d\mu_{T_j}^x = \lim_{j \rightarrow \infty} \frac{1}{T_j} \int_0^{T_j} \mathcal{P}_{t+s} \phi(x) \, ds = \lim_{j \rightarrow \infty} \frac{1}{T_j} \int_t^{T_j+t} \mathcal{P}_s \phi(x) \, ds \\
 &= \lim_{j \rightarrow \infty} \left(\frac{1}{T_j} \int_0^{T_j} \mathcal{P}_s \phi(x) \, ds + \frac{1}{T_j} \int_{T_j}^{T_j+t} \mathcal{P}_s \phi(x) \, ds - \frac{1}{T_j} \int_0^t \mathcal{P}_s \phi(x) \, ds \right) \\
 &= \lim_{j \rightarrow \infty} \int_{\mathcal{X}} \phi \, d\mu_{T_j}^x = \int_{\mathcal{X}} \phi \, d\mu,
 \end{aligned}$$

so that invariance follows from the arbitrariness of ϕ . □

References

1. Billingsley, P.: *Convergence of Probability Measures*, Second, Wiley Series in Probability and Statistics: Probability and Statistics. Wiley, New York (1999). A Wiley-Interscience Publication
2. Breit, D., Hofmanová, M.: Stochastic Navier–Stokes equations for compressible fluids. *Indiana Univ. Math. J.* **65**(4), 1183–1250 (2016)
3. Breit, D., Feireisl, E., Hofmanová, M.: Incompressible limit for compressible fluids with stochastic forcing. *Arch. Ration. Mech. Anal.* **222**(2), 895–926 (2016)
4. Breit, D., Feireisl, E., Hofmanová, M., Maslowski, B.: Stationary solutions to the compressible Navier–Stokes system driven by stochastic forces (2017). [arXiv:1703.03177](https://arxiv.org/abs/1703.03177)
5. Breit, D., Feireisl, E., Hofmanová, M.: Local strong solutions to the stochastic compressible Navier–Stokes system. *Commun. Partial Differ. Equ.* **43**(2), 313–345 (2018)
6. Bricmont, J., Kupiainen, A., Lefevere, R.: Ergodicity of the 2D Navier–Stokes equations with random forcing. *Commun. Math. Phys.* **224**(1), 65–81 (2001)
7. Bricmont, J., Kupiainen, A., Lefevere, R.: Exponential mixing of the 2D stochastic Navier–Stokes dynamics. *Commun. Math. Phys.* **230**(1), 87–132 (2002)
8. Chang, T., Hsiao, L.: *The Riemann Problem and Interaction of Waves in Gas Dynamics*, Longman Scientific & Technical, Harlow. Wiley, New York (1989)
9. Chen, G.-Q., Glimm, J.: Global solutions to the compressible euler equations with geometrical structure. *Commun. Math. Phys.* **180**, 153–193 (1996)
10. Da Prato, G., Debussche, A.: Differentiability of the transition semigroup of the stochastic Burgers equation, and application to the corresponding Hamilton–Jacobi equation. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.* **9**(4), 267–277 (1999)
11. Da Prato, G., Debussche, A.: Ergodicity for the 3D stochastic Navier–Stokes equations. *J. Math. Pures Appl.* (9) **82**(8), 877–947 (2003)
12. Da Prato, G., Debussche, A.: On the martingale problem associated to the 2D and 3D stochastic Navier–Stokes equations. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **19**(3), 247–264 (2008)
13. Da Prato, G., Gatarek, D.: Stochastic Burgers equation with correlated noise. *Stoch. Stoch. Rep.* **52**(1–2), 29–41 (1995)
14. Da Prato, G., Zabczyk, J.: *Ergodicity for Infinite-Dimensional Systems*, London Mathematical Society Lecture Note Series, vol. 229. Cambridge University Press, Cambridge (1996)
15. Desjardins, B., Grenier, E., Lions, P.-L., Masmoudi, N.: Incompressible limit for solutions of the isentropic Navier–Stokes equations with Dirichlet boundary conditions. *J. Math. Pures Appl.* (9) **78**(5), 461–471 (1999)
16. Feireisl, E.: *Dynamics of Viscous Compressible Fluids*, vol. 26. Oxford University Press, Oxford (2004)
17. Feireisl, E., Novotný, A.: The low Mach number limit for the full Navier–Stokes–Fourier system. *Arch. Ration. Mech. Anal.* **186**(1), 77–107 (2007)

18. Feireisl, E., Petzeltová, H.: Asymptotic compactness of global trajectories generated by the Navier–Stokes equations of a compressible fluid. *J. Differ. Equ.* **173**(2), 390–409 (2001)
19. Feireisl, E., Petzeltová, H.: On the long-time behaviour of solutions to the Navier–Stokes–Fourier system with a time-dependent driving force. *J. Dyn. Differ. Equ.* **19**(3), 685–707 (2007)
20. Feireisl, E., Maslowski, B., Novotný, A.: Compressible fluid flows driven by stochastic forcing. *J. Differ. Equ.* **254**(3), 1342–1358 (2013)
21. Flandoli, F., Gatarek, D.: Martingale and stationary solutions for stochastic Navier–Stokes equations. *Probab. Theory Relat. Fields* **102**(3), 367–391 (1995)
22. Flandoli, F., Maslowski, B.: Ergodicity of the 2-D Navier–Stokes equation under random perturbations. *Commun. Math. Phys.* **172**(1), 119–141 (1995)
23. Flandoli, F., Romito, M.: Markov selections for the 3D stochastic Navier–Stokes equations. *Probab. Theory Relat. Fields* **140**(3–4), 407–458 (2008)
24. Gas, O.M.O.A.V.: Mesure invariante pour l'équation stochastique du mouvement d'un gaz visqueux en une dimension avec la discrétisation du domaine. *Rev. Roum. Math. Pures Appl.* **58**(2), 149–162 (2013)
25. Goldys, B., Maslowski, B.: Exponential ergodicity for stochastic Burgers and 2D Navier–Stokes equations. *J. Funct. Anal.* **226**(1), 230–255 (2005)
26. Hairer, M.: Exponential mixing properties of stochastic PDEs through asymptotic coupling. *Probab. Theory Relat. Fields* **124**(3), 345–380 (2002)
27. Hairer, M., Mattingly, J.C.: Ergodicity of the 2D Navier–Stokes equations with degenerate stochastic forcing. *Ann. Math. (2)* **164**(3), 993–1032 (2006)
28. Hoff, D.: Global existence for 1D, compressible, isentropic Navier–Stokes equations with large initial data. *Trans. Am. Math. Soc.* **303**(1), 169–181 (1987)
29. Hoff, D.: Global solutions of the equations of one-dimensional, compressible flow with large data and forces, and with differing end states. *Z. Angew. Math. Phys.* **49**(5), 774–785 (1998)
30. Hoff, D., Serre, D.: The failure of continuous dependence on initial data for the Navier–Stokes equations of compressible flow. *SIAM J. Appl. Math.* **51**, 887–898 (1991)
31. Hoff, D., Smoller, J.: Non-formation of vacuum states for compressible Navier–Stokes equations. *Commun. Math. Phys.* **216**(2), 255–276 (2001)
32. Hoff, D., Ziane, M.: The global attractor and finite determining nodes for the Navier–Stokes equations of compressible flow with singular initial data. *Indiana Univ. Math. J.* **49**(3), 843–889 (2000)
33. Hoff, D., Ziane, M.: Finite-dimensional attractors and exponential attractors for the Navier–Stokes equations of compressible flow. *SIAM J. Math. Anal.* **34**(5), 1040–1063 (2003). (electronic)
34. Kazhikhov, A.V., Shelukhin, V.V.: Unique global solution with respect to time of initial-boundary-value problems for one-dimensional equations of a viscous gas. *J. Appl. Math. Mech.* **41**, 273–282 (1977)
35. Kelley, J.L.: *General Topology*. D. Van Nostrand Company Inc, Toronto (1955)
36. Kuksin, S., Shirikyan, A.: Stochastic dissipative PDEs and Gibbs measures. *Commun. Math. Phys.* **213**(2), 291–330 (2000)
37. Kuksin, S., Shirikyan, A.: *Mathematics of Two-Dimensional Turbulence*, vol. 194. Cambridge University Press, Cambridge (2012)
38. Lions, P.-L.: *Mathematical Topics in Fluid Mechanics*, Vol. 2, vol. 10. The Clarendon Press/Oxford University Press, New York (1998)
39. Lions, P.-L., Masmoudi, N.: Incompressible limit for a viscous compressible fluid. *J. Math. Pures Appl.* (9) **77**(6), 585–627 (1998)
40. Masmoudi, N.: Examples of singular limits in hydrodynamics. In: *Handbook of Differential Equations: Evolutionary Equations*, vol. III, pp. 195–275. North-Holland, Amsterdam (2007)
41. Masmoudi, N., Young, L.-S.: Ergodic theory of infinite dimensional systems with applications to dissipative parabolic PDEs. *Commun. Math. Phys.* **227**(3), 461–481 (2002)
42. Mattingly, J.C.: Exponential convergence for the stochastically forced Navier–Stokes equations and other partially dissipative dynamics. *Commun. Math. Phys.* **230**(3), 421–462 (2002)
43. Mellet, A., Vasseur, A.: Existence and uniqueness of global strong solutions for one-dimensional compressible Navier–Stokes equations. *SIAM J. Math. Anal.* **39**(4), 1344–1365 (2008)
44. Sinaï, Ya G.: Two results concerning asymptotic behavior of solutions of the Burgers equation with force. *J. Stat. Phys.* **64**(1–2), 1–12 (1991)
45. Smith, S.A.: Random perturbations of viscous, compressible fluids: global existence of weak solutions. *SIAM J. Math. Anal.* **49**(6), 4521–4578 (2017)

46. Solonnikov, V.A.: The solvability of the initial-boundary value problem for the equations of motion of a viscous compressible fluid. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **56**, 128–142 (1976). 197. Investigations on linear operators and theory of functions, VI
47. Tornatore, E.: Global solution of bi-dimensional stochastic equation for a viscous gas. *NoDEA Non-linear Differ. Equ. Appl.* **7**(4), 343–360 (2000)
48. Tornatore, E., Yashima, H.F.: One-dimensional equations of a barotropic viscous gas with a not very regular perturbation. *Ann. Univ. Ferrara Sez. VII (N.S.)* **7**(40), 137–168 (1996)
49. Tornatore, E., Yashima, H.F.: One-dimensional stochastic equations for a viscous barotropic gas. *Ricerche Mat.* **46**(2), 255–283 (1998)
50. Wang, D., Wang, H.: Global existence of martingale solutions to the three-dimensional stochastic compressible Navier–Stokes equations. *Differ. Integral Equ.* **28**(11–12), 1105–1154 (2015)
51. Weinan, E., Khanin, K., Mazel, A., Sinai, Ya.: Invariant measures for Burgers equation with stochastic forcing. *Ann. Math. (2)* **151**(3), 877–960 (2000)
52. Weinan, E., Mattingly, J.C., Sinai, Ya.G.: Gibbsian dynamics and ergodicity for the stochastically forced Navier–Stokes equation. *Commun. Math. Phys.* **224**(1), 83–106 (2001)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.