

## Erratum to: Characterization of two proline dipeptidases (prolidases) from the hyperthermophilic archaeon *Pyrococcus horikoshii*


Casey M. Theriot · Sherry R. Tove · Amy M. Grunden

Published online: 27 October 2009  
© Springer-Verlag 2009

**Erratum to: Appl Microbiol Biotechnol**  
DOI 10.1007/s00253-009-2235-x

This article unfortunately contained a mistake.

Figure 2 was presented incorrectly. The correct Fig. 2 which shows Phprol (PH1149) and PhIprol (PH0974) relative activities over pH range 4–10 is displayed below.



**Fig. 2** Activity of *P. horikoshii* prolidases over a pH range of 4.0–10.0. Prolidase assays contained 14.8 ng for Phprol and 6.2 ng PhIprol, Met-Pro (4 mM) and CoCl<sub>2</sub> (1.2 mM). The following buffers were used for each pH at a final concentration of 100 mM: pH4.0–5.0, Sodium Acetate; pH6.0–8.0, MOPS; pH9.0, CHES; pH10.0, CAPS. 100% specific activity corresponds to 2,321 U/mg for Phprol and 3,357 U/mg for PhIprol

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00253-009-2235-x>.

C. M. Theriot · S. R. Tove  
Department of Microbiology, North Carolina State University,  
4548 Gardner Hall, Campus Box 7615, Raleigh,  
NC 27695-7615, USA

A. M. Grunden (✉)  
Department of Microbiology, North Carolina State University,  
4550A Gardner Hall, Campus Box 7615, Raleigh,  
NC 27695-7615, USA  
e-mail: amy\_grunden@ncsu.edu