



# O<sub>2</sub> versus N<sub>2</sub>O respiration in a continuous microbial enrichment

Monica Conthe<sup>1</sup> · Camiel Parchen<sup>1</sup> · Gerben Stouten<sup>1</sup> · Robbert Kleerebezem<sup>1</sup> · Mark C. M. van Loosdrecht<sup>1</sup>

Received: 14 April 2018 / Revised: 13 July 2018 / Accepted: 15 July 2018 / Published online: 27 July 2018  
© The Author(s) 2018

## Abstract

Despite its ecological importance, essential aspects of microbial N<sub>2</sub>O reduction—such as the effect of O<sub>2</sub> availability on the N<sub>2</sub>O sink capacity of a community—remain unclear. We studied N<sub>2</sub>O vs. aerobic respiration in a chemostat culture to explore (i) the extent to which simultaneous respiration of N<sub>2</sub>O and O<sub>2</sub> can occur, (ii) the mechanism governing the competition for N<sub>2</sub>O and O<sub>2</sub>, and (iii) how the N<sub>2</sub>O-reducing capacity of a community is affected by dynamic oxic/anoxic shifts such as those that may occur during nitrogen removal in wastewater treatment systems. Despite its prolonged growth and enrichment with N<sub>2</sub>O as the sole electron acceptor, the culture readily switched to aerobic respiration upon exposure to O<sub>2</sub>. When supplied simultaneously, N<sub>2</sub>O reduction to N<sub>2</sub> was only detected when the O<sub>2</sub> concentration was limiting the respiration rate. The biomass yields per electron accepted during growth on N<sub>2</sub>O are in agreement with our current knowledge of electron transport chain biochemistry in model denitrifiers like *Paracoccus denitrificans*. The culture's affinity constant ( $K_S$ ) for O<sub>2</sub> was found to be two orders of magnitude lower than the value for N<sub>2</sub>O, explaining the preferential use of O<sub>2</sub> over N<sub>2</sub>O under most environmentally relevant conditions.

**Keywords** Nitrous oxide · Mixotrophy · Enrichment · Chemostat

## Introduction

Coping with rising levels of the potent greenhouse gas nitrous oxide (N<sub>2</sub>O) in the atmosphere calls for the development of mitigation strategies to reduce N<sub>2</sub>O accumulation and emission in soil management and wastewater treatment (WWT). The presence and activity of N<sub>2</sub>O-reducing organisms in fertilized soils and WWT plants, such as bacteria and archaea harboring *nosZ*-type genes, may be key in such mitigating strategies (Thomson et al. 2012). Nitrous oxide reductase (N<sub>2</sub>OR), the enzyme encoded by the *nosZ* gene, is a terminal reductase present in some microbial respiratory electron transport chains (ETC) that catalyzes the only microbial reaction known to consume N<sub>2</sub>O, converting it to innocuous N<sub>2</sub> (which constitutes 79% of the Earth's atmosphere). Although N<sub>2</sub>O reduction is

generally associated to denitrifying organisms, many N<sub>2</sub>O reducers lack reductases other than N<sub>2</sub>OR (i.e., nitrate-, nitrite-, or nitric oxide-reductase; Hallin et al. 2018). However, most, if not all, denitrifiers—and presumably N<sub>2</sub>O reducers—are facultative aerobes, having the terminal oxidases necessary for O<sub>2</sub> respiration (van Spanning and Richardson 2007).

Based on what is known on the biochemistry of model organisms like *Paracoccus denitrificans*, N<sub>2</sub>O and O<sub>2</sub> respiration presumably share the core of the ETC (Chen and Strous 2013), with electrons branching out to O<sub>2</sub> (via cytochrome oxidases), N<sub>2</sub>O (via N<sub>2</sub>OR), or other NO<sub>x</sub> (in denitrifying N<sub>2</sub>O reducers) depending on electron acceptor availability. It is a common notion that, when both N<sub>2</sub>O and O<sub>2</sub> are available, N<sub>2</sub>O reducers will consume O<sub>2</sub> preferentially over N<sub>2</sub>O (and other N oxides; Shapleigh 2013). Even though N<sub>2</sub>O is a stronger electron acceptor than O<sub>2</sub> in terms of thermodynamics, a number of authors have shown that N<sub>2</sub>O respiration is energetically less efficient than aerobic respiration, resulting in lower biomass growth yields per substrate (Koike and Hattori 1975; Stouthamer et al. 1982; Beun et al. 2000). We cannot rule out the existence of a more energy-efficient N<sub>2</sub>O reduction process (Conthe et al. 2018a), considering the broad phylogenetic diversity of N<sub>2</sub>O reducers and our limited knowledge regarding non-denitrifying N<sub>2</sub>O reducers in particular. However, given the growth yields reported in literature, it

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00253-018-9247-3>) contains supplementary material, which is available to authorized users.

✉ Monica Conthe  
M.conthecalvo-24@tudelft.nl

<sup>1</sup> Department of Biotechnology, Delft University of Technology, Van der Maasweg 9, 2629 HZ Delft, The Netherlands

would make evolutionary sense for microorganisms to favor aerobic respiration over the respiration of N compounds to optimize energy conservation in the cell. Intriguingly, the physical mechanism directing electrons to O<sub>2</sub> preferentially over other N compounds, when both electron acceptors are available, remains unclear.

Regulatory systems on a transcriptional or post-transcriptional level have been shown to shut down denitrification in the presence of oxygen in a variety of organisms (Zumft 1997). For instance, the NosZ protein of *Paracoccus denitrificans* and *Pseudomonas stutzeri* is inhibited by O<sub>2</sub> in vitro (Coyle et al. 1985; Alefounder and Ferguson 1982), which could be a form of allosteric regulation in vivo. It has also been proposed that N<sub>2</sub>OR is—for reasons unknown—less competent than the cytochrome oxidases involved in respiration of O<sub>2</sub> in the “competition” for electrons in the ETC (Qu et al. 2015). Nevertheless, diverse studies have reported the occurrence of denitrification in the presence of O<sub>2</sub> (termed aerobic denitrification; Chen and Strous 2013 and references therein). Regarding N<sub>2</sub>O reduction more specifically, a significant degree of N<sub>2</sub>OR transcription and activity has been found under aerated conditions (Körner and Zumft 1989; Qu et al. 2015).

From a greenhouse gas mitigation point of view, it is interesting to study O<sub>2</sub> and N<sub>2</sub>O mixotrophy—or the capability of microorganisms to simultaneously respire O<sub>2</sub> and N<sub>2</sub>O—in order to understand how frequent oxic-anoxic shifts during nitrogen removal from wastewater, in space or time, may affect the N<sub>2</sub>O-reducing capacity of activated sludge. WWTP design and operation vary greatly, but universal questions to address are, e.g., (a) if N<sub>2</sub>OR activity can persist in aerated zones consuming nitrification-derived N<sub>2</sub>O potentially minimizing greenhouse gas emissions or (b) if, on the contrary, N<sub>2</sub>OR is relatively less active than the other NO<sub>x</sub> reductases in the presence of O<sub>2</sub>, leading to N<sub>2</sub>O accumulation in the aerobic-anoxic transition zones.

We explored O<sub>2</sub> versus N<sub>2</sub>O respiration in a continuous enrichment culture selected and grown with N<sub>2</sub>O as the sole electron acceptor and fully characterized—in terms of stoichiometry and community composition—in a previous study (Conthe et al. 2018b). The culture had been found to be composed of a relatively simple microbial community dominated by *Dechlorobacter*-like *Betaproteobacteria*. In this study, operation of the chemostat was continued and the N<sub>2</sub>O-limited steady-state conditions were intermittently interrupted to perform short-term batch experiments in situ, with varying concentrations of N<sub>2</sub>O, O<sub>2</sub>, or both N<sub>2</sub>O and O<sub>2</sub> simultaneously, to determine (i) whether O<sub>2</sub> is, in fact, preferentially consumed over N<sub>2</sub>O when both electron acceptors are available, (ii) under which O<sub>2</sub> concentrations (if any) N<sub>2</sub>O consumption can take place, and (iii) to begin to unravel the mechanism governing the electron flow in the ETC to O<sub>2</sub> or N<sub>2</sub>O.

## Materials and methods

### Chemostat operation

Following the work presented in Conthe et al. (2018b), a microbial enrichment using acetate as a carbon and energy source and exogenous N<sub>2</sub>O as the sole electron acceptor was maintained under N<sub>2</sub>O-limiting conditions in a continuous culture at 20 °C, pH 7, and a dilution rate of  $0.026 \pm 0.001 \text{ h}^{-1}$ . The reactor set-up, operation, sampling, and medium composition are described in detail in Conthe et al. (2018b, c). One hundred percent pure N<sub>2</sub>O gas diluted in Argon gas was fed to the chemostat at a total flow rate of 200 ml/min and the offgas from the reactor was recirculated at a rate of 700 ml/min, resulting in an incoming N<sub>2</sub>O concentration of roughly 0.30%. The stability of the culture in terms of conversion rates and microbial community composition was monitored by regular sampling of the broth and biomass and via online monitoring of the acid (1 M HCl) dosing (a proxy for acetate consumption in the system) and offgas composition.

### Batch experiments

The steady-state conditions of the culture were briefly interrupted on different operation days in order to perform batch experiments in situ and determine the maximum conversion rates of the enrichment under non-limiting conditions (Figure S1). The medium and effluent pumps were switched off and the gas supply rates of O<sub>2</sub> (from a bottle of pure O<sub>2</sub>) and/or N<sub>2</sub>O were modified to achieve different electron acceptor concentrations within the system in random steps. Two main types of batches were performed: (1) supplying a single electron acceptor—either N<sub>2</sub>O or O<sub>2</sub>—at different concentrations or (2) supplying N<sub>2</sub>O and O<sub>2</sub> simultaneously, keeping the N<sub>2</sub>O gas supply rate constant and varying that of O<sub>2</sub>. Additionally, we performed a batch test in which a constant O<sub>2</sub> gas supply rate was maintained while varying that of N<sub>2</sub>O as well as short batch tests with either NO<sub>3</sub><sup>−</sup> or NO<sub>2</sub><sup>−</sup> to assess the denitrifying capacity of the culture. Note that gas recirculation was maintained during the experiments, causing an apparent delay between the conversions in the chemostat and the offgas concentration values measured. To avoid acetate depletion, a concentrated solution of sodium acetate was added to the broth at the start of the experiments and the 1 M HCl solution used for pH control during continuous operation was replaced by 1 M acetic acid for the duration of the experiment. For the batch tests with NO<sub>3</sub><sup>−</sup> and NO<sub>2</sub><sup>−</sup>, these compounds were supplied as 1 M KNO<sub>3</sub> or 1 M KNO<sub>2</sub>.

### Analytical procedures

Samples from the reactor for analysis of acetate and NH<sub>4</sub><sup>+</sup> were immediately filtered after sampling (0.45-μm pore size

poly-vinylidene difluoride membrane, Merck Millipore, Carrigtohill, Ireland). Acetate was measured with a Chrompack CP 9001 gas chromatograph (Chrompack, Middelburg, The Netherlands) equipped with an HP Innowax column (Agilent Technologies, Santa Clara, CA, USA) and a flame ionization detector. Ammonium,  $\text{NO}_3^-$ , and  $\text{NO}_2^-$  concentrations were determined spectrophotometrically using cuvette test kits (Hach Lange, Düsseldorf, Germany). For the estimation of biomass concentration, the volatile suspended solids (VSS) concentration was determined by centrifuging 0.2 L of the enrichment, drying the pellet overnight at 105 °C, and then burning the pellet at 550 °C for 2 h to determine the ash content. Additionally, the optical density of the culture (at a wavelength of 660;  $\text{OD}_{660}$ ) was monitored. Concentrations of  $\text{N}_2\text{O}$ ,  $\text{N}_2$  and  $\text{CO}_2$ , Argon, and  $\text{O}_2$  in the headspace of the reactor were measured online via mass spectrometry (Prima BT, Thermo Scientific). The dissolved  $\text{O}_2$  concentration in the broth during the batch tests with  $\text{O}_2$  was measured with two types of oxygen sensors: a Clark electrode calibrated in the range of 0–20.8% and an optical oxygen probe calibrated in range 0–2% (Presens, Regensburg, Germany).

## Calculations

Elemental and electron balances during steady state were set up as described in Conthe et al. (2018a, b, c). During the batch tests, the conversion rates ( $r$ , in  $\text{mol h}^{-1}$ ) for  $\text{O}_2$  and  $\text{N}_2\text{O}$  were calculated from the measured ingoing and outgoing gas composition and the argon supply rate (see Figures S2–S6 and Tables S2–S6 for details). The average biomass concentration value for each experimental step was derived from the ammonium uptake rates (see for example Figure S4b) and used to calculate the corresponding biomass specific rates ( $q$ , in  $\text{mol CmolX}^{-1} \text{h}^{-1}$ ). A standard and constant biomass composition of  $\text{CH}_{1.8}\text{O}_{0.5}\text{N}_{0.2}$  (Roels 1980). The  $q_{\text{O}_2}$  and  $q_{\text{N}_2\text{O}}$  obtained for each step were plotted against the corresponding concentration of dissolved  $\text{O}_2$  or  $\text{N}_2\text{O}$  in the broth in order to determine the  $q_{\text{max}}$  and  $K_s$  of the enrichment for  $\text{O}_2$  and  $\text{N}_2\text{O}$ .

The concentration of dissolved  $\text{O}_2$  was obtained experimentally with the DO probes while the concentration of dissolved  $\text{N}_2\text{O}$  was estimated given a  $k_{\text{LaN}_2\text{O}}$  of  $180 \text{ h}^{-1}$ —obtained by scaling the experimentally derived  $k_{\text{LaO}_2}$  (Janssen and Warmoeskerken 1987) and deriving the corresponding  $K_{\text{La}}^{\text{broth}}$  and  $K_{\text{La}}^{\text{headspace}}$  assuming a  $t_{\text{broth}}$  of 6 s (1800 and  $50 \text{ h}^{-1}$ , respectively). A Monod model fitting the results was obtained by minimizing the sum of squared errors using the Microsoft Excel software.

The thermodynamic efficiency of metabolic growth using acetate as an electron donor and  $\text{O}_2$ ,  $\text{N}_2\text{O}$ , or  $\text{NO}_3^-$  as an electron acceptor can be interpreted by the Gibbs free energy ( $\Delta G_{01}$ ) dissipated per C mole of biomass growth or per electron-equivalent used for respiration. These values were calculated based on Kleerebezem and van Loosdrecht (2010) and using the thermodynamic values found in Thauer et al. (1977)—please refer to Table S7 for more details.

## DNA extraction and 454 amplicon sequencing of 16S rRNA gene

The taxa-based community composition of the enriched culture during the period of operation presented in this study was determined by 454 amplicon sequencing of the 16S rRNA gene following the procedure described in Conthe et al. (2018a, b, c) and the sequences are available at NCBI under BioProject accession number PRJNA413885.

## Results

### Continuous operation and microbial community composition of the $\text{N}_2\text{O}$ -reducing enrichment

A culture enriched from activated sludge using acetate as a carbon source and electron donor and exogenous  $\text{N}_2\text{O}$  as the sole electron acceptor was studied for a total period of 155 days (> 100 volume changes) in a chemostat under electron acceptor ( $\text{N}_2\text{O}$ ) limiting conditions (Figure S1). The start-

**Table 1** Average biomass-specific conversion rates during steady state and the batch experiments

	Compound biomass specific conversion rates ( $\text{mmol}/\text{mmol}_X \text{ h}^{-1}$ )			
	$q_{\text{N}_2\text{O}-\text{N}}$	$q_{\text{NO}_3-\text{N}}$ or $q_{\text{NO}_2-\text{N}}$	$q_{\text{N}_2-\text{N}}$	$q_{\text{Acetate}-\text{C}}$
Steady state	$-0.033 \pm 0.001^b$		$0.034 \pm 0.001^b$	$-0.017 \pm 0.001^b$
$\text{N}_2\text{O}$ batch	$-0.131 \pm 0.004^b$		$0.126 \pm 0.008^b$	$-0.067 \pm 0.009^c$
$\text{NO}_3^-$ batch		$-0.007 \pm 0.000^c$	$0.004 \pm 0.000^c$	$-0.003 \pm 0.000^c$
$\text{N}_2\text{O} + \text{NO}_2^-$ batch <sup>a</sup>	$-0.033 \pm 0.000^c$		$0.042 \pm 0.000^c$	

<sup>a</sup>  $\text{N}_2\text{O}$  gas supply was kept on during addition of 1 mM  $\text{KNO}_2^-$

<sup>b</sup> Standard deviation calculated from at least three independent measurements

<sup>c</sup> Standard deviation calculated by LINEST least squares method

**Table 2** Experimentally determined biomass yields per mole of electron donor or per mole of electron equivalents respired during growth with  $N_2O$ ,  $NO_3^-$ , and  $O_2$  as an electron acceptor and corresponding Gibbs free energy dissipation values based on these yields

Parameter	Units	Growth on electron acceptor			
		$N_2O^a$	$NO_3^-^b$	$O_2^c$	
$Y_{XS}$	Biomass yield on acetate	$Cmol_X/Cmol_{Ac}$	$0.36 \pm 0.03$	0.38	0.45
$Y_{Xe}$	Biomass yield on $e^-$ transported in catabolic process	$Cmol_X/mol_{e^-}$	$0.16 \pm 0.01$	0.15	0.19
$\Delta G_{MET}^{01}$	Metabolic energy change per mole donor <sup>d</sup>	$kJ/Cmol_X$	-1078	-620	-479
$\Delta G_{e\text{ CAT}}^{01}$	Metabolic energy change per electron transferred in catabolism	$kJ/mol_{e^-}$	-159	-96	-101

<sup>a</sup> Steady state data, this study

<sup>b</sup> Steady state data—no significant accumulation of intermediates (Conthe et al.; data unpublished)

<sup>c</sup> Batch experiment data in  $N_2O$  reducing enrichment, this study


up and characterization of the enrichment during the first 70 days of operation, in terms of conversion rates, stoichiometry, and microbial community composition, are described in Conthe et al. (2018b). During the subsequent period reported here, the conversion rates and corresponding biomass yields remained consistent with the previous period, characterized by steady-state growth on acetate oxidation coupled to  $N_2O$  reduction to  $N_2$  (Tables 1 and 2). Furthermore, 454 amplicon sequencing of the 16S rRNA gene of the microbial community confirmed the continued prevalence of a *Dechlorobacter*-like OTU (Figure S1), transiently co-occurring (around day

100) with two other closely related OTUs classified as *Azonexus* and uncultured *Rhodocyclaceae*.


### $O_2$ vs. $N_2O$ batch tests: affinity and yields

Batch experiments with varying supply rates of either  $N_2O$  or  $O_2$  were performed on days 106 and 132, respectively (Fig. 1). The maximum biomass specific conversion rates of  $N_2O$  ( $q_{N_2O}^{max}$ ) and acetate were identified by increasing the  $N_2O$  supply rate to non-limiting conditions. The  $q_{N_2O}^{max}$  values

**Fig. 1** Offgas data from the batch experiments with varying concentrations of **a**  $N_2O$ ; day 106, **b** and  $O_2$ ; day 132. For the experiment with  $O_2$ , the dissolved oxygen concentration (DO) was measured both with a Clark electrode (DO\_1) and an optical sensor (DO\_2). The affinity of the culture for  $N_2O$  and  $O_2$  was determined from these experiments (see Fig. 5). The asterisk mark time points at which acetate had been depleted and was added to the culture



**Fig. 2** Biomass specific transfer rates of electron equivalents ( $q_e$ ) as a function of the electron acceptor concentration (either  $N_2O$ , in black, or  $O_2$ , in gray), along with the fitting Monod model (with the corresponding  $q_e^{\max}$  and  $K_s$  parameters). The inset is an enlargement of the graph at low  $O_2/N_2O$  concentrations. The rates presented were obtained from the experiments shown in Fig. 2



identified were roughly fourfold higher than the actual biomass specific conversion rates during steady state (Table 1). When exposed to varying concentrations of  $O_2$ , the culture was able to switch to aerobic respiration in the order of seconds. The maximum  $O_2$  reducing capacity ( $q_{O_2}^{\max}$ ) was comparable to  $N_2O$  respiration when expressed per mole electron accepted.  $NO_3^-$  and  $NO_2^-$  reducing capacities were much lower compared to  $N_2O$  or  $O_2$  (< 15% of the maximum  $N_2O$  or  $O_2$  reduction rate; Table 1).


Plotting the biomass-specific electron transfer rate ( $q_e^-$ ) at different dissolved  $O_2$  (DO) or  $N_2O$  concentrations, we could determine the apparent  $K_s$  for  $O_2$  or  $N_2O$  by fitting a Monod model to the data (Fig. 2). Given the confidence intervals, the absolute value for this parameter could not be identified accurately, but the results demonstrate clearly that the  $K_s$  value for  $O_2$  is 1 or 2 orders of magnitude smaller compared to  $K_s-N_2O$ . The maximum biomass-specific conversion rate of  $O_2$  ( $q_{O_2}^{\max}$ )

was roughly two times lower than that of  $N_2O$  ( $q_{N_2O}^{\max}$ ) per mole of electron acceptor but the conversion rates expressed as electron equivalents ( $q_e^{\max}$ ) were comparable for both processes, since double the electrons are taken up during the reduction of  $O_2$  to  $H_2O$  compared to  $N_2O$  to  $N_2$ .

The biomass yields per mole of electron donor (determined from the steady-state growth on  $N_2O$  in the chemostat, and from the batch experiments with  $O_2$  as the sole electron acceptor) are presented in Table 2.

### Simultaneous $O_2$ and $N_2O$ batch tests


Batch experiments with excess  $N_2O$  and varying concentrations of  $O_2$ , supplied simultaneously, were performed on days 110 and 155 (Figs. 3 and 4). The maximum electron transfer rate ( $q_e^{\max}$ )—combining the electron transfer capacities of  $N_2O$  and  $O_2$ —summed up to a value comparable with the



**Fig. 3** Offgas data from the batch experiments with excess  $N_2O$  and varying concentrations of  $O_2$  on **a** day 125 and **b** day 155. The dissolved oxygen concentration (DO) was measured both with a Clark electrode (DO\_1) and an optical sensor (DO\_2). The biomass specific electron transfer rates to either  $N_2O$  or  $O_2$  during these experiments are shown in Fig. 4. The asterisk marking the last two steps of the batch experiment on day 155 indicates the culture ran out of  $NH_4^+$  for growth,

and thus the rates during these steps was not considered. **c** Offgas data of batch experiment with excess  $O_2$  and varying concentrations of  $N_2O$  on day 113. Detailed data from these experiments can be found in the Supplementary Materials—Tables xxx–xxx and Figures xxx to xxx. **d** Detailed view of one of the steps from the batch experiment depicted in (**a**) showing the simultaneous consumption of  $O_2$  and  $N_2O$ , and subsequent production of  $N_2$


**Fig. 4** Biomass specific transfers rate of electron equivalents ( $q_e$ ) (a) to  $O_2$  and b  $N_2O$  and c percentage of total electrons being shuttled to  $N_2O$  vs.  $O_2$  at varying  $O_2$  concentrations during the batch tests on day 125 (in black) and day 155 (in gray). The Monod model of  $O_2$  consumption in the absence of  $N_2O$  (shown in Fig. 3) is included in (a) for comparison. The inset in (c) is an enlargement of the graph at low  $O_2$  concentrations



$q_e^{\max}$  found during the  $N_2O$ - or  $O_2$ -only experiments.  $N_2O$  reduction to  $N_2$  co-occurred with aerobic respiration only at relatively low concentrations of  $O_2$  (Fig. 3d). The experiments performed on days 110 and 155 differed regarding the  $O_2$  concentration range at which  $N_2O$  reduction could co-occur (roughly  $< 4$  and  $< 1.5 \mu\text{M}$   $O_2$  on days 110 and 155, respectively) but, nevertheless,  $N_2O$  reduction in the presence of  $O_2$  contributed to no more than a small fraction of the total electron acceptor capacity (generally  $< 20\%$  of  $q_e^{\text{tot}}$ ; Fig. 4). An additional batch experiment on day 113, with a constant supply of  $O_2$  and a varying supply of  $N_2O$ , also showed that  $N_2O$  reduction was undetectable in the presence of relatively high concentrations of  $O_2$  ( $\approx 5 \mu\text{M}$ ; Fig. 3c).

## Discussion

Aerobic respiration was distinctly favored over  $N_2O$  respiration in the enrichment despite the fact that the culture had been operated for an extensive number of generations with  $N_2O$  as only electron acceptor. Upon a sudden change in supply from  $N_2O$  to  $O_2$ , the culture readily switched to  $O_2$  respiration and, when both electron acceptors were available,  $N_2O$  reduction was only observed at relatively low concentrations of  $O_2$  ( $< 4 \mu\text{M} = 0.13 \text{ mg } O_2/\text{L}$ ). Under conditions of electron acceptor excess ( $N_2O$  and/or  $O_2$ ), growth in the system was likely limited by the electron supply rate to the electron transport chain (see Fig. 5) and not by the capacity of  $N_2O$ OR or  $O_2$  reductases.



**Fig. 5** Simplified representation of the proportional distribution of electrons ( $e^-$ ) in the electron transport chain (ETC) during batch tests with only  $N_2O$  (open arrow) versus batch tests with the simultaneous addition of  $O_2$  and  $N_2O$  (black arrows) showing that there is a preferential shuttling of electrons to  $O_2R$  than to  $N_2OR$ . This simplified schematic is based on the assumptions that (i) both enzymes share a common electron pool (/quinone pool) and (ii) that all cells have a similar electron distribution among terminal reductases (whereas it would be possible for the majority of cells to switch fully to aerobic respiration, and a small fraction to continue respiring  $N_2O$ )

This was inferred from the fact that the maximum electron acceptor capacity of the culture was comparable for  $N_2O$  and  $O_2$  respiration (i.e.,  $q_{e-N_2O}^{max} \approx q_{e-O_2}^{max}$ ), and could be due to kinetic limitations in acetate uptake, acetate oxidation in the citric acid cycle, or in some shared component of the ETC itself.

The overall electron transfer capacity during the simultaneous respiration of  $N_2O$  and  $O_2$  (i.e.,  $q_{e-TOT}^{max}$ ) was comparable to  $q_{e-N_2O}^{max}$  or  $q_{e-O_2}^{max}$ . This suggests that “aerobic  $N_2O$  respiration” (by analogy to aerobic denitrification) generally occurs if the electron supply rate to the ETC exceeds the electron accepting capacity of the  $O_2$  reductases. In other words,  $N_2O$  respiration complements aerobic respiration primarily when  $O_2$  is limiting. Nonetheless, our results indicate that, under  $O_2$ -limiting conditions,  $N_2O$  reducers can use  $O_2$  and  $N_2O$  mixotrophically as proposed by Chen and Strous 2013 (Fig. 5). We cannot exclude heterogeneity in electron acceptor use within the population in our bioreactor leading for example to most of the culture respiring  $O_2$  and a side population reducing  $N_2O$ . Under the microscope, we did not observe formation of aggregates or biofilms which could create anoxic niches in spite of the  $O_2$  supply (data not shown), yet oxygen gradients and anoxic microzones could still form around suspended cells if  $O_2$  diffusion rate is slower than the respiration rate. Nevertheless, with the strong sparging and mixing conditions imposed on the culture, we would expect that most cells would be exposed to comparable environmental conditions.

The  $K_s$  values of the enrichment culture were in the same range as the  $K_m$  values reported for purified  $N_2OR$  and different  $O_2$  reductases in literature, i.e., in the  $\mu M$  range for  $N_2O$  and  $nM$  range for  $O_2$  (Pouvreau et al. 2008 and references therein, Yoon et al. 2016). The relatively high  $K_{S,N_2O}$  (two orders of magnitude higher than for  $O_2$ ) is noteworthy in a culture presumably well-adapted to  $N_2O$ -limiting conditions. Also the observation that, even after a prolonged absence of  $O_2$  in the environment, the

cellular machinery specific for aerobic respiration (i.e., cytochrome oxidases) was constitutively present (in contrast to  $NO_3^-$  and  $NO_2^-$  reductases). According to these results, the preferential use of  $O_2$  over  $N_2O$  in natural systems could be attributed to a difference in affinity ( $\mu_{max}/K_s$ ) for  $O_2$  and  $N_2O$ .

With regard to efficiency of  $N_2O$  respiration versus  $O_2$  respiration, our chemostat enrichment cultures corroborate studies in literature (Koike and Hattori 1975; Stouthamer et al. 1982; Beun et al. 2000) and predictions based on our knowledge of the ETC in model denitrifiers (Chen and Strous 2013): with biomass yields per mole of acetate during growth with  $N_2O$  (or  $NO_3^-$ ) roughly 1/3 lower than yields during  $O_2$  respiration (Table 2). The relatively low growth yields on  $N_2O$  imply that  $N_2O$  reduction to  $N_2$  is, thermodynamically, a very inefficient process with high energy dissipation. Thus, ensuring the maximization of energy conservation during microbial growth may be the evolutionary driver behind the preferential flow of electrons to  $O_2$  over  $N_2O$ .

We cannot provide a conclusive answer regarding which cellular mechanism governs the preferential use of  $O_2$  in the presence of excess  $N_2O$  observed. However, the instantaneous switch from  $N_2O$  to  $O_2$  respiration suggests that the preference for  $O_2$  over  $N_2O$  is regulated at the metabolome level and is independent from transcriptional regulation, e.g., by control of enzyme activity, like allosteric inhibition of  $N_2OR$ , or simply a higher affinity of  $O_2$  reductases for the electrons coming from a common quinone pool.

Translated to the environmental conditions in a WWT plant, the results from this study suggest that oxic-anoxic transitions are unlikely to result in  $N_2O$  emissions associated to denitrification as a result of  $N_2OR$  inhibition by  $O_2$  since the enrichment culture readily switched back and forth between  $O_2$  and  $N_2O$  respiration. This implies that (a) either  $N_2OR$  is not directly inhibited by  $O_2$  in vivo or (b) inhibition is readily reversible once  $O_2$  is depleted.

On the other hand, the fact that aerobic respiration is so strongly favored over  $N_2O$  respiration would make it a challenge to exploit the  $N_2O$  sink capacity of activated sludge in the aerated/nitrification zones of WWT plants. The range in which significant  $N_2O$  consumption co-occurred with  $O_2$  consumption in our experiments was narrow: roughly up to 1.5–4  $\mu M$   $O_2$ , i.e., 0.05–0.13 mg  $O_2/L$ , presumably below common DO values in the aerated tanks of WWTP (Tchobanoglous and Burton 2002). The very high affinity for oxygen minimizes the range of dissolved oxygen concentrations in which  $O_2$  and  $N_2O$  respiration could occur in parallel. However, a beneficial difference in full-scale systems compared to our enrichment, in terms of avoiding  $N_2O$  accumulation, may be that mass transfer limitation induced oxygen limitation within the activated sludge flocs provide anoxic zones, prone to  $N_2O$  reduction, even when  $O_2$  is present in the bulk liquid (Picioreanu et al. 2016). This, together with the fact that  $N_2O$  is much more soluble than  $O_2$ , could perhaps be exploited to enhance the  $N_2O$  sink capacity of activated sludge.

**Acknowledgements** The authors would like to thank Gijs Kuenen for his comments on the manuscript and Mitchell Geleijnse and Ben Abbas for their great help with the molecular analysis of microbial community composition.

**Funding** This work was funded by the European Commission (Marie Curie ITN NORA, FP7-316472).

## Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

**Ethical approval** This article does not contain any studies with human participants or animals performed by any of the authors.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Alefunder PR, Ferguson SJ (1982) Electron transport-linked nitrous oxide synthesis and reduction by *Paracoccusdenitrificans* monitored with an electrode. *Biochem Biophys Res Commun* 104 (3):1149–1155
- Beun JJ, Verhoef EV, Van Loosdrecht MCM, Heijnen JJ (2000) Stoichiometry and kinetics of poly- $\beta$ -hydroxybutyrate metabolism under denitrifying conditions in activated sludge cultures. *Biotechnol Bioeng* 68:496–507
- Chen J, Strous M (2013) Denitrification and aerobic respiration, hybrid electron transport chains and co-evolution. *Biochim Biophys Acta Bioenerg* 1827:136–144
- Conthe M, Kuenen JG, Kleerebezem R, van Loosdrecht MCM (2018a) Exploring microbial  $N_2O$  reduction: a continuous enrichment in nitrogen free medium. *Environ Microbiol Rep* 10:102–107. <https://doi.org/10.1111/1758-2229.12615>
- Conthe M, Wittorf L, Kuenen JG, Kleerebezem R, Hallin S, van Loosdrecht MCM (2018b) Growth yield and selection of nosZ clade II-types in a continuous enrichment culture of  $N_2O$  respiring bacteria. *Environ Microbiol Rep* 10:239–244. <https://doi.org/10.1111/1758-2229.12630>
- Conthe M, Wittorf L, Kuenen JG, Kleerebezem R, van Loosdrecht MCM, Hallin S (2018c) Life on  $N_2O$ : deciphering the ecophysiology of  $N_2O$  respiring bacterial communities in a continuous culture. *ISME J*. <https://doi.org/10.1038/s41396-018-0063-7>
- Coyle CL, Zumft WG, Kroneck PMH, Körner H, Jakob W (1985) Nitrous oxide reductase from denitrifying: *Pseudomonas perfectomarina* purification and properties of a novel multicopper enzyme. *Eur J Biochem* 153:459–467
- Hallin S, Philippot L, Löffler FE, Sanford RA, Jones CM (2018) Genomics and ecology of novel  $N_2O$ -reducing microorganisms. *Trends Microbiol* 26:43–55
- Janssen LPBM, Warmoeskerken MMCG (2006) Transport phenomena data companion, Third edit edn. VSSD, Delft
- Kleerebezem R, van Loosdrecht MCM (2010) A generalized method for thermodynamic state analysis of environmental systems. *Crit Rev Env Sci Technol* 40(1):1–54
- Koike I, Hattori A (1975) Energy yield of denitrification: an estimate from growth yield in continuous cultures of *Pseudomonas denitrificans* under nitrate-, nitrite- and oxide-limited conditions. *J Gen Microbiol* 88:11–19
- Körner H, Zumft WG (1989) Expression of denitrification enzymes in response to the dissolved oxygen level and respiratory substrate in continuous culture of *Pseudomonas stutzeri*. *Appl Environ Microbiol* 55:1670–1676
- Picioreanu C, Erez JP, Van Loosdrecht MCM (2016) Impact of cell cluster size on apparent half-saturation coefficients for oxygen in nitrifying sludge and biofilms. <https://doi.org/10.1016/j.watres.2016.10.017>
- Pouvreau LAM, Strampraad MJF, Van Berloo S, Kattenberg JH, de Vries S (2008)  $NO$ ,  $N_2O$ , and  $O_2$  reaction kinetics: scope and limitations of the Clark electrode. *Methods Enzymol* 436:97–112
- Qu Z, Bakken LR, Molstad L, Frostegård Å, Bergaust L (2015) Transcriptional and metabolic regulation of denitrification in *Paracoccus denitrificans* allows low but significant activity of nitrous oxide reductase under oxic conditions. *Environ Microbiol* 18: 2951–2963. <https://doi.org/10.1111/1462-2920.13128>
- Roels JA (1980) Simple model for the energetics of growth on substrates with different degrees of reduction. *Biotechnol Bioeng* 22:33–53
- Shapleigh JP (2013) Denitrifying Prokaryotes. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (eds) *The prokaryotes: prokaryotic physiology and biochemistry*. Springer Berlin Heidelberg, Berlin, pp 405–425
- Stouthamer AH, Boogerd FC, van Verseveld HW (1982) The bioenergetics of denitrification. *Antonie Van Leeuwenhoek* 48:545–553
- Thauer RK, Jungermann K, Decker K (1977) Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* 41:100–180
- Tchobanoglous G, Burton F, Stensel HD (2006) *Wastewater engineering—treatment and reuse*. McGraw-Hill, New York
- Thomson AJ, Giannopoulos G, Pretty J, Baggs EM, Richardson DJ (2012) Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philos Trans R Soc Lond Ser B Biol Sci* 367: 1157–1168
- van Spanning RJM, Richardson DJ, Ferguson SJ (2007) Introduction to the biochemistry and molecular biology of Denitrification. *Biology of the Nitrogen Cycle* 3–20. <https://doi.org/10.1016/B978-044452857-5.50002-3>
- Yoon S, Nissen S, Park D, Sanford RA, Löffler FE (2016) Nitrous oxide reduction kinetics distinguish bacteria harboring clade I NosZ from those harboring clade II NosZ. *Appl Environ Microbiol* 82:3793–3800
- Zumft WG (1997) Cell biology and molecular basis of cell biology and molecular basis of Denitrification. *Microbiology* 61(4):533