

A chronic thigh mass in a 69-year-old man

Simon A. S. Beggs · Awatif Al-Nafussi ·
C. Michael Lambert · Daniel Porter · J. T. Patton

Published online: 18 June 2010
© ISS 2010

Question

A 69-year-old man presented with a solitary, painless swelling on the lateral aspect of his left thigh that had been present, unchanged for 10 years. His medical history included chronic obstructive pulmonary disease, hypertension and bilateral knee osteoarthritis.



Fig. 1 Magnetic resonance imaging of the thigh, including **a** axial T1-weighted, **b** coronal T2-weighted and **c** axial fat-suppressed T1-weighted images after gadolinium administration, was followed by ultrasound-guided core biopsy

The diagnosis can be found at doi: 10.1007/s00256-010-0973-6.

S. A. S. Beggs (✉)
Edinburgh University College
of Medicine and Veterinary Medicine,
The Chancellor's Building 2nd Floor, 49 Little France Crescent,
Edinburgh EH16 4SB, UK
e-mail: beggs1@googlemail.com

A. Al-Nafussi
Department of Pathology, Royal Infirmary,
Edinburgh EH16 4SA, UK

C. M. Lambert
Department of Rheumatology, Western General Hospital,
Edinburgh EH4 2XU, UK

D. Porter · J. T. Patton
Department of Orthopaedics, Royal Infirmary,
Edinburgh EH16 4SA, UK