Multiple bilateral forearm nodules

Robert W. Morris¹ • Youssef Al Hmada² • Joseph R. Chaudry¹ • Kathryn M. Nutter¹ • Chris D. Reed¹

Published online: 3 August 2019
© ISS 2019

44 year-old female with several month history of firm, painful nodules involving the bilateral forearms. MRI with contrast of the left (Fig. 1) and right (Fig. 2) forearms was performed. Excision of a painful right forearm nodule was also performed (Fig. 3).

The diagnosis can be found at <https://doi.org/10.1007/s00256-019-03296-9>

✉ Robert W. Morris
rwmorris@umc.edu

¹ Department of Radiology, University of Mississippi Medical Center, 2500 North State Street, Jackson, MS 39216, USA

² Department of Pathology and Laboratory Medicine, University of Mississippi Medical Center, 2500 North State Street, Jackson, MS 39216, USA

Fig. 1 **a** Axial T1, **b** axial PD fat sat and **c** axial T1 fat sat with contrast of the left forearm

Fig. 2 **a** Coronal T1, **b** coronal PD fat sat and **c** coronal T1 fat sat with contrast of the right forearm

Fig. 3 Hematoxylin and eosin stained sections of the excised right forearm nodule at **a** 40x and **b** 100x magnification

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.